

		I	905.			19	08.	18		19	10.	
Star.	E.		W.		E.		W.		E.		W.	
	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.
ζ Octantis γ Chamæleontis Lacaille 4510 η Octantis β Chamæleontis ι Octantis κ Octantis	+0'19 +0'23 +1'11 +0'54	3°4 3°9 2°7 0°7 3°3 3°7	-0.86 +0.68 +0.32 -0.40 -0.55	2.4 2.8 2.9 3.8 2.1	$ \begin{array}{r} -0^{2}29 \\ -0^{2}23 \\ +0^{0}4 \\ +0^{4}5 \\ -0^{0}6 \\ +0^{4}5 \\ \dots\end{array} $	2.7 1.9 1.3 1.9 2.0 2.2	-0.67 +0.45 -0.72  -0.48 -0.60	2·2 0·7 2·2 0.8 2·4	+0.53	0.2	"	0.7
$\begin{array}{l} \theta \text{ Apodis} \\ a \text{ Apodis} \\ \zeta \text{ Apodis} \\ \text{Lacaille } 6077 \\ \rho \text{ Octantis} \\ \delta^1 \text{ Apodis} \end{array}$	+0 <sup>.85</sup>	2.0	- 0°44	0.8	-0.53  +0.85 -0.08 +0.47	0.8 1.3 0.8 1.3	- 1.07 - 1.15 - 0.20 - 1.06 - 0.77 + 0.07	1.7 1.2 1.5 1.3 2.9 2.0	+0.19		+ 0.71	0.2
γ Apodis Lacaille 6545 β Apodis χ Octantis Lacaille 8094 Lacaille 8257	-0°18  	1.5	+0.24 0.09	1.2 0.9	+0.74 +0.22 +0.55 +0.44 -0.54 +0.47	2.0 2.5 2.0 2.3 2.7 2.9	-1.38 - 0.42 - 0.70 - 0.26 - 0.89 + 0.61	0.8 1.6 1.7 1.5 1.2 2.5	-0.66  +0.41	1.0	+0.39	1'2
$\mu^{1} \text{ Octantis}$ $\mu \text{ Octantis}$ $\nu \text{ Octantis}$ $\nu (C) \text{ Octantis}$ $\tau \text{ Octantis}$ Lacaille 9494	-0'12 +0'13	3 <sup>.9</sup> 3 <sup>.2</sup> 0 <sup>.7</sup>	+0°14 +0°04 -0°09	3.0 2.7 1.3	+0.11 -0.02 -0.78  -0.27 -0.25	1.6 2.2 1.3 2.2 2.2	-0.65 -0.24 -0.09 -0.43 -0.59 -0.74	1.6 2.2 1.9 3.5 1.2 1.2	+ 1.68	0.2		
$\theta$ Octantis		2.0	+0.10	2.0	+0.05	0.2			+0.15	0.2	-0.20	1.0

TABLE XIV.—continued.

whence we derive in the mean

 $\Delta\delta$  (above - below).

Roberts I	Clamp E.	Weight.	Clamp W.	Weight.
1905	+0.17	45	$ \begin{array}{r} -0^{*1}5 \\ -0^{*4}9 \\ -0^{*2}7 \end{array} $	40
1908	+0.04	65		62
1910	+0.47	10		8

The differences E - W give in the mean the value

which corresponds very closely with the value of  $2B_{II}$  at the pole, as previously determined.

Applying the corrections  $-2(A \mp B_{II})$  respectively to results from Clamp E and Clamp W, we obtain the following values :---

	Δ	δ (abo	ve-below).	
			Clamp E.	Clamp W.
1905			+0.07	+0.23
1908			-0.06	-0.11
1910			+0.32	+0.11

and the corresponding corrections to the declinations on account of the combined effects of latitude and flexure :---

		Clamp E.	Clamp W.
1905		-0'04	-0'12
1908		+0.03	+0.06
1910		-0.18	-0.06

The parts of these quantities due to flexure alone are respectively

1905		-0.27
1908		-0.14
1910		-0.02

whence the derived values for the latitude correction referred to the mean system are

	1		Clamp E.	Clamp W.
1905			+0.23	+0.12
1908			+0.12	+0'20
1910		,	-0.13	-0.01

Collecting the various determinations, we find as the latitude correction referred to the homogeneous system,  $[\frac{1}{2}(I + II) : \frac{1}{2}(E + W)]$ .

Period of Observations.	Position.	Clamp.	$\Delta \phi.$	Weight.
1906-10 " 1905 " 1908 " 1910 " 1911	I. I. II. II. II. II. II. II. II. II. I	E W E W E W E and W	$ \begin{array}{r} + & 0^{\circ}12 \\ + & 0^{\circ}08 \\ + & 0^{\circ}23 \\ + & 0^{\circ}15 \\ + & 0^{\circ}17 \\ + & 0^{\circ}20 \\ - & 0^{\circ}13 \\ - & 0^{\circ}01 \\ + & 0^{\circ}17 \end{array} $	217 192 45 • 65 62 10 8 309


The weighted mean of these results gives as the definitive latitude correction applicable to the mean system of the Ledgers


 $\Delta \phi = + 0^{"} \cdot 14 \pm 0^{"} \cdot 012.$ 

C. F. C., 1900.

xxxiii

5







# FUNDAMENTAL CATALOGUE

ÓF∙

## 1293 STARS

FOR THE EQUINOX

## 1900

FROM OBSERVATIONS MADE AT THE

## ROYAL OBSERVATORY, CAPE OF GOOD HOPE,

DURING THE YEARS

## 1905 - 1911:

UNDER THE DIRECTION OF

SIR DAVID GILL, K.C.B., LL.D., D.Sc., F.R.S., Hon. F.R.S.Ed., &c., FORMERLY HIS MAJESTY'S ASTRONOMER,

AND

S. S. HOUGH, M.A., F.R.S., HIS MAJESTY'S ASTRONOMER AT THE CAPE.

Epe of Good Hope . Royal observatory

Published by Order of the Lords Commissioners of the Admiralty, in obedience to His Majesty's Command.

#### EDINBURGH:

PRINTED UNDER THE AUTHORITY OF HIS MAJESTY'S STATIONERY OFFICE By NEILL & CO., Limited, Bellevue.

To be purchased, either directly or through any Bookseller, from WYMAN AND SONS, LIMITED, 29 BREAMS BUILDINGS, FETTER LANE, LONDON, E.C., and 54 ST MARY STREET, CARDIFF; or II.M. STATIONERY OFFICE (SCOTTISH BRANCH), 23 FORTH STREET, EDINBURGH; or E. PONSONBY, LIMITED, 116 GRAFTON STREET, DUBLIN; or from the Agencies in the British Colonies and Dependencies, the United States of America and other Foreign Countries of T. FISHER UNWIN, LIMITED, LONDON, W.C.

1915.

[Crown Copyright Reserved.] Price Five Shillings.

6B6 C35

Astron. Dept.

thers of Calfornaal

gift of art. Soc. of Pacific


ASTRONOMY DEPT.

## CONTENTS.

						PAGE
I. REVISION OF CLOCK STAR SYSTEM-						
System used for the Ledgers 1905-11 .						v
Comparison of I. E., I. W., II. E., II. W.			•			ix
Comparison with Newcomb's Catalogue						x
Daylight Observations of Clock Stars .		•		0.C	•	xi
Comparison of Day and Night Observations	•		•	•		xii
II. REVISION OF DECLINATION SYSTEM-						
Comparison of Observations Clamp E. and W.	. 80					xxvi
Comparison of Observations Positions I. and II.						xxvii
Flexure Observations				•		xxviii
Reduction to uniform system						xxviii
Latitude and flexure corrections			1.45		• 1 1	xxix
Systematic corrections to the Declinations				•	•	xxxiv
III. FORMATION OF DEFINITIVE CATALOGUE PLACES-						
Corrections applied to Ledger places .						XXXV
Weights used						XXXV
Explanation of separate columns of the Catalogue.						XXXV
IV. FURTHER CORRECTIONS TO CATALOGUE RIGHT ASCENSIONS-						
Examination of Pivot Errors						xl
Correction to Equinox						xl
V. FURTHER CORRECTIONS TO DECLINATIONS-						
					-	<sup>1</sup> xliii
Latitude and Refraction .						
m- Contraction in		-				1-27
THE CATALOGUE						

701156

Wt. 21906/121 & 163-600-8/15 -N. & Co., Ltd Gp. XV.



### INTRODUCTION.

THIS Catalogue is based on meridian observations of stars made with the new reversible transit circle from the year 1905, when the instrument was first brought into regular use, to the end of the year 1911. Details of the observations, together with a full account of the methods of reduction, will be published in the volumes of *Cape Meridian Observations* covering the same period which are now being passed through the press. A full description of the instrument itself is contained in the *History and Description of the Cape Observatory*, to which reference may be made for detailed particulars. It is thus only necessary here to give an account of the processes employed for the formation of the Catalogue subsequently to the collection of separate results contained in the ledgers.

#### I.—REVISION OF CLOCK-STAR SYSTEM.

The entries in the ledgers, as contained in the *Cape Meridian Observations*, depend on Clock Errors derived with Newcomb's places for the standard clock stars. The observed Right Ascensions of the clock stars themselves were only retained as determinations and transferred to the ledgers in cases where at least five such were observed within a watch, which generally did not exceed four hours in duration. Thus, though the individual star places obtained by combining the separate results will not accurately conform with those of Newcomb's Catalogue, it may be anticipated that the combination will reproduce in entirety any systematic errors of Right Ascension of Newcomb's Catalogue dependent on the Right Ascension itself, except such as involve fluctuations contained within narrow limits of Right Ascension. The latter will be to a large extent smoothed out in the process of combination.

Observations have been made in four different conditions of the instrument, distinguished as I. E., I. W., II. E., II. W. The symbols I. II. refer to the relative positions of the object glass and eye-end, I. denoting that the object glass is adjacent to the reading 0° on the fixed circle, and II. that it is adjacent to the reading 180°. The symbols E. and W. (East and West) refer to the position of the Clamp. Observations in position I. were made in the years 1906, 1907, 1909, 1910, 1911, and those in position II. in the years 1905, 1908, 1910.

The entries in the ledgers were all obtained by referring the results of the separate observations to the epoch and equinox 1900 with Newcomb's proper motions. The means of the derived right ascensions of the clock stars dependent on the four different conditions of the instrument, together with their combination derived by taking the simple mean of the four without weighting, are contained in the following Table :---

							1								
Star.	Dec.	R	light Asce	onsion 190	0.01		Corr. to New-	Star.	Dec.	R	light Asce	nsion 190	0'0.		Corr. to New-
Utat.	1900.	I. E.	I. W.	II. E.	II. W.	Mean.	comb.	Stal.	1900.	I. E.	I. W.	II. E.	II. W.	Mean.	comb.
γ Pegasi	+14 38	h m s o 8 5.116	8 5°112	s 5.110	s 5.102	s 5.111	\$ - '024	7 Arietis	+20 47	h m s 3 15 27 · 151	s 27°128	s 27°090	8 27°110	s 27°120	s - • 019
ı Ceti		14 19.970	19.990	19.973	19.988	19.980	008	o Tanri		19 25.855	25.830	25.848	25.853		+ .003
44 Piscium		20 16.578	16.550	16.565	16.547	16.560	016	٤ Tauri		21 44'917	44.903	44'916	44.910	44.912	
12 Ceti		24 56.118	56.121	56.114	56.130	56.121	010	f Tauri		25 21.074	21.036	21.048	21'042	21.020	Product State
13 Ceti		30 6.034	6.033	6.054	6.070	6.048	+ .015	e Eridani		28 13.106	13.120	13.124	13.138	5	+ .011
Lacaille 147	-25 19	32 12.551	12'551	12.526	12.525	12.538	+ .067	τ <sup>5</sup> Eridani		29 22.232	22.210	22.233	22.273		+ .053
ß Ceti	-18 32	38 34.226	34.228	34.222	34.218	34.224	- '002	δ Eridani		38 27.434	27.460	27.420	27.430		024
(Andromedæ	+23 43	42 2.178	2.180	2.160	2.142	2.165	030	η Tauri		41 32.301	32.315	32.288	32.318		
δ Piscium	1	43 29.614	29.590	29.610	29.578	29.598	010	τ <sup>6</sup> Eridani		42 32.728	32.704	32.738	32.753		+ .004
20 Ceti		47 53.835	53.818	53.811	53.813		+ .031	γ Eridani		53 21.808	21.811	21.822	21.833		002
e Piscium	+ 7 21	57 45.130	45.154	45.146	45.122	45.138	011	λ Tanri		55 8.354	8.347	8.320	8.346	8.342	
η Ceti	-10 43	I 3 33.580	33.563	33.546	33.565	33.564	-0.001	v Tauri		57 50'180	50.163	50.182	50.143	50.167	•000
( Piscium pr	+7 3	8 30.342	30.336	30.330	30.387	30.349	016	A Tauri		58 46.883	46.930	46.908	46.886	· ·	-0.018
θ Ceti	- 8 42	19 1.498	1.482	1.477	1.202	1.490	+ .006	43 Tauri		4 3 20.364	20.342	20.330	20.328	20'341	009
η Piscium	+14 50	26 7.865	7.860	7.848	7.860	7.858	+ .000	o <sup>2</sup> Eridani		10 40.158	40'162	40'185	40.165		+ .008
v Piscium	+ 4 59	36 13.580	13. 567	13.567	13.563	13.569	023	γ Tauri		14 6.072	6.005	6.095	6.078		008
τ Ceti		39 25.340	25.328	25.345	25.341	25.339	+ .012	δ Tauri		17 10.006	10.012	10.000	9.993	~	+ .002
o Piscinni	+ 8 39	40 6.720	6.707	6.716	6.730	6.718	000	e Tauri		22 46.563	46.558	46.562	46.550	46.558	029
Ç Ceti	-10 50	46 31.465	31.460	31.462	31.470	31.464	001	v Eridani		31 19.308	19.310	19:339	19:330		
E Piscium	+ 2 42	48 22.685	22.670	22.697	22.653	22.676	+ .008	53 Eridani		33 36.040	36.033	36.023	36.062	36.047	
& Arietis	+20 19	49 6.846	6.858	6.855	6.850	6.852	+ .016	<i>τ</i> Tauri	-	36 14 529	14.217	14.493	14.492		- •021
v Ceti	-21 34	55 17.598	17.608	17.612	17.620	17.010	+ .031	μ Eridani		40 30.112	30.110	30.125	30.110	30'117	1
a Arietis	+22 59	2 1 32.044	32.050	32.043	32.057	32.049	003	$\pi^3$ Orionis		44 24 .684	24.682	24.685	24.628	24.670	and the second second
ξ1 Ceti	+ 8 23	7 41 . 923	41.906	41.914	41.930	41.918	+ .003	π <sup>5</sup> Orionis		49 2.519	2.233	2.524	2.543	2.530	
67 Ceti	- 6 53	11 59.710	59.736	59.725	59.733	59.726	+ .028	. Tauri		57 7.057	7.035	7.042	7.038		
o Ceti	- 3 26	14 17.637	17.624	17.653	17.650	17.641	- '021	e Leporis		5 1 13.647	13.656	13.650	13.660	13.653	
ξ <sup>2</sup> Ceti	+ 8 1	22 50.450	50.458	50.455	50.475	50.460	- ·001	ß Eridani	-	2 55.988	55.996	55.998	56.028	56.003	013
v Ceti	+ 5 9	30 37.548	37.542	37.505	37.530	37.531	+ .020	µ Leporis		8 26.354	26.371	26.325	26.343	26.348	010
v Arietis		33 8.123	8.173	8.145	8.154	8.149	016.	o Orionis		16 39.433	39.421	39.433	39.441	39.432	+ .027
δ Ceti	- 0 6	34 21.345	21.345	21.355	21.345	21.348	- '025	δ Orionis	-	26 53.873	53.844	53.865	53.870		
γ Ceti seq	+ 2 49	38 7.150	7.076	7.093	7.140	7.112	+ .029	a Leporis	1	28 19.174	19.174	19.200	19.160	19.177	006
π Ceti		39 21.785	21.785	21.815	21.793	21.795	+ .036	1 Orionis		30 32.468	32.494	32.454	32.480	32.474	002
μ Ceti	+ 9 42	39 32.100	32.082	32.123	32:114	32.105	+ .007			31 40.070	40.058	40.080	40.065	40.068	014
σ Arietis		45 58.218	58.198	58.186	58.194	58.199		ζ Leporis		42 25.455	25.456	25.450	25.449	25.453	+ .010
τ <sup>2</sup> Eridani	-21 25	46 30.121	30.166	30.125	30.144	30°147	+ .052	κ Orionis		43 0.836	0.830	0.830	0.826		
η Eridani		51 32.498	32.201	32.506	32.505	32.203	018	I Geminorum		58 2.477	2.486	2.458	2.200	2.480	1.
a Ceti	+ 3 42	57 3.092	3.078	3.100	3.070	3.085	+ .019	v Orionis		6 I 51.726	51.732	51.755	51.730	51.736	027
δ Arietis	+19 21	3 5 54 540	54.555	54.560	54.538			η Geminor. seq		8 50.472	50.467	50.488	50.443	50.468	
Contraction in the						01 01-	1		1	5-47-			5 115		1

TABLE I.-Right Ascensions of Clock Stars derived from Cape Ledgers.

vi

TABLE I.—continued.

Star, Dee		Right Ase	ension 190	0'0.		Corr. to New-	Star.	Dec.	F	light Asce	nsion 190	oʻo.		Corr. to New-
Star. 1900	I. E.	1. W.	11. E.	11. W.	Mean.	Comb.	Star.	1900.	1. E.	`I. W.	II. E.	II. W.	Mean.	Comb.
$\mu$ Geminorum $+22$	4 6 16 54.63	o 54.660	8 54·643	s 54.653	s 54.647	s - • 016	v Hydræ	-15 40	h m s 10 44 41 416	s 41°440	s 41.433	s 41.448	s 41°434	8 + ·027
B Canis Majoris 17			17.710	17.733	17.740	- '010	d Leonis		55 23.807	23.800	23.788	23.824		+ .012
8 Monocerotis + 4			28.160	28.145	28.153	000	χ Leonis		59 51.560	51.220	51.586	51.280		+ .004
10 Monocerotis 4		1	1.270	1.306	1.585	047	B Crateris		11 6 44.324	44.345	44'334	44.356		+ '015
v Geminorum +20		-	1.528	1.232	1.237	+ .003	δ Leonis		8 47 453	47.445	47.458	47 440	47.449	- '041
ξ Canis Majoris22			51.884	51.001	51.892	028	θ Leonis		8 59.605	59.605	59.613	59.628	59.613	
15 Monocerotis + 9			28.280	28.273	28.275	+ .011	δ Crateris		14 20.420	20.443	20.445	20.442	20.438	+ .011
ξ Geminorum +13	0 39 40.67		40.598	40.652	40.638	+ .007	σ Leonis		15 58.823	58.822	58.830	58.833	58.827	- '014
18 Monocerotis + 2			38.798	38.803	38.806	+ .021	$\tau$ Leonis		22 47 703	47'702	47.683	47.704	47.698	+ .002
θ Canis MajorisII			32.655	32.657	32.649	001	v Leonis		31 49 721	49.719	49.736	49.740	49.729	+ .004
y Canis Majoris15			14.064	14.070	14.062	007	ß Leonis		43 57 560	57.562	57.576	57.590	57.572	015
51 Geminorum + 16			37.770	37.720	37.754	- '041	ß Virginis		45 29.189	29.183	29.182	29.203	29.189	+ .000
A Geminorum +16			20.806	20.809	20.804	007	π Virginis		55 44.926	44.966	44.930	44.945	44.942	+ '026
δ Geminor. seq +22	-		9.066	9.070	9.073	034	o Virginis	+ 9 17	12 0 6.939	6.917	6.931	6.931	6.930	005
B Canis Minoris + 8 :			43.680	43.715	43.698	004	e Corvi		4 58.843	58.842	58.849	58.860	58.849	+ .007
25 Monocerotis 3			18.388	18.406	18.403	+ .034	η Virginis		14 47.361	47.357	47:374	47.363	47.364	028
26 Monocerotis 9		8 28.194	28.207	28.211	28.200	+ .053	δ Corvi seq	-15 58	24 41 320	41.323	41.350	41'311	41.326	035
к Geminor. seq +24	8 38 24 68	8 24.698	24.690	24.684	24.690	- '022	20 Comæ	+21 27	24 41.840	41.803	41.814	41.830	41.822	072
ξ Argûs seq24.	7 45 5'3	6 5.316	5.332	5.329	5.326	+ .008	ß Corvi		29 7 977	8.010	7.996	7.986	7.992	+ .041
9 Puppis m 13			8.480	8.453	8.477	009	24 Comæ seq		. 30 6.853	6.832	6.836	6.820	6.835	008
ρ Argûs24	1 8 3 17.11		17.134	17.164	17.136	+ .027	ρ Virginis	+10 47	36 49.398	49.383	49.378	49.391	49.388	- '026
20 Puppis15 :	9 8 44 18	5 44.197	44.200	44.301	44.196	+ .010	δ Virginis	+ 3 56	50 33.980	33.969	33.962	33.950	33.965	+ .002
β Cancri 9 :		7 5.581	5.584	5.547	5.575	+ .014	e Virginis	+11 30	57 11.944	11.928	11.930	11.938	11.935	008
Bradley 1197 3 :			39.880	39.853	39.862	009	θ Virginis	- 5 0	13 4 46 294	46.297	46.303	46.317	46.303	+ .014
η Cancri+20	7 26 55.61	3 55.629	55.290	55.613	55.611	019	γ Hydræ		13 29.031	29.038	29.052	29.050	29.043	+ .027
δ Hydræ + 6	3 32 21.77	7 21.793	21.768	21.793	21.783	- · 001	i Virginis	-12 II	21 26.126	26.147	26.131	26.116	26.130	+ .013
δ Cancri +18		5 0.220	0.302	0'210	0.208	100. +	ζ Virginis	- o 5	29 35.838	35.811	35.821	35.823	35.823	+0.008
e Hydræ AB + 6			28.862	28.872	28.871	006	m Virginis		36 21.772	21.737	21.776	21.758	21.761	+ .012
14 Hydræ – 3		-	20.241	20.248	20'244	027	au Boötis		42 30.616	30.602	30.296	30.288	30.601	002
ζ Hydræ+ 6 :		8 6.491	6.473	6.211	6.201	029	89 Virginis		44 26.187	26.206	26.233	26.209	26.209	+ .023
α Caneri +12			I.110	1.115	I.155	023	η Boötis		49 55.383	55.366	55.387	55.406	55.386	012
к Cancri+11	4 9 2 19.92		19.902	19.915	19.910	008	$\tau$ Virginis		56 33.406	33.300	33.384	33.373	33.388	- '015
$\theta$ Hydræ + 2			9.754	9.752	9.749	003	94 Virginis		14 0 59.983	59.980	59.978	59.994	59.984	+ .002
83 Cancri	8 13 24.10		24.098	24.070	24.085	- '021	κ Virginis		7 33.607	33.605	33.630	33.613	33.614	000
a Hydræ 8			40.445	40.443	40.440	+ .018	· Virginis		10 46.196	46.198	46.208	46.172	46.194	+ .015
ξ Leonis+11			33.438	33.403	33.421		λ Virginis		13 41.842	41.847	41.873	41.867	41.857	+ .023
и Hydræ13 к Hydræ+10		-	30.773	30.750	30.759		f Boötis		21 48.268	48.252	48.225	48.256	48.250	- '022
6 Sextantis 3			48.906	48.873	48.883	-	ζ Boötis m		36 22.360	22.354	22.338	22.378	22.358	042
$\pi$ Leonis			11.716	11.210	11.712	-	μ Virginis 109 Virginis		37 47 348	47'342	47.352	47.372	47 354	- 100
$\pi$ Leonis			55.767	55.817	55'791				41 11.552	11.224	11.220	11.222	11.242	- '020
λ Hydræ			52.898	52.906	52.915 42.781		a Libræ 15 Libræ		45 20.691	20.704	20.688	20.723		+ .013
22 Sextantis 7			42.774	42.795			Piazzi XIV. 221		51 20.450	20.463	20.438	20.445		+ .015
μ Hydræ				39.664	39.675		: Libræ		51 30.002	30°024 31°180	29.987	29.980		+ ·006 + ·009
p Leonis			15°244 32°767	15·250 32·776	32.786	1 .	β Libræ		15 6 31·167 11 37·472	37.482	31.201	31.170	37.482	002
33 Sextantis 1			18.965	18.985	18.983	and the second se	30 Libræ		17 27.037	27.059	37.482	37.490		+ .001
34 Sextantis + 4			27.698	27.703	27.696	1.	32 Libræ		22 36.922	36.914	36.928	36.920	36.921	011
l Leonis+11		_	0'100	0.076	0.092		γ Libræ		29 55.842	55.853	55.870	55.854	55.855	- '026
	· · · · · · · · · · · · · · · · · · ·				1						00-1-	55 51	55 55	

vii

TADLE I CONCONCUCA.	TABLE .	[.—continued	
---------------------	---------	--------------	--

City of the second seco	Dec.	I	Right Asc	ension 190	0.00		Corr. to		Dec.	1	Right Asc	ansion 19	00'0.		Corr. to
Star.	1900.	I. E.	I. W.	п. е.	II. W.	Mean.	New- Comb.	Star.	1900.	I. E.	I. W.	II. E.	II. W.	Mean.	New- Comb.
a Serpentis	+ 6 44	h m a 15 39 20.502	8 20.510	8 20°515	9 20°493	s 20*505	8 · - ·002	8 Aquilæ	+ 6 9	h m a	8	8 24°084	S	S	8
ß Sarpentis		41 34.288	34.314	34.310	34.294	34.302	052	γ Sagittæ		19 50 24.107 54 18.576	24.077	18.592	24.063 18.583	24.083	
« Serpentis		44 14.278	14'254	14.268	14°252	14.263	013	θ Aquilæ		20 6 8.743	8.758	8.752	8.757	18·579 8·753	- ·008 + ·018
μ Serpentis		44 24.039	24.044	24.032	24.035	24.038	+ '005	a <sup>2</sup> Capricorni		12 30.437	30.468	30.442	30.412	30.440	+ .021
e Serpentis	+ 4 47	45 49 820	49.804	49.804	49.838	49.817	013	A Capricorni		15 23.645	23.633	23.630	23.640	23.637	0021
γ Serpentis		51 49.990	49.992	49.994	50'002	49'995	033	e Delphini		28 26.142	26.112	26.113	26.144	-	- '018
δ Scorpii	-22 20	54 25.144	25.137	25.144	25.141	25.142	+ .015	ß Delphini		32 51.590	51.564	51.295	51.575	51.582	036
B Scorpii pr	-10 32	59 37.254	37.290	37.263	37.279	37.272	+ .036			34 21.483	21.478	21.513	21.203		+ .010
δ Ophiuchi		16 9 6.273	6.265	6.250	6.258	6.262	+ .002	α Delphini		34 59.600	59.573	59.600	59.598	59.593	010
e Ophiuchi	- 4 27	13 1.775	1.749	1.752	1.763	1.760	+ .002	e Aquarii		42 15.824	15.824	15.820	15.823		+ .010
γ Herculis	+10 23	17 30.477	30.493	30.506	30.496	30.493	015	μ Aquarii	- 9 22	47 15.663	15.676	15.683	15.663		+ .010
λ Ophinchi m	+ 2 12	25 52.198	52.158	52.146	52.168	52.168	+ .013	32 Vulpeculæ		50 17.839	17.836	17.862	17.842	17.845	039
ß Herculis	+21 42	25 55.212	55.240	55.255	55.208	55.229	+ .010	θ Capricorni		21 0 19.642	19.625	19.636	19.612		+ .023
Ç Ophinchi		31 39.078	39.086	39.090	39.102	39.089	- '001	v Aquarii	-11 47	4 8.848	8.866	8.880	8.865		+ .019
49 Herculis		47 31.654	31.637	31.649	31.657	31.649	025	α Equulei		10 49 506	49.542	49.535	49.528	_	+ '012
к Ophiuchi	+ 9 32	52 56.064	56.052	56.070	56.058	56.061	000	· Capricorni		16 40.780	40.764	40.786	40.787	40.779	008
η Ophiuchi m	-15 36	17 4 38.554	38.558	38.540	38.560	-	+ .032	1 Pegasi		17-27.680	27.708	27.680	27.662	c0.	- '024
δ Herculia	+24 57	10 55.417	55.428	55.393	55.365	55.401	025	ζ Capricorni		20 57 546	57.556	57.540	57.554	57 549	- '010
θ Ophiuchi		15 52.049	52.060	52.038	52.048	52.049	+ .010	B Aquarii		26 17.727	17.731	17.735	17:745		+ .019
d Ophiuchi	-29 47	20 58.080	58.102	58.084	58.080	58.087	+ .071	ξ Aquarii		32 25 755	25.755	25.740	25.748	25.750	- ·co4
σ Ophiuchi	+ 4 14	21 33.157	33.152	33.138	33.128	33.121	020	γ Capricorni	-17 7	34 33'090	33.079	33.073	33.085		- '017
α Ophiuchi	+12 38	30 17.552	17.553	17.532	17.530		+ .002	e Pegasi		39 16.480	16.465	16.476	16.498		+ '012
ξ Serpentis		31 51.602	51.618	51.612	51.610		+ .024	δ Capricorni	-16 35	41 31.350	31.344	31.376	31.388		+ .021
ß Ophiuchi		38 31.944	31.934	31.932	31.932		011	16 Pegasi		48 30.660	30.675	30.668	30.675		035
v Ophiuchi	- 9 46	53 31.284	31.272	31.265	31.266		+ .002	α Aquarii		22 0 38.898	38.897	38.886	38.914		+ .003
67 Ophiuchi	+ 2 56	55 38.196	38.170	38.170	38.168	0 -	- '045	( Aquarii		1 2.268	2.260	2.233	2.255		+ .027
72 Ophinchi	+ 9 33	18 2 36.533	36.520	36.516	36.538		+ .013	θ Pagasi		5 9.340	9.330	9.308	9.328	9.326	024
μ Sagittarii	-21 5	7 46.985	46.982	46.988	46.980	-	+ .019	θ Aquarii		11 33.458	33.448	33.447	33.473		+ .013
η Serpentis		16 8.130	8.132	8.130	8.128	8.130	+0.028	γ Aquarii	- 1 53	16 29.523	29.508	29'470	29.530		+ .012
109 Herculis		19 26.163	26.190	26.165	26.193	26.178	012	σ Aquarii		25 21.352	21.362	21.362	21.347		014
λ Sagittarii		21 47 966	47 967	47.948	47.980	47.965	+ '001	η Aquarii		30 13.123	13.080	13.081	13'103		+0.011
Scuti 4 H		36 47 909	47.904	47 912	47 923	47.912	032	ζ Pagasi	+10 19	36 28.455	28.442	28.453	28.440	28.448	026
φ Sagittarii	-27 6	39 24.552	24.564	24.554	24.578	24.562	+ .031	λ Pegasi		41 42.798	42.814	42.780	42.764		025
110 Herculis		41 21.448	21.478	21.483	21.443	21.463	+ .000	λ Aquarii	- 8 7	47 23.896	23.898	23.883	23.894	23.893	+ .013
θ Serpentis pr		51 14.924	14'900	14.918	14.908	14.913	+ .029	δ Aquarii	-16 21	49 20.658	20.650	20.667	20.668	20.661	+ .033
ξ Sagittarii		51 45.881	45.882	45.863	45.870	45.874	+ .019	a Pegasi	+14 40	59 46.743	46.773	46.740	46.725	46.745	+ .004
e Aquilæ	+14 56	55 5.047	5.033	5.030	.5.025	5.034		c <sup>2</sup> Aquarii		23 4 6.939	6.973	6.941	6.953		+ .025
ζ Aquilæ pr		19 0 48.848	48.823	48.803	48.808	48.821		γ Piscium		11 58.873	58.875	58.870	58.873	58.873	.000
λ Aquilæ		0 56.230	56.543	56.553	56.537	56.541	+ .013	7 Pegasi	+23 12	15 41.185	41.170	41.188	41.170	41.178	002
$\pi$ Sagittarii		3 49.035	49.050	49.058	49.077	49.055	+ .020	v Pagasi	+22 51	20 23 204	23.237	23.226	23.220	23.222	018
ψ Sagitarii		9 24.572	24.567	24.593	24.598	24.583		к Piscium		21 48.360	48.370	48.367	48.388	48.371	010
ω Aquilæ		13 7.362	7.320	7.356	7.362	7.350		70 Pegasi		24 5.812	5.820	5.813	5.792	5.809	+ .010
δ Aquilæ		20 27.408	27.405	27.408	27.415	27.409		e Piscium		34 48.370	48.392	.48.390	48.372	48.381	013
μ Aquilæ		. 29 12.255	12.234	12.242	12.265	12.248		ω <sup>2</sup> Aquarii pr		37 32.223	32.229	32.233	32.193	32.220	+ :002
54 Sagittarii		34 59 703	59.705	59.713	59.698	59.705		φ Pegasi		47 23.998	23.995	23.968	23.964	23.981	+ .005
f Sagittarii		40 31.776	31.748	31.790	31.797	31.778		ω Piscium		54 10.542	10.227	10.220	10.218	000 1	015
γ Aquilæ		41 30.340	30.330	30.363	30.344	30.344		2 Ceti	-17 54	58 37.048	37.063	37.053	37.063	37.057	+ .000
δ Sagittæ	+18 17	42 55.690	55'740	55.720	55.212	55.716	- '026						1.526-		
			1		1										

viii

On comparing the entries in columns 3, 4, 5, and 6 of this table with the mean contained in column 7, the discordances between right ascension observations made in the four conditions of the instrument may be summarised as follows :---

R.A.		۵	α.		No. of
	I. E.	I. W.	II. E.	II. W.	Stars.
h h O— I	s + 0.004	8 + 0'002	8 - 0.001	- 0°006	II
I — 2	+ .003	- :005	003	+ .002	11
2-3	'000	003	- :002	+ .004	15
3	+ .001	'000	004	+ '002	14
4-5	+ .001	+ .001	+ .004	006	12
5-6	001	•000	- *004	+	11.
6 7	+ .002	002	004	+ .001	13
7-8	+ .003	+ .001	100' -	- '003	9
8 9	003	+ .006	005	100' +	II
9-10	+ '004	009	+ .002	- °002	9
1011	- '002	.000	- '003	+ .003	II
I I I 2	007	.000	001	+ .008	IO
12-13	+ .002	004	100. +	003	IO
13-14	100. +	005	+ '004	.000	9
14-15	100	+ .005	003	+ '002	11
15-16	006	• 000•	+ .002	100	13
16—17	+ •002	004	+ .001	.000	. 8
17-18	+ .006 .	+ .002	002	006	10
18—19	.000	+ .001	003	+ .002	11
19—20	001	008	+ .002	+ .004	13
20—2 I	.000	004	+ .004	100. —	10
2 I 2 2	004	•000	+ .001	+ .003	12
22-23	+ .002	+ '002	110	+ .001	12
23-24	- 0.003	+ 0.006	+ 0.003	- 0 <b>.0</b> 02	II

 TABLE II.—Discordances between Time Determinations in the Four

 Different Conditions of the Transit Circle.

There appears to be no sensible trace of systematic run in these residuals, as may be expected since each column is necessarily constrained to follow the system of C. F. C., 1900. 2 Newcomb's Catalogue. If we similarly compare the mean results with those of Newcomb's Catalogue, the comparison may be summarised as follows :----

 
 TABLE III.—Comparison of Cape Ledgers with Newcomb's Catalogue in order of Right Ascension.

R.A.	Δα.	No. of Stars.	R.A.	Δα.	No. of Stars.
h h	8		h h	S	
0— I	-0.001	11	12-13	-0.013	IO
I— 2	+ .003	II	13-14	+ .002	9
2-3	+ .004	15	14-15	003	11
3-4	003	14	15-16	002	13
4-5	+ .001	12	16-17	003	8
5-6	003	II	17-18	+ .002	IO
6-7	- '012	13	18-19	+ '010	11
7-8	- '002	9	19-20	+ '002	13
8-9	006	II	20-21	- '004	IO
9-10	+ .006	9	21-22	+ .001	12
10-11	+ .000	II	2223	+ '002	12
II—I2	+0.005	10	23-24	-0*001	. 11

(Cape Ledgers-Newcomb).

Here again the differences are insignificant. If, however, we arrange the stars in order of declination, we derive the following summary of results :---

### TABLE IV.—Comparison of Cape Ledgers with Newcomb's Catalogue in order of Declination.

Limits of Declination.	Δα.	No. of Stars.	Limits of Declination.	Δα.	No. of Stars.
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$ \begin{array}{r} 8 \\ -0.023 \\ -0.028 \\ -0.013 \\ -0.04 \\ -0.015 \\ -0.013 \\ -0.00 \\ -0.002 \\ -0.003 \\ +0.003 \\ \end{array} $	1 I 1 I 1 I 1 I 1 I 1 I 1 I 1 I	$\begin{array}{cccccccccccccccccccccccccccccccccccc$		I1           I1           I1           I1           I2           I1           I2           I1           I1

(Cape Ledgers-Newcomb).

Thus the observations indicate a correction to the adopted clock star places, dependent on the declination, which is zero at, or slightly to the north of, the equator,

but which increases southwards at the rate of about 0<sup>s</sup> 001 per degree of declination. The effects of such an error on the periodic errors in R.A. will, however, be insignificant, as the clock stars in the higher declinations are fairly uniformly distributed in right ascension, as is evidenced by the following table, showing the distribution of clock stars in declination.

R.A.	Mean Dec.	No. of Stars.	R.A.	Mean Dec.	No. of Stars.
$ \begin{array}{c}     h & h \\     0 I \\     I 2 \\     2 3 \\     3 4 \\     4 5 \\     5 6 \\     6 7 \\ \end{array} $	$ \begin{array}{r} - 1 \\ - 1 \\ + 3 \\ + 4 \\ + 8 \\ - 4 \\ + 3 \end{array} $	11 11 15 14 12 11 13	h h 12-13 13-14 14-15 15-16 16-17 17-18 18-19	+ $i- 3- 2- 4+ 6- 5- 3$	10 9 11 13 8 10 11
7 8 8 9 910 1011 1112	+ 4 + 3 + 4 + 3 + 3	9 11 9 11 10	$19-20 \\ 20-21 \\ 21-22 \\ 22-23 \\ 23-24$	+ 4 - 4 - 3 + 3	13 10 12 12 11

TABLE V.—Distribution of Clock Stars.

In consideration of this approximate symmetry of distribution any errors in the right ascensions of Newcomb's Catalogue dependent on the declinations may be regarded as sensibly eliminated from the mean results of the Cape Ledgers, and so far as such errors are concerned the latter may be regarded as defining an independent fundamental system. In so far, however, as the errors of the original system depend on the right ascension they will only be partially smoothed out, the more wide-spread features being reproduced almost in their entirety. To examine such errors recourse must be had to additional observations which have not otherwise been included in the formation of the Catalogue.

### Discussion of Daylight Observations of Clock Stars.

In addition to the observations made in the night watches directly for the purposes of the Catalogue, regular observations of the Sun and inferior planets have been made by day. These have always been accompanied by observations of bright stars for the determination of clock error. The stars used are for the most part contained in the above clock star list, but include also the following additional stars, the places quoted being derived from the Cape Ledgers in the same manner as for the former stars.

Star.	Dec. 1900.	R. A. 1900'0.	Corr. to New- comb.	Star.	Dec. 1900.	R. A. 1900'0.	Corr. to New- comb.
a Tauri $\beta$ Orionis	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	h m s 4 30 10'914 5 9 43'936 5 19 46'044 5 23 57'607 5 31 8'356 5 49 45'502 6 16 28'457 6 31 56'119 6 40 44'633 6 54 41'744 7 4 19'542 7 20 8'360 7 34 4'089 7 39 11'845 9 40 10'585 10 3 2'857 11 28 4'927		γ Corvi         a Virginis	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	h m s 12 10 39'740 13 19 55'432 14 11 6'017 14 58 12'932 15 52 48'050 16 15 6'518 16 23 16'488 16 43 41'157 17 59 23'072 18 14 35'542 18 17 32'136 18 49 3'929 18 56 14'998 19 45 54'263 22 52 7'580 22 58 55'496	

TABLE VI. - Additional Clock Stars used for Daylight Observations.

Clock errors have been derived at the instant of each daylight transit, utilising the places of the stars as contained in the above tables and Newcomb's proper motions. Care has always been taken to control the level and azimuth variations by simultaneous reference to the nadir trough and meridian marks.

The clock errors at the same instants have also been derived by interpolation from adjacent night watches, assuming a uniform rate derived from observations approximately twenty-four hours, or a multiple thereof, apart. The details of the comparison between the two methods of determination are contained in the following table :--

	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day— Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day— Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.
	1908.		h m	h m	S	1908.		h m	h m	8	1908.		b m	h m	8
M	lay 4	М	0 39	11 23	-0.03	May 13	Μ	0 39	12 12	-0.05	May 20	RC	5 10	11 42	-0.03
	5		5 10	23	+ .02	14		5 10	22	+ .06	20		5 20	,,	+ .07
	5	215	5 20	>>	.00	17	JJ	0 39	12 7	01	20	C	0 39	11 38	07
	11	C	0 39	11 6	03	17		2 57	,,	+ .08	20		1 19	"	+ .01
	12		4 30		02	18	5.45	5 10	,,	+ .06	2 I		5 10	,,	03
	12		5 10	29	04	18		5 20	"	+ .02	21		5 20	,,,	+ .02
	12	AW	0 39	11 22	+ .02	18	M	0 39	12 15	- '01	26	AW	2 57	13 2	.00
	13		5 3	>>	+ .08	19		5 10	>>	+ .02	27		6 18	>>	+ .05
	13		5 10	,,	+ .06	19		5 20	>>	+ .08	27		6 54	,,	+ '02
30	13		5 20	,,,	.00	19	RC	0 39	11 42	03	27		7 4	>>	+ .03

TABLE VII.—Comparison of Day and Night Determinations of Clock Error.

TABLE VII.—continued.

i			-							1	1				
	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.
	1908. June 5	M	h m 6 4 I 6 54	h m 12 49	s +0.01 02	1908. July 15 15	RC	h m 5 32 5 43	h m 1637	8 +0.01 + .03	1908. Sept. 7 13	RC M	h m 10 3 9 23	h m 20 14 19 52	8 +0.05 + .02
	555	JW	1 49 2 1 2 57	13 O "	+ ·03 + ·07 + ·02	16 16 16	AW	5 32 5 43 6 19	16 37 "	+ ·04 + ·05 - ·01	14 - 14 15	AW JJ	9 23 10 3 9 23 10 3	20 4 20 38	+ ·12 + ·08 + ·07 + ·03
	9 9 10 10		$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	13 40 "" "	$ \begin{array}{r} - & \cdot 04 \\ + & \cdot 02 \\ + & \cdot 02 \\ - & \cdot 01 \\ \end{array} $	19 19 19 19	JJ	5 10 5 20 5 32 5 43	15 29 " "	$+ \cdot 03$ + $\cdot 02$ + $\cdot 01$ + $\cdot 06$	I 5 2 1 2 1 2 2	JJ AP	10 3 9 23 10 3 9 23	,, 20'31 ,, 20'49	+ .03 + .03 + .13 + .01
	15 15 16 16		5 3 5 10 7 4	14 10 "	+ .02 + .05 01	20 20 20	M C	6 55 7 5 7 34 5 20	15 29 " 16 20	+ ·03 + ·08 + ·05 + ·04	22 23 23	AW M	IO 3 9 23 IO 3	21 2 ,, 20 57	- 01 + 03 - 01 + 01
	16 17 17	M	7 20 5 10 7 5 7 20	" 14 10 "	$- \frac{.02}{.00}$ + $\frac{.03}{.00}$	22 22 22 23	AP	5 20 5 31 5 43 5 10	10°20 ,, 16°18	+ .05 + .05 + .03	24 24 27 29	JJ JW	10 3 10 3 9 23	20 57 21 14 20 56	+ ·02 + ·01 - ·03
	17 18 18		5 10 7 5 7 20 2 57	14 25 ,, ,, 14 40	$- \cdot 01$ $- \cdot 06$ $- \cdot 01$ $+ \cdot 03$	23 23 23 26	JW	5 20 5 32 5 43 5 10	" " 16 26	$+ \cdot 02$ + $\cdot 03$ + $\cdot 06$ + $\cdot 03$	29 30 30 Oct. 1	AP	10 3 9 23 10 3 14 11	21 10 ,,	+ ·03 - ·03 - ·03
	19 19 19		3 19 3 29 4 30	27 T 33 33	- ·02 - ·05 ·00	26 26 27	AW	7 20 7 34 5 32	" 16 <sup>"</sup> 33	+ <sup>•04</sup> + <sup>•01</sup> + <sup>•08</sup>	1 1 5 6	RC M	9 23 10 3 10 3	21 19 ,, 21 56	10 00.
*	19 22 22 23	C	5 10 5 10 5 43 7 20	", 14 40 ",	+ .04 + .10 + .07 + .07	27 27 29 29	м	5 43 6 18 6 41 6 55	" 16 <sup>°°</sup> 53	+ .10 + .02 01 01	. 6 7 7	AW C	11 9 11 44 14 11 10 3	21 56 " 22 30	$- \cdot 02$ $- \cdot 03$ $+ \cdot 03$ $+ \cdot 03$
	28 28 29		5 20 5 32 7 34	15 O ,; *,;	+ '12 + '09 + '03	29 29 30	AP	7 5 7 34 7 4	" 17 <sup>°</sup> 0	- ·01 - ·05 + ·01	8 8 8 8	AP	14 11 10 3 11 9	22 30 "	+ .03 + .05 + .04 06
	29 29 29 29		5 10 5 20 5 32 5 43	14 57 ""	+ ·03 + ·01 - ·05	30 Aug. 9 9 9	JJ	7 34 6 19 6 55 7 34	19°0 "	+ ·04 ·00 - ·01	9 15 18	JJ C	11 44 14 11 11 44 11 9	" 22 20 21 31	- ·10 - ·06 - ·02
	30 July 5 5 6	JJ	7 34 5 32 5 43 4 30	" 15 <sup>°</sup> 20 " 15 <sup>°</sup> 20	·00 + ·08 + ·02 + ·10	- IO II II II	M RC	7 34 6 55 7 20 7 34	18 0 17 46 ,,	- ·03 + ·04 + ·04 + ·05	18 21 22 22	M AW	11 44 11 44 11 9 11 44	"22 50 23 9	·00 - ·07 + ·04 + ·09
	6 6 6		5 3 5 10 5 20	25 20 27 27 27	+ ·05 + ·02 + ·08	14 14 14	AW JW	11 9 6 32 6 41	18 <sup>°</sup> 30 1848 "	- ·08 - ·03 ·00	22 Nov. 2 2	JJ	12 11 11 44 12 11	" 23 13 "	+ .06 02 + .02
	6 7 7 7	AP	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	" 15 <sup>20</sup> "	+ ·04 + ·04 + ·05 + ·03	14 16 16 20	JJ M	6 55 6 41 7 5 7 20	18 <sup>"</sup> 16 18 <sup>"</sup> 16	- ·02 + ·06 + ·04 - ·06	I 2 1 2 1 5 1 7	JW JJ C	13 20 14 11 13 20 12 29	23 40 " 23 54 0 6	$ \begin{array}{r} - & 0 \\ - & 0 \\ + & 0 \\ + & 0 \\ + & 0 \\ \end{array} $
	7 8 8 8 8 8		4 30 5 10 5 20	15 20 ,, ,,	+ ·03 - ·02 - ·03	20 20 21	AW	7 34 7 39 6 55	" 18 <sup>°</sup> 0	- ·05 - ·01 - ·04 - ·05	17 19 19	RC JJ	13 20 13 20 14 11	" 0 52 "	+ .01 + .10 + .08
	35		5 32 5 20	16"37	10 <sup>.</sup> +	21 Sept. 7	RC	7 34 9 23	20'14	- ·05 - ·05	22 22	00	13 20 14 11	0 30 "	+ .04 + .06

Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day— Night.
1908.         Nov. 23         23         23         Dec. 1         3         6         7         13         13         13         13         13         13         13         13         13         13         13         13         13         13         13         13         13         13         14         20         20         20         20         20         21         20         21         20         21         20         21         24         25         27         28         29         20         21         24         25         27         28         29         31         31         31         32	C C JW AP JW JJ M C M JJ RC JW AW M JJ JC AP	18 22 18 50 18 57	h m o 30 " 1 35 2 19 3 5 3 34 " 2 33 " 2 33 " 2 54 " 2 45 2 45 2 45 2 45 2 45 5 0 " 5 0 " 5 0 " 5 14 " 5 24 " 5 24 " 5 24 5 39 " 6 1 " " 5 50 " " " 5 50 " " " 5 0 " " " 2 45 5 50 " " " 5 0 " " " " 2 45 5 50 " " " " " 2 45 5 50 " " " " " " " " " " " " " " " " " " "	- '07 - '04 - '10 - '08 - '10 - '15	5	AW C AP	23 0 21 40 22 1 23 0 22 1 22 53	h m 5 56 6 24 5 47 3 47 3 47 3 47 3 6 39 3 6 46 6 45 7 53 7 50 7 40 7 27 8 8 3 7 50 7 40 7 27 8 8 3 8 3 4 3 8 3 4 3 8 3 4 3 9 11 3 9 45 9 45 10 12 10 16	$\begin{array}{c} s\\ -0.01\\ -0.01\\ -0.01\\ -0.01\\ -0.02\\$	1909.         Apr.       9         12       13         15       15         15       15         16       10         23       23         23       23         23       23         25       26         May       2         6       6         7       7         9       10         10       10 <td< td=""><td>AW JJ AW C AW JJ AW C JJ AW C JJ AW JJ AP RC AP JJ JW C JJ K C AP RC AP RC</td><td>h m 0 399 0 399 22 53 23 0 0 399 22 53 0 399 22 53 0 399 23 0 0 990 39922 23 0 0 990 39922 23 0 0 990 3900 3000 3000 30</td><td>h m 10 16 10 36 10 59 10 14 " 10 54 " 10 54 " 10 49 " 9 57 10 19 " 11 22 " 11 28 12 35 12 35 13 8 " 12 35 13 8 " 12 35 13 8 " 12 35 13 8 " 13 4 " " 13 23 12 54 " 13 4 " " 13 23 12 54 " 13 4 " " 13 23 12 54 " " 13 4 " " " " " " " " " " " " "</td><td>s + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 +</td></td<>	AW JJ AW C AW JJ AW C JJ AW C JJ AW JJ AP RC AP JJ JW C JJ K C AP RC AP RC	h m 0 399 0 399 22 53 23 0 0 399 22 53 0 399 22 53 0 399 23 0 0 990 39922 23 0 0 990 39922 23 0 0 990 3900 3000 3000 30	h m 10 16 10 36 10 59 10 14 " 10 54 " 10 54 " 10 49 " 9 57 10 19 " 11 22 " 11 28 12 35 12 35 13 8 " 12 35 13 8 " 12 35 13 8 " 12 35 13 8 " 13 4 " " 13 23 12 54 " 13 4 " " 13 23 12 54 " 13 4 " " 13 23 12 54 " " 13 4 " " " " " " " " " " " " "	s + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 +

TABLE VII.—continued.

TABLE VII.—continued.

İ	Da	te	Observer.	S.T. of Day	Mean S.T. of	Diff. Day-	Date.	Observer.	S.T. of Day	Mean S.T. of	Diff. Day-	Date.	Observer.	S.T. of Day	Mean S.T. of	Diff. Day-
	Du		Obse	Obs.	Night Obs.	Night.	2	Obse	Obs.	Night Obs.	Night.		Obsi	Obs.	Night Obs.	Night.
	190 June	1 2 2	RCC	h m 6 41 5 3 5 10	14 18 14 18 ,,	+0.02 + .03 + .04	1909. Aug. 12 12	C JJ	h m 7 35 8 4 7 35	h m 18 32 18 33	8 + 0°03 + °01 + °03	1909. Oct. 18 19 19	AP C	h m 15 53 15 12 15 55	h m 22 24 22 21 ,,	8 +0.02 07 + .04
		8 9 10 10	JW JW C	5 32 7 5 2 57 6 32 6 41 4 30	13 55 13 58 " 14 11	$+ \cdot 02 + \cdot 05 + \cdot 02 - \cdot 04 + \cdot 03 + \cdot 02$	16 16 16 16 19 20	JJ C RC	11 44 6 55 7 35 8 4 11 44 11 44	" 19 <sup>°</sup> 20 " 19 <sup>°</sup> 6 18 <sup>°</sup> 51	$+ \cdot 01$ $- \cdot 04$ $+ \cdot 04$ $- \cdot 04$ $- \cdot 02$ $- \cdot 04$	19 20 20 21 21 21	RC JJ	16 24 16 0 16 24 15 40 15 55 16 0	"22 17 22 29 "	$ \begin{array}{r} - \cdot 02 \\ - \cdot 03 \\ + \cdot 01 \\ \cdot 00 \\ - \cdot 04 \\ + \cdot 04 \end{array} $
		11 11 18 18 30	11 11	6 19 6 32 2 57 4 31 5 10		- ·08 + ·06 - ·01 + ·01 + ·03	20 23 24 24 24	C	12 5 9 23 11 44 12 5 12 5	18 <sup>°</sup> 13 <sup>°°</sup> 17 <sup>°</sup> 57	$ \begin{array}{r} - & \cdot 02 \\ + & \cdot 03 \\ + & \cdot 03 \\ + & \cdot 04 \\ + & \cdot 03 \end{array} $	22 22 22 25 25	M AP RC	15 55 16 0 13 20 15 55 16 24	22 35 " 22 33 20 54 "	06 03 + .10 + .04 + .07
	Jul <b>y</b>	30 30 4 4 5	AP C	5 20 5 32 5 10 5 20 5 32 5 20	" 15 37 " 15 57	$+ \cdot 06$ $- \cdot 03$ $- \cdot 03$ $+ \cdot 05$	26 26 26 27 Sept. 1	AP AP	11 44 12 5 12 11 12 11 9 23 10 4	19 15 " 19 1 19 23	+ .01 01 + .01 + .02 + .04	29 Nov. 1 5 5 7	RC M AW	13       20         16       24         17       5         12       58         13       20         12       30	0 32 23 0 " 23 31 " 23 20	$ \begin{array}{r} - & 02 \\ - & 01 \\ - & 09 \\ + & 06 \\ + & 10 \\ - & 02 \end{array} $
		56666	JJ	5 50 4 31 5 3 5 10 5 20	" 15 50 " "	+ .09 + .07 + .05 + .08 + .04	2 3 3 9 9	M C	12 25 12 25 12 30 12 30 13 14	" 19 <sup>"</sup> 12 " 19 <sup>"</sup> 30 "	+ ·04 + ·07 - ·01 + ·04	7 7 8 8 Dec. 8	JJ	12 58 13 20 16 44 17 5 18 50	" " " " I 54	+ '01 + '01 + '07 + '04 '00
1		7 7 11 11 16	RC JW AP	5 3 5 10 5 20 5 10 5 20 5 10 5 10	15 43 " 16 <sup>°</sup> 24 16 <sup>°</sup> 52	$ \begin{array}{r}         :00 \\         + :03 \\         + :07 \\         + :03 \\         + :06 \\         $	13 13 13 14 14	JJ RC C	12 30 13 14 13 20 13 14 13 20 13 20 12 30	19 44 "" 19 44 "20 18	+ ·03 + ·05 + ·05 + ·01	8 8 13 14 14	M AW JW	19 42 19 46 19 46 19 42 19 46	" 1 30 1 45 "	+ .04 + .02 09 06 + .04
		16 16 16 18 18	JJ	5 10 5 20 5 32 6 19 6 19 6 32	10'32 "" 16'38	+ 00 + 00 + 00 + 00 + 00 + 00 + 00 + 0	15 15 16 16	AP	12 30 13 14 13 20 12 30 13 14 13 20	1, ,, 20 <sup>°</sup> 52 ,, ,,	04 04 07 + .04 02 03	15 16 16 21 21	RC JJ	19 46 19 42 19 46 19 46 20 16	1 57 2 22 " 2 35 "	- ·07 - ·12 - ·05 - ·05 - ·01
		19 20 20 20 22	JJ RC	10 4 5 10 5 32 5 44 5 32	" 15 <sup>°</sup> 50 " 17 <sup>°</sup> 11	+ .09 + .0602 + .02 + .05	17 17 21 21 27	C ]]	13 14 13 20 13 14 13 20 11 44	20 7 ,,, 20 13 ,,, 21 35	+ .07 + .03 + .01 + .03 + .02	1910. Jan. 6 6 6	JJ	15 55 16 24 16 32 16 44	23 23 23	- ·02 + ·02 - ·01 + ·11
	Aug.	22 22 30 30 2 2	RC C	6 19 6 41 5 44 5 50 5 20 5 32	" 17"45 " 17"34	$\begin{array}{r} + & \cdot 02 \\ - & \cdot 01 \\ + & \cdot 03 \\ + & \cdot 08 \\ + & \cdot 04 \\ + & \cdot 06 \end{array}$	28 28 Oct. 1 1 8 8	AP C	13 20 14 12 10 4 11 9 13 20 14 46	20°42	$ \begin{array}{r} - \cdot 04 \\ + \cdot 03 \\ + \cdot 12 \\ \cdot 00 \\ - \cdot 07 \\ - \cdot 02 \\ \end{array} $	9 9 9 10 10		15 55 16 24 17 5 15 55 16 24 16 32	3 51	- ·04 - ·04 + ·01 - ·01 - ·04 - ·07
		2 3	AP	5 44 10 4 7 35	>> >>	+ '07 + '02 + '04	8 18 18		15 12	22 24	- ·04 - ·09 - ·06	10 11	M	16 32 16 44 17 31 16 24	3 53 -	+ ·03 - ·22 + ·03

TABLE VII.—continued.

Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day— Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.
Igio. Jan. I I I I I I I I I I I I I I I I I I I	3 AP 3 AP 3 G JW 6 G 9 AP 9 P 1 C 1 5 JJ 5 JJ 5 KC 6 6 6 7 M 7 8 AW 8 8	obs.           h         m           16         44           17         5           16         24           16         32           17         5           18         15           18         22           18         15           18         22           17         31           17         39           18         15           18         22           17         31           17         39           18         22           18         18           18         22           18         18           18         22           18         18           18         22           18         18           17         39           18         22           17         31           17         39           18         22           18         50           19         46           19         46           19         46	Night Obs. h m 4 5 3 53 " 5 6 " 5 3 " 5 5 3 " 5 5 2 " 4 55 " 8 34 8 25 " 8 34 8 25 " " 9 0 " 9 19 " 9 19 " 9 19 " " 11 17 " " 13 18 " " 13 18 " " 13 2 " 13 2 " 13 3 ]	$ \begin{array}{c} \text{Night.} \\ * & 0.08 & 0.7 & 5.0 & 0.9 & 0.0 &$	Date. 1910. Mar. 6 6 8 8 9 16 16 20 20 22 22 23 28 28 29 30 30 30 30 30 30 30 30 30 30	RC M AW JW RC M JW RC M JJ JW AW AW JJW AW JW AP AP RC JW C M AW RC		Night	N ight. $s$ $s$ $-0.033$ $0.090$ $0.022$ $0.002$ $0.070$ $0.002$ $0.022$ $0.002$ $0.022$ $0.002$ $0.022$ $0.002$ $0.022$ $0.002$ $0.022$ $0.002$ $0.022$ $0.002$ $0.022$ $0.002$ $0.022$ $0.0022$ $0.022$ $0.0022$ $0.022$ $0.0022$ $0.022$ $0.0022$ $0.0222$ $0.0022$ $0.0222$ $0.0022$ $0.0222$ $0.0022$ $0.0222$ $0.0022$ $0.0222$ $0.0022$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.0222$ $0.02222$ $0.0222$ $0.02222$ $0.02222$ $0.022222$ $0.022222$	Date. 1910. May 2 5 6 8 8 9 11 12 12 12 13 13 13 13 13 14 15 15 15 19 10 26 26 27 27 30 July 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Lago C AP AW AW AW AW AW AP RC M AP RC M AP RC M AP RC M AP AW		Night	$\begin{array}{c c c c c c c c c c c c c c c c c c c $

TABLE VII.—continued.

		_								-	-	-	-	1	-
Date		Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day— Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.
	25 25 25	AP	h m 5 10 5 20 5 29	h m I 9 ,,	8 +0·10 + ·11 + ·05	1910. Sept. 21 21 28	JW RC	h m 9 23 10 4 9 23	h m 1 48 "3 6	s 0.00 05 + .04	1911. Jan. 25 25 26	JJ	h m 18 15 18 22 21 40	h m 4 34 ,,	8 +0.02 + .01 01
	25 25 26 26	RC	5 43 5 50 5 20 5 29	,, ,, 1 23	+ ·07 + ·07 + ·11 + ·07	28 29 29 30	M AP AW	11     9       10     4       11     9       10     4	,,, 3 3 ,, 3 2	·00 + ·04 - ·02 + ·06	30 30 30 31	AW RC	18 15 18 22 19 46 18 15 18 22	5 47 " 5 9	+ ·03 - ·07 - ·08 - ·03 - ·08
Aug.	26 31 31 31 31 1	AW RC	5 50 6 19 6 32 5 43	2 I 20 ,, 20 30	+ .05 + .09 + .04 + .05 + .03	3 10 10 18	M C	11 9 10 4 11 44 11 44	3 9 2 <sup>°</sup> 38 <sup>°</sup> 217	+ .14 + .03 02 05 + .02	3 I Feb. I 3 3	M JJ	18 22 18 50 18 50 19 46	5 <sup>"15</sup> 6 <sup>"</sup> 18	·00 ·00 - ·07 - ·06
	1 1 5 5 5 5	AP	6 55 7 4 5 43 5 50 6 19	" 19 <sup>°</sup> 48 "	+ ·03 + ·05 + ·09 + ·11 + ·09	21 21 23 23 24	JW JW AW	12 5 12 11 12 5 12 11 12 11	4 0 2 58 "2 55	$ \begin{array}{r} - & 01 \\ + & 02 \\ + & 01 \\ + & 06 \\ + & 02 \\ \end{array} $	7 7 8 8 9	AW M JJ	22 53 22 59 22 53 19 4 22 53	6 27 ,, 5 44 5 42 ,,	$ \begin{array}{r} - & 0.03 \\ + & 0.07 \\ - & 0.02 \\ & 0.00 \\ - & 1.1 \\ \end{array} $
	5 16 16 16 16	C AP	7 21 6 55 7 5 7 45 6 55	21 54 " 21 55	+ .05 + .04 + .01 + .05 + .03	26 26 Nov. 3 3 8	RC JW JW	12       5         13       20         12       29         13       20         12       30	3 44 2 35 1 51	+ '11 + '08 - '06 - '05 + '07	9 9 9 16 16	RC AP	18 50 19 4 19 46 18 50 19 46	5 42 " 6 <sup>"</sup> 51	- ·04 - ·03 + ·04 - ·03 + ·07
	17 19 23 23	M M	7 5 8 4 12 5 12 11	21 <sup>"</sup> 52 23 6 "	+ ·04 ·00 + ·02 - ·03	8 13 13 14	JW C	13 20 13 20 14 11 13 20	2 <sup>"16</sup> "219	$+ \cdot 02$ $- \cdot 04$ $+ \cdot 04$ $+ \cdot 02$	17 19 19 20	AW AP	22 53 19 42 19 46 19 46	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	- ·03 - ·04 - ·08 + ·01
Sept.	I I 2 2 2 2	AW M	8 4 9 23 12 11 8 51 9 41	22 3 ,, 22 3 ,,	+ ·08 + ·05 + ·05 - ·07 - ·01	14 16 16 23 27	AW C JW	14 11 13 20 14 11 14 11 13 20	"3 24 "2 46 4 20	$+ \cdot 04$ + $\cdot 03$ $- \cdot 01$ $- \cdot 02$	20 24 24 27 27	J] C	20 16 19 42 19 46 19 42 19 46	7 <sup>°</sup> 2 <sup>°°</sup> 7 <sup>°</sup> 39 <sup>°°</sup>	01 05 04 + .09 + .03
	7 7 8 9	AW JW AP	9 23 10 4 9 23 8 4	23 13 ,, 23 25 20 41	+ .08 + .01 08 + .01 + .02	28 28 29 29 Dec. 9	AP RC C	13 20 14 11 14 11 14 46	4 20 " 1 27 " 2 30	+ .06 + .02 + .09 + .12 + .07	27 27 Mar. 1 1	AW	20 7 20 16 19 42 19 46	" " 7 42 "	$+ \cdot 03$ $+ \cdot 08$ $- \cdot 06$ $- \cdot 07$ $- \cdot 01$
	9 9 11 11 11	AP	9 23 8 51 9 23 10 4	" 22 46 "	$+ \cdot 03$ + $\cdot 04$ + $\cdot 06$ + $\cdot 06$	11 11 11	AP	16 24 14 11 14 46 16 24	2 30 I 27 ,,	+ <sup>.07</sup> + <sup>.06</sup> + <sup>.04</sup> + <sup>.12</sup>	I I I 2	S	19 51 19 55 20 7 20 16 19 46	7 35	$- \cdot 03$ $- \cdot 06$ $+ \cdot 02$ $- \cdot 14$
	12 12 13 13 16	AW	9 23 10 4 12 58 13 20 9 23	22 45 ,, ,, 21 28	+ •03 + •11 + •16 + •09 + •06	1911. Jan. 17 17 22 22	JJ AP	18 0 18 49 17 31 17 39	3 50 ,,, 5 43	+ .02 + .00 + .00	Apr. 2 2 6 7 10	RC C JJ	2 I 42 22 I 22 53 2 58 22 53	8 29 ,,, 7 54 ,, 7 52	·00 + ·02 - ·10 - ·05 - ·06
	16 18 18 19	C AP	10 4 9 23 10 4 9 23 10 4	23 <sup>"18</sup> "22 <sup>"40</sup>	+ ·06 + ·01 + ·01 + ·01 + ·01	23 23 23 24 25	AW	18 15 18 22 18 50 21 40 21 40	5 39	·00 ·00 + ·04 - ·04 - ·10	11 11 12 24 28	C AW C	2 58 22 53 2 58 4 31 0 <b>3</b> 9	7 58	+ ·03 - ·03 - ·03 + ·07 - ·04
	-		.				3								

C. F. C., 1900.

3

	Date	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day— Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.	Date.	Observer.	S.T. of Day Obs.	Mean S.T. of Night Obs.	Diff. Day- Night.
M	ay	 SolutionCJJAWJJRCAWAPCMJJAPCJJAPCJJAP					Image: Second system      AP      C      AW      JJ      C      C      AP      JJ      RC      AP      S      AW      M      RC      AW      C      RC      AW      M      RC      AW      C      RC      AW      C      RC      RC	h         m           h         m           8         4           5         10           5         20           7         40           5         20           7         40           5         20           7         40           5         20           7         40           5         20           9         23           10         4           9         23           10         4           9         23           10         4           9         23           10         4           9         23           10         4           5         50           6         12           11         9           12         11           15         11           16         33           6         55           7         5           11         15           12         11           12         11	Night Obs. h m 12 41 15 12 " " 14 53 " " 14 53 " " 15 29 " 15 33 " 15 29 " 15 33 " 15 33 " 15 33 " 15 57 16 " 0 " 15 49 16 29 " " 15 49 16 29 " " 15 49 16 50 " " 16 39 " " 17 41 17 58 17 29 " " 17 12 "	s         o	Ight, Aug. 30 30 30 31 31 Sept. 5 6 6 8 8 12 12 19 28 29 Oct. 2 2 2 2 4 6 6 6 9 20 20 20 20 20 24 25 31 Nov. 8 8 13 13 14 14 16 16 16 17 7 7 20 22 22 22 22 22 22 22 22 22 22 22 22	Image: second systemJJMAWSAWCAWRCJJMCJJMCJJSCAPJJRCAWJJRCAWJJJJRCAWJJRCAWJJ		Obs. h m 20 6 18 16 " 19 33 18 59 18 59 18 42 20 25 " 19 16 "	

TABLE VII.—continued.

The periodic character of the differences Day—Night is at once evident from inspection of this table. In order, however, to subject it to a closer analysis it has been assumed that it is primarily due to a periodic error in the clock-star system used, which may be expressed analytically by the formula

 $\Delta a = A_1 \cos a + B_1 \sin a + A_2 \cos 2a + B_2 \sin 2a.$ 

xviii

This has further been regarded as possibly associated with a diurnal periodicity, either due to different habits of the observers in daylight observing as contrasted with night observing, or to a diurnal change in the conditions of the transit circle or a diurnal period in the clock rate. Thus each of the differences Day—Night has been equated to an expression of the form

#### $K + A_1(\cos a_1 - \cos a_2) + B_1(\sin a_1 - \sin a_2) + A_2(\cos 2a_1 - \cos 2a_2) + B_2(\sin 2a_1 - \sin 2a_2),$

where  $a_1$  denotes the R.A. of the day star and  $a_2$  the mean R.A. of the night stars on which the comparison depends. The quantities  $a_1$ ,  $a_2$  are given under the headings S.T. (sidereal time) of *Day* Observations and Mean S.T. of *Night* Observations in columns 3 and 4 of Table VII. While the quantities  $A_1$ ,  $B_1$ ,  $A_2$ ,  $B_2$  have been regarded as constant throughout the series of observations, the quantity K has been considered as possibly variable with the observer or the method of observing. We give in Table VIII. partial normal equations obtained by grouping the observations according to the observer and the year of observation and combining the separate equations with equal weight.

## TABLE VIII.—Partial Normals for the Determination of Periodic Errors in Right Ascension.

					v.	υ'.
30 <i>K</i>	$- 6A_1 + 63$	$ \begin{array}{r}     14B_{1} \\     + 26 \\     + 44 \end{array} $	908. $- 14A_2$ - 16 - 18 + 22	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r}         s \\         + 0.19 \\         - 0.18 \\         + 0.36 \\         - 0.01 \\         - 0.47         $	$ \begin{array}{r} & & & \\ & + & \circ \cdot \circ 4 \\ & - & \circ \cdot \circ 2 \\ & + & \circ \cdot \circ 5 \\ & & \circ \cdot \circ 0 \\ & - & \circ \cdot 1 1 \end{array} $
55 <i>K</i>	$+ 5A_{1}$	$+ 10B_1$ - 10 87	909. $-$ $+$ $7A_2$ - $24+$ $44^{\circ}$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.12 \\ + & 1.55 \\ + & 0.39 \\ - & 0.10 \\ - & 0.69 \\ \end{array} $	$ \begin{array}{r} - & 0.02 \\ + & 0.15 \\ + & 0.04 \\ - & 0.02 \\ - & 0.11 \\ \end{array} $
20K	$- \frac{16A_1}{33}$	$\begin{array}{c} + 5B_1 \\ - 2 \\ 33 \end{array}$	$\begin{array}{c} + & 5A_2 \\ + & 6 \\ + & 3 \\ & 21 \end{array}$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} + & 0.05 \\ - & 0.11 \\ - & 0.01 \\ + & 0.09 \\ + & 0.03 \end{array} $	+ 0.01 - 0.02 0.00 + 0.02 + 0.01
31 <i>K</i>	$+ 5A_{1}$ 43	$+ 23B_1 \\ - 2 \\ 57$	$\begin{array}{r} - & 8A_2 \\ - & 4 \\ - & 26 \\ + & 24 \end{array}$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} + 0.02 \\ + 0.07 \\ + 0.16 \\ - 0.01 \\ - 0.61 \\ \end{array} $	0.00 + 0.01 + 0.02 - 0.00 - 0.11

Observer C.

## TABLE VIII.—continued.

Observer AP.

			and the second	v.	<i>v</i> .'
30 <i>K</i>	- 3A <sub>1</sub> 44	$ \begin{array}{rcrcrcr} & & & & & & & & & \\ & + & & & & & & & & $	$ \begin{array}{rcrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r}                                     $	$ \begin{array}{r} 8 \\ - 0.09 \\ + 0.07 \\ + 0.02 \\ - 0.01 \\ - 0.07 \\ \end{array} $
49 <i>K</i>	+ 20A <sub>1</sub> 92	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.56 \\ - & 1.06 \\ + & 0.37 \\ + & 0.15 \\ + & 0.06 \end{array} $	- 0.08 - 0.11 + 0.04 + 0.04 + 0.01
47 <i>K</i>	- 34 <i>A</i> 1 59	$ \begin{array}{r}     1910. \\     + 17B_1 - 27A_2 \\     - 12 - 7 \\     79 - 29 \\     71 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	+ 1.04  - 1.55  + 0.72  - 1.12  + 0.27	$\begin{array}{r} + & 0.12 \\ - & 0.20 \\ + & 0.08 \\ - & 0.13 \\ + & 0.05 \end{array}$
24 <i>K</i>	$+ \frac{1A_1}{17}$	$ \begin{array}{r}     1911. \\     + 8B_1 - 11A_2 \\     + 5 - 3 \\     58 - 25 \\     22 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} 0.00 \\ - 0.15 \\ - 0.17 \\ - 0.24 \\ - 0.15 \end{array} $	0.00 - 0.04 - 0.02 - 0.05 - 0.03

)	bs	er	v	e	r	F	2	C	
	NO	GT	v	C	L	L	6		

and the second se					
18 <i>K</i>	$-3A_{I}$ + 26	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.22 \\ - & 0.28 \\ - & 0.56 \\ + & 0.17 \\ + & 0.26 \end{array} $	- 0.05 - 0.06 - 0.09 + 0.05 + 0.08
29 <i>K</i>	- 6A <sub>1</sub> 31	$ \begin{array}{r}     1909. \\ + 10B_1 + 6A_2 \\ - 8 - 4 \\ 60 + 10 \\ 27 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.20 \\ + & 0.38 \\ + & 0.15 \\ + & 0.05 \\ + & 0.19 \\ \end{array} $	$ \begin{array}{r} - & 0.04 \\ + & 0.07 \\ + & 0.02 \\ + & 0.01 \\ + & 0.04 \\ \end{array} $
38 <i>K</i>	+ 5A1 39	$ \begin{array}{r}     1910. \\     + 5B_1 - 7A_2 \\     + 1 - 8 \\     45 + 9 \\     62 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} + & 0.45 \\ - & 0.99 \\ - & 0.34 \\ - & 0.31 \\ + & 0.01 \end{array} $	+ 0.07 - 0.16 - 0.05 - 0.04 0.00
22 <i>K</i>	- 8A <sub>1</sub> 23	$ \begin{array}{rcrr}  & 1911. \\ - & 10B_1 & + & 18A_2 \\ - & 2 & - & 21 \\  & 40 & - & 2 \\  & & & 32 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	0.00 - 0.06 - 0.08 + 0.04 + 0.08	0.00 - 0.01 - 0.01 + 0.01 + 0.02

xx

## TABLE VIII.—continued.

Observer AW.

	12182			v.	<i>v</i> .'
41 <i>K</i>	$+ \frac{5A_1}{63}$	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r}                                     $	8 - 0'10 + 0'01 - 0'10 + 0'06 + 0'06
		1909		-	
34 <i>K</i>	$+ {}^{14A_1}_{67}$	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.07 \\ - & 0.23 \\ + & 0.02 \\ & 0.00 \\ + & 0.57 \end{array} $	- 0.01 - 0.03 0.00 + 0.13
54 <i>K</i>	$- \frac{19A_1}{73}$	$ \begin{array}{r} 1910 \\ + 21B_1 & - 20A_2 \\ - 25 & + 18 \\ 76 & - 10 \\ 91 \\ \end{array} $	$- IB_2 = + 2.48$ - I8 = - 2.23 + 43 = + 2.24 - 8 = - I.63 55 = + 0.86	$ \begin{array}{r} + & 0.72 \\ - & 0.66 \\ + & 0.36 \\ + & 0.01 \\ - & 0.02 \\ \end{array} $	+ 0.10 - 0.08 + 0.04 0.00 0.00
49 <i>K</i>	+ 4A <sub>1</sub> 55	$ \begin{array}{r}     1911 \\     + \circ B_1 + 16A_2 \\     - 26 - 25 \\     99 + 9 \\     49 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} 0.00 \\ - 0.36 \\ + 0.39 \\ + 0.21 \\ + 0.49 \end{array} $	0.00 - 0.02 + 0.04 + 0.03 + 0.08

Observer M	
------------	--

the second se					
35 <i>K</i>	+ 8 <i>A</i> 1 45	$ \begin{array}{rcrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{vmatrix} + & 0.09 \\ + & 0.56 \\ - & 0.10 \\ - & 0.52 \\ + & 0.07 \end{vmatrix} $	+ 0'02 + 0'09 - 0'01 - 0'09 + 0'02
16 <i>K</i>	- 5 <i>A</i> 1 14	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.01 \\ + & 0.09 \\ + & 0.03 \\ + & 0.54 \\ + & 0.02 \\ \end{array} $	$ \begin{array}{r} 0.00 \\ - 0.02 \\ + 0.01 \\ + 0.13 \\ + 0.01 \end{array} $
39 <i>K</i>	+ 5 <i>A</i> <sub>I</sub> 54	$ \begin{array}{r}     1910. \\ + 7B_1 + 3A_2 \\ - 4 + 2 \\ 50 + 20 \\ 37 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.07 \\ + & 0.16 \\ + & 0.33 \\ + & 0.26 \\ + & 0.43 \end{array} $	$ \begin{array}{r} - & 0.01 \\ + & 0.02 \\ + & 0.05 \\ + & 0.04 \\ + & 0.05 \end{array} $
24 <i>K</i>	- 9 <i>A</i> <sub>1</sub> 25	$ \begin{array}{rcrr}  & & & & & & & & & & & & & & & & & & &$	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.02 \\ + & 0.16 \\ + & 0.09 \\ + & 0.08 \\ - & 0.40 \end{array} $	$ \begin{array}{r} 0.00 \\ + 0.03 \\ + 0.01 \\ + 0.02 \\ - 0.09 \end{array} $

## TABLE VIII.—continued.

Observer JJ.

1					
				<i>v</i> .	<i>v</i> ′.
27 <i>K</i>	- 12 <i>A</i> <sub>1</sub> 49	$ \begin{array}{r}     1908 \\     + 28B_1 & - 7A_2 \\     - 2 & - 13 \\     53 & - 7 \\     14 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r}                                     $	8 0.00 + 0.04 - 0.04 - 0.05 - 0.01
54 <i>K</i>	$+ 9A_{1} 67$	$   \begin{array}{r}     1909 \\     + 20B_1 - 4A_2 \\     + 12 - 12 \\     106 0 \\     33   \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - & 0.15 \\ - & 0.08 \\ - & 0.27 \\ + & 0.32 \\ & 00 \end{array} $	$ \begin{array}{r} - & 0.02 \\ - & 0.01 \\ - & 0.03 \\ + & 0.06 \\ 0.00 \end{array} $
19 <i>K</i>	+ $\frac{6A_1}{18}$	$ \begin{array}{r}     1910 \\ - 18B_1 - 3A_2 \\ + 8 - 1 \\     40 + 14 \\     31 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} + & 0.21 \\ + & 0.16 \\ + & 0.08 \\ + & 0.41 \\ - & 0.09 \\ \end{array} $	$\begin{array}{r} + & 0.05 \\ + & 0.04 \\ + & 0.01 \\ + & 0.08 \\ - & 0.03 \end{array}$
37 <i>K</i>	- 1 <i>A</i> 1 40	$ \begin{array}{r}     1911 \\     - 12B_1 + 7A_2 \\     - 18 - 9 \\     72 - 9 \\     32 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{c} + & 0.01 \\ + & 0.16 \\ - & 1.21 \\ + & 0.04 \\ + & 0.17 \end{array} $	0.00 + 0.03 - 0.14 + 0.01 + 0.03

Observer JW.

	19 <i>K</i>	- 10A <sub>1</sub> 35	$ \begin{array}{rcrcrcr}  & & & & & & & \\ & + & 8B_1 & - & & & & \\ & + & 13 & - & 2 \\ & + & 13 & - & 2 \\ & 40 & - & 0 \\ & & & & & & & \\ & & & & & & & 2 \end{array} $	$ \begin{array}{rcl} - & 1B_2 &= & 0.00 \\ - & 1 &= & + & 0.49 \\ - & 3 &= & + & 0.36 \\ - & 1 &= & - & 0.03 \\ & 6 &= & - & 0.02 \end{array} $	- 0'05 + 0'48 - 0'04 - 0'06 - 0'01	- 0.01 + 0.08 - 0.01 - 0.04 0.00
A DELLARD	23 <i>K</i>	+ 24A <sub>1</sub> 37	$ \begin{array}{rcrcrcr}  & & & & & & & & & & & & & & & & & & &$	$ \begin{array}{rcrcrc} - & 6B_2 &= & - & 0.41 \\ + & 1 &= & - & 0.37 \\ + & 1 &= & + & 1.01 \\ + & 7 &= & + & 0.11 \\ & 9 &= & + & 0.11 \end{array} $	$ \begin{array}{r} + \ 0.03 \\ + \ 0.24 \\ + \ 0.16 \\ + \ 0.29 \\ + \ 0.11 \end{array} $	+ 0.01 + 0.04 + 0.05 + 0.06 + 0.06
	35 <i>K</i>	- 12A <sub>1</sub> 64	$ \begin{array}{r}     1910. \\     - 15B_1 + 11A_2 \\     + 30 + 2 \\     36 + 9 \\     37 \end{array} $	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$ \begin{array}{r} - 0.07 \\ + 0.04 \\ - 0.11 \\ - 0.14 \\ + 0.10 \\ \end{array} $	$ \begin{array}{r} - & 0.01 \\ + & 0.01 \\ - & 0.02 \\ - & 0.02 \\ + & 0.02 \\ \end{array} $

#### TABLE VIII.—continued.

Observer S.

			v.	υ'.
$7K - \frac{1}{8}A_1$	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$- 4B_2 = - 0.08$ $0 = + 0.44$ $- 5 = + 0.45$ $- 1 = - 0.32$ $5 = - 0.17$	$ \begin{array}{r} 8 \\ + 0.01 \\ + 0.49 \\ + 0.35 \\ - 0.25 \\ - 0.21 \end{array} $	s 0°00 + 0°17 + 0°10 - 0°07 - 0°09

For the further combination of these equations the observations in the different years were first treated independently. By means of the normal in K, the quantity K was first eliminated, and reduced partial normals in  $A_1$ ,  $B_1$ ,  $A_2$ ,  $B_2$  were derived. The reduced partial normals for the separate observers were then combined by addition and the resulting complete normals solved. The results for the separate years are as follows:—

TABLE IX.—Coefficients of Periodic Errors in the Clock-Star System.

	<i>A</i> <sub>1</sub> .	<i>B</i> <sub>1</sub> .	A 2.	B <sub>2</sub> .
1908 1909	8 -0.0057 + .0020	s +0.0122 + .0177	8 	s -0.0046 + .0047
1910 1911	- ·0149 - ·0090	+ .0165 + .0122	- ·0103 - ·0148	+ •0010 - •0069

The observations made by each observer during the years 1908-10 were next regarded as furnishing homogeneous groups, which were combined among themselves in like manner, those of 1911 however being excluded, as a different method of observing was used in this year. The results from the separate groups are as follows :---

 TABLE IXA.—Coefficients of Periodic Errors in Clock-Star System (1908–10)

 grouped according to Observers.

Observer.	A 1.	<i>B</i> <sub>1</sub> .	A <sub>2</sub> ,	B <sub>2</sub> .
С	s -0.0013	8 +0.0181	s -0.0124	s -0'0237
AP	- '0231	+ .0176	- '0234	- '0079
RC AW	- '0172 - '0095	+ .0080	- 0137 - 0082	+ '0058
M	.0000	+ '0164	0052	+ .0066
JJ JW	- ·0037 - ·0026	+ '0111 + '0127	- °0003 - °0086	0024 + .0053
	S			

According to either method of grouping, the values of the quantities  $A_1$ ,  $B_1$ ,  $A_2$ ,  $B_2$  appear to be persistent, indicating real periodic errors in the Cape Ledger system. The definitive values have been derived by combining by addition all the reduced partial normals  $A_1$ ,  $B_1$ ,  $A_2$ ,  $B_2$ , which result after the elimination of K from each homogeneous group. The final complete normals are as follows:—

with the solution

 $A_1 = -0.0085 , \text{ weight } 1183, \\ B_1 = +0.0148 , ..., 1486, \\ A_2 = -0.0103 , ..., 741, \\ B_2 = +0.0006 , ..., 663.$ 

If we substitute these values in the respective partial normals in K, we derive the following values for K, which represent the personal discordances in time determinations by day as compared with those of the mean observer by *night*.

TABLE X.—Discon	rdances between Da	ay and Night Dete	erminations of Clo	ck Error
(in s	sense Day-Night),	grouped according	g to Observers.	

Day Observer.		Year of Observation.			
Duy Obbirron.	1908. 1909.		1910.	1908-10.	1911.
C AP RC AW M JJ JW S		$ \begin{array}{r}         8 \\         0'000 \\         - '021 \\         - '002 \\         + '018 \\         - 023 \\         + '009 \\         - '009 \\          $			
Mean	0026	0040	+ .0026	0004	+ .0110

The quantities here derived, except in so far as they are due to purely accidental causes, may be attributed in part to personal and partly to instrumental causes. If we give equal weight to the determinations in each of the four years involved, the mean result derived from all the observations amounts only to +0.80023 for the mean observer. It follows that there can be little or no danger of the determinations of the periodic errors in R.A. being vitiated by periodic errors due to diurnal changes in the instrument or the clock.

xxiv

The quantities contained in the two final columns of Table X. have been adopted as definitive, and, together with the finally derived values of  $A_1$ ,  $B_1$ ,  $A_2$ ,  $B_3$ , have been substituted in the original equations of condition. From the sum of the squares of the residuals thus formed the probable accidental error corresponding to weight unity has been derived as  $\pm 0^{\circ} \cdot 031$ ; whence, with the weights derived, the probable accidental errors of  $A_1, B_1, A_2, B_2$ , amount to  $\pm 0^{\circ} \cdot 0009, \pm 0^{\circ} \cdot 0008, \pm 0^{\circ} \cdot 0011, \pm 0^{\circ} \cdot 0012$ . The agreement between the derived values of these same quantities from the groups of observations, either arranged according to time or according to the observers, does not confirm this high estimate of the precision, doubtless on account of cumulative systematic errors. To obtain a more reliable estimate of the probable errors, both accidental and systematic, of the results, the derived values have been substituted in the partial normals (Table VIII.); the residuals are given in the last column but one of this Table. Now it is evident that if any one of these partial normals be written in the symbolical form

 $(aa)x + (ab)y + (ac)z + \ldots = (an),$ 

where each of the quantities n is of weight unity, the square of the mean error of the absolute term will be  $(\alpha \alpha)\epsilon^2$ ,  $\epsilon$  denoting the mean error corresponding to unit weight.

Hence we may reduce the equations to equal weight unity by multiplying by the factor  $1/\sqrt{\alpha\alpha}$ . The final column in Table VIII. gives the residuals from the equations thus reduced.

Now if the quantities thus obtained represented true errors, instead of residual phenomena, since each has the same weight unity, the mean of their squares would give a determination of the square of the mean error for unit weight, but in that the derived phenomena depend on the equations themselves, we may anticipate that the average residual will be less than the average error.

On the other hand, the sum of the squares of the residuals will exceed that which would be derived from a least square combination of the partial normal equations regarded as equations of condition. But, according to the usual conventions of least squares, if  $\Sigma v^2$  denote the sum of the squares of the residuals, m the number of equations, and n the number of unknown quantities,

 $m\epsilon^2 = \Sigma v^2 + n\epsilon^2.$ 

Hence if v' denote residuals from a solution other than a least square solution,

 $m\epsilon^2 < \Sigma v'^2 + n\epsilon^2$ .

Applying this formula to the present case, a superior limit to the probable error corresponding to unit weight is found to be  $\pm 0^{\circ}.043$  and the corresponding probable errors of  $A_1$ ,  $B_1$  do not exceed  $\pm 0^{\circ} \cdot 0013$ , those of  $A_2$ ,  $B_2$ ,  $\pm 0^{\circ} \cdot 0018$ . 4

C. F. C., 1900.

On the basis of this determination the probable error, inclusive of residual systematic error, as well as purely accidental error of the quantity

#### $A_1 \cos \alpha + B_1 \sin \alpha + A_2 \cos 2\alpha + B_2 \sin 2\alpha$ ,

amounts at a maximum in any right ascension to  $\pm 0^{8} \cdot 0022$ .

As regards the actual values derived for the coefficients  $A_1$ ,  $B_1$ ,  $A_2$ ,  $B_2$ , confirmation has been sought from comparison with approximately simultaneous series of observations made in other observatories, with results that support the values here derived (see *Monthly Notices*, January 1913). For the purposes of the present Catalogue it has, however, been thought desirable, in order to maintain its fundamental character, to avoid the introduction of extraneous evidence.

Thus the definitive corrections which have been applied to the Ledger right ascensions in order to eliminate the errors in the system of right ascension originally adopted for their formation are

 $\Delta a = +0^{s} \cdot 0085 \cos a - 0^{s} \cdot 0148 \sin a + 0^{s} \cdot 0103 \cos 2a - 0^{s} \cdot 0006 \sin 2a.$ 

For reasons which will be discussed later no constant correction has been applied. Thus the equinox of reference corresponds with that of Newcomb's Catalogue.

#### II.—REVISION OF DECLINATION SYSTEM.

The declinations in the Ledgers have been derived from the nadir readings, with the Pulkowa refractions and with an assumed value for the mean latitude of the transit circle, viz. :--

### -33° 56′ 2″.5.

Except for the year 1911, they have received corrections on account of the motion of the Earth's axis from data supplied by Albrecht from the latitude determinations at the International Geodetic Stations. The same applies to the time stars of 1911, but not to the circumpolar stars, the observations of which are contained in a separate ledger, and which have formed the subject of a special discussion (*Cape Annals*, vol. xi., part iii.). No corrections for instrumental flexure have been applied prior to the formation of the Ledgers.

Before considering the corrections on account of latitude and flexure, a comparison was first made between the results derived in the four conditions of the instrument I. E., I. W., II. E., II. W. A summary of this comparison, based on observations during the years 1905-10, is given in the following tables.

xxvi

.

INTRODUCTION.

		and the second se		(2	,		
Mean Dec.	0 <sup>h</sup> -4 <sup>h</sup> .	4 <sup>h</sup> -8 <sup>h</sup> .	8 <sup>h</sup> -12 <sup>h</sup> .	12 <sup>h</sup> -16 <sup>h</sup> .	16 <sup>h</sup> -20 <sup>h</sup> .	20h_oh.	Mean.
$\begin{array}{r} + 27 \\ + 15 \\ + 5 \\ - 5 \\ - 25 \\ - 35 \\ - 45 \\ - 55 \\ - 75 \\ - 85 \\ - 95 \\ - 105 \end{array}$	$\begin{array}{r} & & & & & & \\ & + & \circ^{\circ}17_{34} \\ & + & \circ^{\circ}26_{16} \\ & + & \circ^{\circ}37_{27} \\ & + & \circ^{\circ}43_{16} \\ & + & \circ^{\circ}38_{10} \\ & + & \circ^{\circ}35_{16} \\ & + & \circ^{\circ}27_{13} \\ & + & \circ^{\circ}34_{26} \\ & + & \circ^{\circ}25_{13} \\ & + & \circ^{\circ}46_{13} \\ & + & \circ^{\circ}62_{5} \end{array}$	$\begin{array}{r} & & & & & & \\ & + & & & & & \\ & + & & & &$	$\begin{array}{r} + & 0^{\circ} \cdot 1 \\ + & 0^{\circ} \cdot 3 \\ 0^{\circ} \cdot 3 \\ + & 0^{\circ} \cdot 3 \\ 0^{\circ} \cdot 2 \\ + & 0^{\circ} \cdot 3 \\ 2 \\ + & 0^{\circ} \cdot 3 \\ 1 \\ + & 0^{\circ} \cdot 5 \\ 1 \\ - & 0^{\circ} \\ 1 \\ - & 0^{$	$\begin{array}{c} - & 0.02 & 10 \\ + & 0.11 & 17 \\ - & 0.01 & 11 \\ - & 0.01 & 17 \\ + & 0.16 & 22 \\ + & 0.17 & 15 \\ + & 0.22 & 14 \\ + & 0.24 & 31 \\ + & 0.39 & 14 \\ + & 0.36 & 7 \\ + & 0.16 & 8 \\ + & 0.43 & 4 \\ + & 0.07 & 4 \\ + & 0.40 & 5 \end{array}$	$\begin{array}{r} + 0.11 \\ + 0.16 \\ 20 \\ + 0.29 \\ 18 \\ + 0.06 \\ 12 \\ + 0.26 \\ 13 \\ + 0.27 \\ 24 \\ + 0.28 \\ 16 \\ + 0.28 \\ 16 \\ + 0.28 \\ 16 \\ + 0.28 \\ 16 \\ + 0.28 \\ 16 \\ + 0.28 \\ 16 \\ + 0.27 \\ 12 \\ + 0.57 \\ 10 \\ - 0.02 \\ 6 \\ + 0.07 \\ 4 \\ - 0.07 \\ 5 \\ + 0.37 \\ 1 \end{array}$	$\begin{array}{r} + \circ^{''}_{24_{25}} \\ + \circ^{''}_{38_{14}} \\ + \circ^{'}_{36_{18}} \\ + \circ^{'}_{43_{24}} \\ + \circ^{'}_{34_{20}} \\ + \circ^{'}_{46_{19}} \\ + \circ^{'}_{46_{19}} \\ + \circ^{'}_{48_{18}} \\ + \circ^{'}_{30_{14}} \\ + \circ^{'}_{51_{12}} \\ + \circ^{'}_{47} \\ 7 \\ + \circ^{'}_{34_{4}} \\ \circ^{'}_{000_{2}} \\ + \circ^{'}_{53_{4}} \\ \end{array}$	$\begin{array}{r} & & & & & & & & & \\ & + & 0.196_{154} \\ & + & 0.267_{110} \\ & + & 0.325_{116} \\ & + & 0.300_{101} \\ & + & 0.303_{91} \\ & + & 0.378_{102} \\ & + & 0.378_{102} \\ & + & 0.396_{154} \\ & + & 0.368_{88} \\ & + & 0.479_{61} \\ & + & 0.364_{49} \\ & + & 0.359_{24} \\ & + & 0.145_{22} \\ & + & 0.578_{19} \end{array}$
		Р	osition II.	$\Delta\delta (E - V)$	V).		
$\begin{array}{r} + & 27 \\ + & 15 \\ + & 5 \\ - & 55 \\ - & 15 \\ - & 25 \\ - & 35 \\ - & 35 \\ - & 45 \\ - & 45 \\ - & 55 \\ - & 75 \\ - & 85 \\ - & 95 \\ - & 105 \end{array}$	$+ 0.48 \\ + 0.65 \\ 16 \\ + 0.52 \\ 27 \\ + 0.63 \\ 16 \\ + 0.49 \\ 10 \\ + 0.62 \\ 16 \\ + 0.43 \\ 13 \\ + 0.43 \\ 13 \\ + 0.43 \\ 13 \\ + 0.33 \\ 13 \\ + 0.39 \\ 11 \\ + 0.44 \\ 4 \\ + 0.64 \\ 4 \\ + 0.48 \\ 5 \end{bmatrix}$	$\begin{array}{r} + \circ \cdot 45_{27} \\ + \circ \cdot 53_{21} \\ + \circ \cdot 27_{17} \\ + \circ \cdot 39_{20} \\ + \circ \cdot 44_{14} \\ + \circ \cdot 38_{19} \\ + \circ \cdot 49_{16} \\ + \circ \cdot 52_{24} \\ + \circ \cdot 34_{16} \\ + \circ \cdot 21_{17} \\ + \circ \cdot 32_{16} \\ + \circ \cdot 21_{17} \\ + \circ \cdot 32_{16} \\ + \circ \cdot 21_{17} \\ + \circ \cdot 2$	$\begin{array}{r} + \circ \cdot 19_{18} \\ + \circ \cdot 33_{22} \\ + \circ \cdot 43_{25} \\ + \circ \cdot 44_{13} \\ + \circ \cdot 39_{11} \\ + \circ \cdot 41_{9} \\ + \circ \cdot 34_{10} \\ + \circ \cdot 23_{10} \\ + \circ \cdot 23_{10} \\ + \circ \cdot 31_{12} \\ - \circ \cdot 66_{8} \\ + \circ \cdot 19_{3} \\ + \circ \cdot 20_{3} \\ + \circ \cdot 24_{2} \end{array}$	$\begin{array}{r} + \circ \cdot 12_{19} \\ + \circ \cdot 27_{17} \\ + \circ \cdot 30_{11} \\ + \circ \cdot 20_{17} \\ + \circ \cdot 33_{22} \\ + \circ \cdot 36_{15} \\ + \circ \cdot 33_{14} \\ + \circ \cdot 19_{14} \\ + \circ \cdot 19_{14} \\ + \circ \cdot 26_{17} \\ + \circ \cdot 19_{18} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ + \circ \cdot 19_{19} \\ + \circ \cdot 35_{18} \\ - \circ \cdot 19_{18} \\ + \circ \cdot 35_{18} \\ - \circ \cdot 19_{18} \\ + \circ \cdot 19_{18} \\ - \circ \cdot 19_{18} \\ + \circ \cdot 19_{18} \\ - \circ \cdot 10_{18} \\ - \circ \cdot 1$	$\begin{array}{r} + \circ \cdot 17_{31} \\ + \circ \cdot 24_{20} \\ + \circ \cdot 4\circ 18 \\ + \circ \cdot 39_{12} \\ + \circ \cdot 47_{13} \\ + \circ \cdot 45_{24} \\ + \circ \cdot 37_{16} \\ + \circ \cdot 37_{26} \\ + \circ \cdot 37_{29} \\ + \circ \cdot 17_{10} \\ + \circ \cdot 29_{12} \\ + \circ \cdot 17_{10} \\ + \circ \cdot 23_{6} \\ - \circ \cdot 14_{3} \\ + \circ \cdot 19_{3} \\ + \circ \cdot 39_{1} \end{array}$	$\begin{array}{r} + 0.21_{25} \\ + 0.24_{14} \\ + 0.51_{19} \\ + 0.37_{24} \\ + 0.46_{20} \\ + 0.62_{18} \\ + 0.47_{15} \\ + 0.31_{17} \\ + 0.30_{14} \\ + 0.30_{12} \\ + 0.16 \\ 7^{\circ} \\ + 0.17 \\ 4 \\ + 0.10 \\ 3 \\ + 0.02 \\ 3 \end{array}$	$\begin{array}{r} + \circ 291_{154} \\ + \circ 378_{110} \\ + \circ 424_{117} \\ + \circ 398_{102} \\ + \circ 421_{90} \\ + \circ 427_{101} \\ + \circ 346_{153} \\ + \circ 366_{88} \\ + \circ 278_{61} \\ + \circ 278_{61} \\ + \circ 219_{49} \\ + \circ 234_{23} \\ + \circ 247_{20} \\ + \circ 377_{18} \end{array}$

TABLE XI.—Comparison of Declinations with opposite positions of the Clamp. Position I.  $\Delta\delta$  (E – W).

Comparison of Declinations with reversed positions of Object Glass and Eye-End.

$\Delta \sigma$ (1 ostion 1.— i ostion 11.; mean of E and w.)										
$\begin{array}{c ccccc} + & 27 & - & 0.1 \\ + & 15 & - & 0.1 \\ + & 5 & - & 0.1 \\ + & 5 & - & 0.1 \\ - & 5 & - & 0.0 \\ - & 15 & - & 0.1 \\ - & 25 & + & 0.1 \\ - & 35 & + & 0.0 \\ - & 35 & + & 0.0 \\ - & 45 & + & 0.1 \\ - & 55 & + & 0.2 \\ - & 65 & + & 0.2 \\ - & 65 & + & 0.2 \\ - & 75 & + & 0.2 \\ - & 85 & + & 0.3 \\ - & 95 & + & 0.1 \\ - & 105 & + & 0.7 \\ \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$ \begin{array}{c} - 0.31_{18} \\ - 0.23_{22} \\ - 0.08_{25} \\ - 0.01_{13} \\ + 0.01_{11} \\ + 0.06_{9} \\ + 0.02_{10} \\ + 0.10_{27} \\ + 0.13_{19} \\ + 0.18_{12} \\ + 0.14_{8} \\ + 0.29_{3} \\ + 0.46_{3} \\ + 0.22_{2} \end{array} $	$\begin{array}{c} - 0.29_{19} \\ - 0.28_{17} \\ 0.00111 \\ - 0.08_{17} \\ - 0.01_{22} \\ + 0.10_{15} \\ + 0.26_{14} \\ + 0.10_{31} \\ + 0.27_{14} \\ + 0.27_{7} \\ - 0.06_{8} \\ + 0.40_{4} \\ + 0.37_{4} \\ + 0.27_{5} \end{array}$	$\begin{array}{c} - 0.37_{31} \\ - 0.31_{20} \\ - 0.06_{18} \\ + 0.01_{12} \\ - 0.01_{13} \\ + 0.08_{24} \\ + 0.19_{16} \\ + 0.21_{28} \\ + 0.15_{12} \\ + 0.32_{10} \\ + 0.17_{16} \\ + 0.23_{3} \\ + 0.21_{3} \\ + 0.01_{1} \end{array}$	$\begin{array}{c c} - & 0.222_{25} \\ - & 0.24_{15} \\ - & 0.15_{18} \\ - & 0.07_{23} \\ - & 0.07_{20} \\ - & 0.01_{19} \\ & 0.00_{14} \\ + & 0.01_{19} \\ + & 0.17_{13} \\ + & 0.10_{12} \\ + & 0.20_{7} \\ + & 0.20_{7} \\ + & 0.20_{2} \\ + & 0.32_{3} \end{array}$	$\begin{array}{r} - 0.235_{154} \\ - 0.250_{111} \\ - 0.070_{116} \\ - 0.024_{101} \\ - 0.027_{90} \\ + 0.080_{102} \\ + 0.112_{155} \\ + 0.112_{155} \\ + 0.112_{155} \\ + 0.158_{87} \\ + 0.212_{61} \\ + 0.158_{49} \\ + 0.320_{24} \\ + 0.250_{19} \\ + 0.394_{18} \end{array}$				

 $\Delta\delta$  (Position I.—Position II. : mean of E and W.)

The suffixes indicate the number of stars in the group.

Fairly pronounced discordances of a systematic character depending on the zenith distance are clearly indicated. These may be in part accounted for by residual division-errors and by the variations in flexure under the different conditions. The separate determinations of the flexure coefficient by means of the horizontal collimators are given in the Introduction to the *Meridian Observations*. A summary of these is here given :--

V	1		11.			
Year.	E.	W.	E.	W.		
1905			+ 0.288	+ 0.363		
1906	+ 0.253	+ 0.310				
1907	+ 0.459	+ 0.335	60			
1908			+ 0.213	+ 0.132		
1909	+ 0'271	+ 0.323				
1910	+ 0.305	+ 0.384	+ 0.001	+ 0.068		
1911	+ 0.298	+ 0.384				
Mean	+ 0.317	+ 0.342	+ 0.187	+ 0.189		

TABLE XII.—Determinations of Mean Flexure Coefficient.

Within the limits of accidental errors of determination these figures indicate no appreciable change due to reversal between the two clamps, but a strongly marked difference between determinations in Positions I. and II. Accordingly the differences E - W, as given above in Table XI., after being smoothed by graphical interpolation, have been adopted as definitive.

To the difference II-I a correction on account of variation in the flexure coefficient, amounting to  $-0'' \cdot 14 \sin \zeta$ , where  $\zeta$  denotes the zenith distance, has been applied, and the results then smoothed in like manner.

Denoting the semi-differences  $\frac{1}{2}(II-I)$  by A, and the semi-difference  $\frac{1}{2}(E-W)$  by  $B_I$  or  $B_{II}$ , the following table gives the smoothed values for these quantities which have been used :—

Dec.	<i>A</i> .	B <sub>L</sub>	B <sub>11.</sub>	Dec.	A.	$B_{\rm I}$	B <sub>II.</sub>	Dec.	А.	B <sub>I</sub> .	$B_{\rm II.}$
$ \begin{array}{r} + 35 \\ 30 \\ 25 \\ 20 \\ 15 \\ 10 \\ + 5 \\ - 5 \\ 10 \\ \end{array} $	$\begin{array}{r} +0.07 \\ +0.07 \\ +0.05 \\ +0.05 \\ +0.03 \\ +0.01 \\ -0.02 \\ -0.03 \\ -0.03 \end{array}$	$\begin{array}{r} -0.08 \\ -1.0 \\ -1.12 \\ -1.12 \\ -1.13 \\ -1.14 \\ -1.15 \\ -1.15 \\ -1.15 \\ -1.15 \\ -1.15 \\ -1.15 \end{array}$	$ \begin{array}{r} -0.12 \\ -14 \\ -16 \\ -18 \\ -19 \\ -20 \\ -2$	- 15 20 25 30 35 40 45 55 55 60	- 0.04 - 0.04 - 0.05 -	$ \begin{array}{c} -0.16 \\ -16 \\ -17 \\ -17 \\ -17 \\ -17 \\ -17 \\ -17 \\ -18 \\ -19 \\ -20 \\ -20 \\ -20 \\ \end{array} $	16	80 85 90 S.P.85 ,, 80 ,, 75	$ \begin{array}{r} - \cdot 05 \\ - \cdot 05 \\ - \cdot 06 \\ - \cdot 06 \\ - \cdot 07 \\ + \cdot 08 \\ \end{array} $	$ \begin{array}{r} -0.21 \\ -20 \\ -10 \\ -10 \\ -118 \\ -118 \\ -117 \\ +17 \\ +17 \\ +17 \\ +18 \\ \end{array} $	$-\frac{0}{13} - \frac{12}{12} - \frac{11}{11} - \frac{11}{11} - \frac{11}{12} + \frac{13}{13} + \frac{14}{14} + \frac{16}{16}$

Table of Systematic Discordances.

xxviii

These corrections have to be applied to the observed declinations with the following signs in order to reduce the whole series to a uniform system :---

Position.	Clamp.	Δδ.
I.	E.	$A + B_{I.}$
I.	W.	$A - B_{I}$
II.	E.	$-A+B_{II.}$
II.	W.	$-A-B_{\rm IL}$

Consider next the latitude corrections. The separate observations have been reduced with the instantaneous nadir reading in combination with an assumed mean latitude and Albrecht's values for the periodic fluctuations of latitude, except in the case of close circumpolar stars observed during 1911. The latter have been separately discussed (*Cape Annals*, xi., part 3), the fluctuations of latitude being derived in this case from the observations themselves. From this discussion it appears that the latitude corrections required to reconcile the above-pole and below-pole observations at the Cape are less than those derived from observations at the International Latitude Stations by 0".18 in the mean, or, in other words, the adopted mean latitude used in the reductions requires to be diminished by 0".18. In deriving this value, however, no account was taken of the instrumental flexure. For the year in question the mean value of the flexure coefficient was +0".34, giving as the amount of flexure in the neighbourhood of the pole -0".28, in the sense in which it is to be applied to declination observations at upper culmination.

Taking

 $\Delta \delta = \Delta \phi + f \sin \zeta \text{ for stars above pole}$  $\Delta \delta = -\Delta \phi - f \sin \zeta \text{ for stars below pole}$ 

where  $\Delta\delta$  denotes the correction required to the declinations of the Ledgers,  $\Delta\phi$  the correction to the adopted latitude, and f the flexure coefficient, the above determinations give

whence

$\Delta \phi + f \sin \phi$	$\zeta = -0'' \cdot 18$	
$f \sin$	$\zeta = -0'' \cdot 28$	

 $\Delta \phi = + o'' \cdot 10.$ this year were all made wit

The observations during this year were all made with the transit circle in Position I. Now we have already seen that there are small systematic discordances between results obtained in Positions I. and II., amounting at the pole to  $+0''\cdot14$ , in the sense I-II. We may refer the latitude to the mean system  $\frac{1}{2}(I+II)$  by adding half this difference.

Thus the correction to the adopted latitude, suitable for application to determinations made by symmetrical observations in the two positions, as derived from the observations of circumpolars in the year 1911, is

 $\Delta \phi = + 0^{\prime\prime} \cdot 17.$ 

When the instrument was used in Position I., in the years 1906-10, direct determinations of flexure indicate that the mean flexure coefficient was sensibly constant. Hence for these years observations of the same star, made in this position of the instrument, have been treated as homogeneous and combined into a single mean. The determinations above and below pole have been thus separately grouped. The mean differences between the results for each star are contained in the following table :---

I OSICION I.											
	Clam	p E.	Clamp	• W.		Clam	р E.	Clamp W.			
Star.	Δδ Above —Below.	Weight.	Δδ Above Below.	Weight.	Star.	Δδ Above —Below.	Weight.	Δδ Above —Below.	Weight.		
ο Octantisβ Hydri Lacaille 505 τ <sup>1</sup> Hydri Lacaille 634 μ Hydri Lacaille 1029 Lacaille 2010 θ Mensæ Lacaille 2274 A Octantis θ Chamæleontis γ Chamæleontis γ Chamæleontis γ Octantis η Octantis η Octantis γ Octantis	$\begin{array}{c} - 0.07 \\ + 0.63 \\ + 1.02 \\ + 0.72 \\ + 1.20 \\ + 0.94 \\ + 0.94 \\ + 1.58 \\ + 0.27 \\ + 0.90 \\ + 0.30 \\ + 0.30 \\ + 0.33 \\ + 0.58 \\ + 0.11 \\ + 0.62 \\ + 1.10 \\ + 0.67 \\ + 0.52 \\ + 0.13 \\ + 0.58 \\ + 1.05 \\ + 0.58 \end{array}$	6 2 3 2 7 2 5 4 3 4 9 3 6 2 7 5 3 2 6 2 5 7	$\begin{array}{c} & & & & & & \\ & & & & & & \\ & & & & & $	7 2 3 2 6 2 6 5 2 3 10 2 7 2 4 2 3 3 2 2 8 2 7 5	β Chamæleontis ι Octantis κ Octantis θ Apodis α Apodis α Apodis α Octantis β Apodis γ Apodis γ Apodis γ Apodis γ Apodis γ Apodis γ Apodis γ Apodis γ Apodis γ Apodis μ Apodis α Octantis α Octantis μ Octantis μ Octantis μ Octantis β Octantis μ Octantis	$+ 1^{2}28 + 0^{4}1 + 0^{6}05 + 0^{4}3 + 0^{6}7 + 1^{6}02 - 0^{4}1 + 0^{6}7 + 1^{6}02 + 0^{6}7 + 1^{6}151 + 1^{6}51 + 1^{6}41 + 1^{6}51 + 1^{6}42$	2 8 7 2 6 2 6 2 9 3 9 5 5 8 2 2 3 3 7 7 2 2 2 3 3 7 7 2 2 2 9 3 9 5 5 5 8 2 2 2 9 3 9 5 5 5 8 7 2 2 2 6 2 2 9 3 9 5 5 5 7 2 2 2 2 9 3 9 5 5 5 7 2 2 2 9 3 9 5 5 5 5 5 5 5 5 5 5 5 7 2 2 2 5 5 5 5 5	$\begin{array}{c} & & & & & & \\ & & & & & & \\ & & & & & $	2 8 7 2 5 2 5 2 5 2 7 2 6 3 4 7 3 2 2 3 4 8 3 3 3		

TABLE	XIII.—Differences	between	Declinations	above	and	below	pole	in
	Ca	ape Ledg	yers (1906–10	).				

Position I.

The weights are derived from the formula

 $\frac{mn}{m+n}$ 

where m, n denote the number of observations made respectively at upper and lower culminations. Taking the means with these weights, we find, from Clamp E,

 $\Delta\delta$  (above - below) = + 0".81, weight 217;

and, from Clamp W,

 $\Delta\delta$  (above - below) = + 0"21, weight 192.

XXX

The difference between these results is in conformity with the discordances already found between declination determinations with reversed clamps. If we had previously applied the corrections represented by A,  $B_1$  above, the above determinations would have been increased respectively by the values of  $2(A \pm B_1)$  at the pole. The resulting corrections to the declinations on account of the combined effects of latitude and flexure would then be

 $-o'' \cdot \mathbf{i}o - (A + B_{I}) = -o'' \cdot \mathbf{i}6 \text{ for Clamp E.}$  $-o'' \cdot \mathbf{i}o - (A - B_{I}) = -o'' \cdot \mathbf{i}o \text{ for Clamp W.}$ 

These results are in reasonably close agreement.

Subtracting the part  $-0^{\prime\prime}$  28 due to flexure alone, we derive from the mean of the two latitude corrections referred to the mean system  $\left[\frac{1}{2}(I + II), \frac{1}{2}(E + W)\right]$ 

$$\Delta\phi = +0^{\prime\prime} \cdot 10,$$

The flexure determinations made in Position II. during the years 1905–10 show variations from year to year. Consequently for this position of the instrument a separate investigation on similar lines has been made for each year. Table XIV. gives the results derived from separate stars.

## TABLE XIV.—Differences between Declinations above and below pole in Cape Ledgers (1905–10).

		1905.				19	08.		1910.			
Star.	E.		w.		E.		W.		E.		W.	
	Δδ	Wt.	Δδ	Wt.	۵۵	Wt.	Δδ	Wt.	Δδ	Wt,	Δδ	Wt.
o Octantis	+ 1'20	1.0	+0'12	2.0	+0.32	2.0		1'2				
β Hydri	T120	19	T012	20	+0.32	1.0	-0.22	12			0.00	1.0
Lacaille 505	-0'25	0.8	-1'27	2.0				301				
$\tau^1$ Hydri Lacaille 634			-0.28	3.3	-0.50	1.7	-0.49	1.3	-0'20	0.8	•••	14
μ Hydri		-					- 1'02	1.2		1.4	- 1.00	0.2
Lacaille 1029 Lacaille 1848				2.7	-0.72	0.2	+0.12	1.3			-0.05	0.7
ι Hydri			+0.68	0.2	-0.33	1.3	-0.22	2.0	•••	2		0 /
Brisbane 593 Lacaille 1707		0.5	-0.44	1.0	+0.81	1.2	-0.64	2.0		_	•••	13.0
y Mensæ		20	-011	1.2	+0.01	3.4	-0.66 -0.87	3.2				
Lacaille 2296		0.9		111	+0.14	2.8	-0.09	1.3	• • • • •			50
к Mensæ Lacaille 2512					+0.33	1.2	+0.48	1.4 0.2	-0.11	0.7		
Lacaille 3274			-1.02	0.9	+0.50	1.7	-1'29	1.3	-0.02	1	-0.39	1.5
$\theta$ Chamæleontis $\eta$ Chamæleontis				Aile	+0.14	2·4 2·2	-0.38 +0.20	1.3	•••	Mei-	-0'21	0.2
7					_090	2.2	70.50	0.2			-021	

## Position II.

は見れたいのほど言語		1905.				1908.				1910.			
Star.	E.	E IT	W.		E.		w.		E.		W.		
	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	Δδ	Wt.	
ζ Octantis		3.4	- o	2.4	-0"29	2.7	-0.67	2'2	+0.53	0.2			
γ Chamæleontis			+0.68	2.8	-0°23	1.9	+0.45	0.7			-1.15	0.2	
Lacaille $4510$ $\eta$ Octantis		3.9	+0.32	2.9	+0.42	1.3	-0-72	2.2					
$\beta$ Chamæleontis		0.7		- 7	-0.06	2.0	-0.48	0.8				-	
i Octantis	+0.24	3.3	-0.40	3.8	+0.42	2.2	-0.60	2'4					
к Octantis	+0.34	3.7	-0.22	2.1		1			+0.28	0.2			
$\theta$ Apodis					-0.23	0.8	- 1.02	1.2				-	
a Apodis					•••		- 1.12	1.5	+0.19	0.8	+0.21	0.2	
ζ Apodis	+0.82	2.0	-0.44	0.8	+0.85		-0.50	1.2					
Lacaille $6077$ $\rho$ Octantis		1.0		60.1C	-0.08	0.8	-0'77	1.3	+1.42	1.5			
$\delta^1$ Apodis		1 2		01	+0.47	1.3	+0.02	2.0	T 1 4*	1.2			
$\gamma$ Apodis					+0.74	2.0	-1.38	0.8					
Lacaille 6545		1.5	+0.24	1.2	+0.22	2.5	-0.42	1.6	-0.66	1.0	+0.39	1'2	
$\beta$ Apodis	· · · · ·	-		-	+0.22	2.0	-0.70	1.7					
x Octantis					+0.44	2.3	-0.26	1.2					
Lacaille 8094			-0.09	0.0	-0.24	2.7	-0.89	1.5					
Lacaille 8257		2.0	•••		+0.42	2.9	+0.01	2.2	+0.41	0.2			
$\mu^1$ Octantis	•••	-			+0.11	1.9	-0.62	1.6	· ···			iente	
a Octantis		-	•••	1111	-0.05	2.2	-0.54	2°2					
$\nu$ Octantis		0.0		2:0	-0.28	1,3	-0.09	1.9	•••		•••		
v (C) Octantis $\tau$ Octantis		3.9	+0.14	3.0	-0.27	2.2	-0.43 -0.29	3.2	+ 1.68	0.7			
Lacaille 9494		3 2	-0.00	1.3	-0'25	2.2	-0.24	1.5	1100	0/		4	
$\theta$ Octantis		2.0	+0.16	2.0	+0.02	0.2	/4		+0.15	0.2	-0.20	1.0	
	- 74					- )				- )			

TABLE XIV.—continued.

whence we derive in the mean

 $\Delta\delta$  (above - below).

	Clamp E.	Weight.	Weight. Clamp W.		
1905	+0 <sup>"17</sup>	45	-0.12	40	
1908	+0°04	65	-0.49	62	
1910	+0°47	10	-0.27	8	

The differences E - W give in the mean the value

+0".47,

which corresponds very closely with the value of  $2B_{II}$  at the pole, as previously determined.

Applying the corrections  $-2(A \mp B_{II})$  respectively to results from Clamp E and Clamp W, we obtain the following values:—

	$\Delta\delta$ (above – below).											
			Clamp E.	Clamp W.								
1905		•	+0.01	+0.23								
1908	•	•	. – 0.06	-0.11								
1910	•	•	+0.32	+0.11								

		Clamp E.	Clamp W.
1905		 -0.04	-0'12
1908		 +0.03	+0.06
1910	?	 -0.18	-0.06

The parts of these quantities due to flexure alone are respectively

1905		 -0'27 .
1908		-0.14
1910		-0.02

whence the derived values for the latitude correction referred to the mean system are

	1	Clamp E.	Clamp W.
1905		+0.23	+ 0.12
1908		+0.12	+0.50
1910		 -0.13	-0.01

Collecting the various determinations, we find as the latitude correction referred to the homogeneous system,  $[\frac{1}{2}(1 + II) : \frac{1}{2}(E + W)]$ .

Period of Observations.	Position.	Clamp.	Δφ.	Weight.
1906-10 " 1905 " 1908 " 1910 " 1911	I. I. II. II. II. II. II. II. II. II. I	E W E W E W E and W	$ \begin{array}{r} + 0.12 \\ + 0.08 \\ + 0.23 \\ + 0.15 \\ + 0.17 \\ + 0.20 \\ - 0.13 \\ - 0.01 \\ + 0.17 \\ \end{array} $	217 192 45 65 62 10 8 309

The weighted mean of these results gives as the definitive latitude correction applicable to the mean system of the Ledgers

C. F. C., 1900.

 $\Delta \phi = + 0'' \cdot 14 \pm 0'' \cdot 012.$ 

## CAPE FUNDAMENTAL CATALOGUE, 1900.

The mean latitude of the transit-circle, as derived with the Pulkowa refraction constant, is therefore

-33° 56' 2".36.

Instead of utilising the mean value of the latitude correction in order to reduce the whole series of observations to a homogeneous system, it has been thought preferable to apply to each homogeneous group of observations the values of the corrections derived solely from the observations contained within the group. Corrections have accordingly been applied in accordance with the following table, which include the combined effects of latitude correction, flexure correction, and the reductions A, B, necessary to refer the whole to a homogeneous mean system.

Dec.	1906	-11.	19	05.	19	08.	19	10.
Dec.	I. E.	I. W.	II. E.	II. W.	II. E.	II. W.	JI∠E.	II. W.
$\begin{array}{r} + 35 \\ 30 \\ 25 \\ 20 \\ 15 \\ 10 \\ + 5 \\ - 5 \\ 10 \\ 15 \\ 20 \\ 25 \\ 30 \\ 35 \\ 40 \\ 45 \\ 50 \\ 55 \\ 60 \\ 65 \\ 70 \\ 75 \\ 80 \\ 85 \\ 90 \\ 85 \\ 85 \\ 85 \\ 85 \\ 85 \\ 85 \\ 85 \\ 8$	$+ \circ'39 + \circ'34 + \circ'31 + \circ'27 + \circ'23 + \circ'18 + \circ'15 + \circ'10 + \circ'07 + \circ'05 - \circ'00 - \circ'02 - \circ'07 + \circ'05 - \circ'00 - \circ'02 - \circ'07 - \circ'10 - \circ'13 - \circ'15 - \circ'19 - \circ'23 - \circ'26 - \circ'29 - \circ'32 - \circ'34 - \circ'36 - \circ'38 - \circ'39 - \circ'41 + \circ'43 + \circ'45 + \circ'48$	$+ \frac{0.55}{0.55} + \frac{0.54}{0.55} + \frac{0.54}{0.55} + \frac{0.53}{0.51} + \frac{0.42}{0.42} + \frac{0.38}{0.30} + \frac{0.30}{0.27} + \frac{0.30}{0.27} + \frac{0.21}{0.21} + \frac{0.13}{0.11} + \frac{0.13}{0.00} + \frac{0.17}{0.15} + \frac{0.13}{0.11} + \frac{0.09}{0.00} + \frac{0.02}{0.00} - \frac{0.004}{0.00} + \frac{0.004}{0.000} + \frac{0.004}{0.0000} + \frac{0.004}{0.000} + \frac{0.004}{0.0000} + \frac{0.004}{0.000} + \frac$	$\begin{array}{c} + & 0^{\circ} 31 \\ + & 0^{\circ} 30 \\ + & 0^{\circ} 27 \\ + & 0^{\circ} 24 \\ + & 0^{\circ} 22 \\ + & 0^{\circ} 21 \\ + & 0^{\circ} 20 \\ + & 0^{\circ} 20 \\ + & 0^{\circ} 19 \\ + & 0^{\circ} 10 \\ + & 0^{\circ} 14 \\ + & 0^{\circ} 10 \\ + & 0^{\circ} 08 \\ + & 0^{\circ} 05 \\ + & 0^{\circ} 03 \\ - & 0^{\circ} 05 \\$	$+ \circ' \circ	$\begin{array}{c} + 0^{\circ} 17 \\ + 0^{\circ} 16 \\ + 0^{\circ} 14 \\ + 0^{\circ} 12 \\ + 0^{\circ} 11 \\ + 0^{\circ} 10 \\ + 0^{\circ} 04 $	$\begin{array}{c} " & 0.41 \\ + & 0.45 \\ + & 0.46 \\ + & 0.47 \\ + & 0.49 \\ + & 0.51 \\ + & 0.51 \\ + & 0.52 \\ + & 0.51 \\ + & 0.50 \\ + & 0.50 \\ + & 0.50 \\ + & 0.50 \\ + & 0.46 \\ + & 0.43 \\ + & 0.46 \\ + & 0.43 \\ + & 0.36 \\ + & 0.31 \\ + & 0.29 \\ + & 0.27 \\ + & 0.22 \\ + & 0.22 \\ + & 0.22 \\ + & 0.22 \\ + & 0.22 \\ + & 0.22 \\ + & 0.22 \\ + & 0.22 \\ - & 0.22 \\ - & 0.29 \end{array}$	$\begin{array}{c} - & 0^{\circ}18 \\ - & 0^{\circ}19 \\ - & 0^{\circ}21 \\ - & 0^{\circ}22 \\ - & 0^{\circ}22 \\ - & 0^{\circ}22 \\ - & 0^{\circ}22 \\ - & 0^{\circ}20 \\ - & 0^{\circ}19 \\ - & 0^{\circ}20 \\ - & 0^{\circ}19 \\ - & 0^{\circ}22 \\ - & 0^{\circ}$	$\begin{array}{c} + & 0 \cdot 06 \\ + & 0 \cdot 10 \\ + & 0 \cdot 12 \\ + & 0 \cdot 15 \\ + & 0 \cdot 16 \\ + & 0 \cdot 18 \\ + & 0 \cdot 19 \\ + & 0 \cdot 20 \\ + & 0 \cdot 21 \\ + & 0 \cdot 21 \\ + & 0 \cdot 22 \\ + & 0 \cdot 23 \\ + & 0 \cdot 23 \\ + & 0 \cdot 23 \\ + & 0 \cdot 22 \\ + & 0 \cdot 21 \\ + & 0 \cdot 17 \\ + & 0 \cdot 12 \\ + & 0 \cdot 17 \\ + & 0 \cdot 14 \\ + & 0 \cdot 12 \\ + & 0 \cdot 11 \\ + & 0 \cdot 09 \\ + & 0 \cdot 07 \\ + & 0 \cdot 01 \\ - & 0 \cdot 11 \\ - & 0 \cdot 13 \\ \end{array}$

TABLE XV.—Table of Systematic Corrections to the Declination.

xxxiv

#### III.—FORMATION OF DEFINITIVE CATALOGUE PLACES.

The systematic periodic corrections to the right ascensions derived in § I. (p. xxvi) and the corrections to the declinations derived in § II. (p. xxxiv) were applied to the Ledger places, and separate means were first formed for the groups of observations in each of the four conditions I. E., I. W., II. E., II. W. These separate means were then combined into a single mean, with weights dependent on the number of observations in each group, in accordance with the following scheme of weights :---

0	0
No. of Observations.	Combining Weight.
I	13
2-3	$\frac{1}{2}$
4-7	I
8-10	$I\frac{1}{2}$
10+	2

In the case of those stars which are contained in Newcomb's Catalogue, the observations in the Ledgers have been referred to the mcan epoch 1900.0 by the application of Newcomb's proper motions. In forming the final Catalogue positions, the proper motions thus introduced have been removed.

In the case of the double stars Sirius, Procyon, and a Centauri, the reductions to epoch include also the reductions from the bright (or observed) component to the centre of gravity of the system. The corrections thus introduced have been removed in like manner, so that the places quoted in the Catalogue represent the position of the actual object observed referred to the equinox 1900.0, but to the mean epoch of observation.

The right ascensions of the close circumpolars observed during 1911 have been adopted without further modification from the discussion of the observations contained in *Cape Annals*, vol. xi., part iii. The declinations of these same stars have been derived from the combination of the results therein with additional observations in other years. These additional observations have first received corrections, as indicated in the last section, and the combination has then been effected by regarding all the observations as of equal weight, *i.e.* the means from the various groups have been combined with weights simply proportional to the number of observations in each.

The entries in the separate columns of the Catalogue have the following significance :---

Column 1.—" No." The rotation number. \* and † attached to a number indicate a footnote, † being used in the case of double stars.

Column 2.—" Mag." The magnitude taken from Boss's Catalogue or the Harvard Publications, or a few, marked with an asterisk, from recent Cape Observations.

Column 3.—" Name." For Bradley stars the name in Auwers' Bradley has been adopted, except in a few cases mentioned in footnotes; for stars south of declination  $-23^{\circ}$ , the C.G.A. has been followed, with the exceptions used by Auwers in vol. xlvii. of

### CAPE FUNDAMENTAL CATALOGUE, 1900.

the Monthly Notices. The names of the stars z Octantis, A Octantis, have been retained in accordance with the usage in previous Cape Catalogues. For stars otherwise unnamed, a Catalogue number is given in the following order of preference :—Bradley; Mayer; Lacaille; Piazzi; Lalande; Brisbane; Catalogo General Argentina (C.G.A); Cape 1880; Gilliss's Circumpolar Zones; Bonn Durchmusterung. m, pr, seq, br in this column signify mass, preceding component, following component, or bright component.

Columns 4 and 9.—" Mean R.A. 1900.0" and "Mean Dec. 1900.0" respectively. The mean right ascension and declination derived from the observations made for the purposes of this Catalogue, and combined according to the methods described above. They are referred to the *mean epoch of observation*, but to the *equinox of* 1900. The third decimal figure is omitted from the Mean R.A. of Polar stars observed in 1911 only. The R.A. is supplied to the nearest second for stars not observed in this element.

Columns 5 and 10.—" $\mu\Delta E$ ." The quantities tabulated in these columns are the corrections on account of proper motion to be applied to the entries in the columns immediately preceding in order to refer the latter to the epoch as well as the equinox of 1900.0. They depend on the values of the proper motions in columns 8 and 13.

Columns 6 and 11.—"Annual Variation 1900." The annual changes in right ascension and declination due to the combined effects of precession and proper motion. Where no entry is contained in the columns immediately preceding, the quantities in these columns represent the annual precession computed from the formulæ

$$\begin{array}{c} p_a = m + n \tan \delta \sin a , \\ p_\delta = n \cos a \end{array} \right\} \qquad . \qquad . \qquad . \qquad . \qquad (A)$$

where, in accordance with Newcomb's values for the precessional motion,

$$m = 3^{s \cdot 07234}$$
  

$$n = 1^{s \cdot 33646}$$
  

$$= 20'' \cdot 0468.$$

Columns 7 and 12.—"Sec. Var. 1900.0." The quantities given in these columns are in general the centennial variations of the annual variations due to the combined effect of the motions of the pole and equinox and the "proper motion" of the star. If we denote by  $\alpha$ ,  $\delta$  the true co-ordinates of a star referred to the *mean* equator and equinox of epoch t, and suppose that t is expressed in terms of the tropical year as unit, the quantities involved are the values for 1900 of the expressions

$$100\frac{d^2a}{dt^2}$$
,  $100\frac{d^2\delta}{dt^2}$ .

Let us suppose that the "proper motion" of the star consists of a motion with uniform velocity along a great circle. In the annexed diagram, let S denote the star's position at time t, S' its position at time  $t + \Delta t$ , and C the pole of the great circle SS". Further let P, Y represent the mean pole and equinox of the epoch t.

xxxvi

Let  $\rho$  denote the amount of the annual proper motion and  $\chi$  its position angle with reference to the pole of epoch t. Then in the diagram below


$$SS' = \rho \Delta t$$

$$< PSC = \frac{\pi}{2} - \chi$$

$$PS = \frac{\pi}{2} - \delta$$

$$< \Upsilon PS = a,$$

The variations in  $\alpha$ ,  $\delta$  due to the precessional motions of P,  $\Upsilon$  are given by the formulæ, (A) above, where however m and n should not be regarded as strictly constant but as



functions slightly variable with the time t. In accordance with Newcomb's determinations these values at epoch 1900 + t are

 $m = 3^{8} \cdot 07234 + 0^{8} \cdot 0000186t$   $n = 1^{8} \cdot 33646 - 0^{8} \cdot 000057t$  $= 20'' \cdot 0468 - 0'' \cdot 0000855t.$ 

If  $\mu_a$ ,  $\mu_\delta$  denote the "proper motions" of the star, in R.A. and declination respectively, referred to the equator and equinox of epoch t;  $\mu_a$ ,  $\mu_\delta$  are the parts of the complete expressions for  $\frac{da}{dt}$ ,  $\frac{d\delta}{dt}$  which cannot be attributed to precession, *i.e.* 

$$\mu_a = \frac{da}{dt} - p_a \quad , \quad \mu_\delta = \frac{d\delta}{dt} - p_\delta \quad . \quad . \quad . \quad . \quad . \quad (B).$$

The changes thus represented by  $\mu_{\alpha}, \mu_{\delta}$  in the interval  $\Delta t$  result solely in the transference of the star from the point S' to the point S', irrespective of any motion which may be attributed to the points  $P, \Upsilon$ . Hence if we draw the perpendicular S'N on PS, we have

$$S'N = \mu_a \Delta t \cos \delta$$
$$SN = \mu_\delta \Delta t.$$

## CAPE FUNDAMENTAL CATALOGUE, 1900.

But we have also, from the triangle SS'N,

whence

Denote by A, D the right ascension and declination of the point C. The motion of the star being along a great circle, this point will be stationary, and therefore any changes in A, D must be solely those resulting from the precessional motions of the pole and equinox of reference. Hence

$$\frac{dA}{dt} = m + n \tan D \sin A$$

$$\frac{dD}{dt} = n \cos A$$
(D).

We have likewise, from (B),

$$\frac{da}{dt} = m + n \tan \delta \sin \alpha + \mu_{\alpha} \left\{ \begin{array}{cccc} & & \\ & & \\ \frac{d\delta}{dt} = n \cos \alpha + \mu_{\delta} \end{array} \right\} \quad . \qquad . \qquad . \qquad . \qquad . \qquad (E).$$

But in the spherical triangle PCS, we have

$$PS = \frac{\pi}{2} - \delta \quad , \quad PD = \frac{\pi}{2} - D \quad , \quad SC = \frac{\pi}{2} \quad , \quad$$

whence

$$\cos \delta \sin \chi = + \sin D \sin \delta \sin \chi = -\cos D \cos (a - A) \cos \chi = +\cos D \sin (a - A)$$
 (F).

and therefore, by means of (C),

$$\begin{array}{l} \mu_{a}\cos^{2}\delta = \rho \sin D, \\ \mu_{\delta} = \rho \cos D \sin (a - A) \end{array} \right\}$$
 (G).

The conditions that the proper motion is uniform along a great circle are expressed by equations (D), together with the additional equation

$$\frac{d\rho}{dt} = 0.$$

Hence, if we differentiate equations (G) and substitute for  $\frac{dA}{dt}$ ,  $\frac{dD}{dt}$  from (D),

we find

$$\frac{d}{dt}(\mu_{\alpha}\cos^{2}\delta) = \rho \cos D (n \cos A)$$

$$\frac{d\mu_{\delta}}{dt} = -\rho \sin D \sin (\alpha - A) (n \cos A) + \rho \cos D \cos (\alpha - A) \left(\frac{d\alpha}{dt} - m - n \tan D \sin A\right),$$

which, by means of (F), reduce to

$$\frac{d}{dt}(\mu_{\alpha}\cos^{2}\delta) = n\rho\left(\cos\chi\sin\alpha - \sin\delta\sin\chi\cos\alpha\right) \qquad .$$
$$\frac{d\mu_{\delta}}{dt} = -n\rho\cos\delta\sin\chi\sin\alpha - \rho\sin\delta\sin\chi\left(n\tan\delta\sin\alpha + \mu_{\alpha}\right)$$
$$= -n\rho\sec\delta\sin\chi\sin\alpha - \mu_{\alpha}\rho\sin\delta\sin\chi.$$

xxxviii

Replacing  $\rho \sin \chi$ ,  $\rho \cos \chi$  by means of (C), we derive

$$\frac{d}{dt}(\mu_{\alpha}\cos^{2}\delta) = n(\mu_{\delta}\sin\alpha - \mu_{\alpha}\sin\delta\cos\delta\cos\alpha),$$
$$\frac{d\mu_{\delta}}{dt} = -n\mu_{\alpha}\sin\alpha - \mu_{\alpha}^{2}\sin\delta\cos\delta.$$

In virtue of the second of equations (E), the first of these gives

$$\frac{d\mu_{\alpha}}{dt} = n\mu_{\delta}\sin\alpha\sec^{2}\delta + n\mu_{a}\tan\delta\cos\alpha + 2\mu_{a}\mu_{\delta}\tan\delta.$$

Finally, on differentiating equations (E) and substituting for

$$\cdot \quad \frac{da}{dt} \quad , \quad \frac{d\delta}{dt} \quad , \quad \frac{d\mu_a}{dt} \quad , \quad \frac{d\mu_b}{dt}$$

from (E) and from the equations just derived, we find

$$\begin{aligned} \frac{d^2a}{dt^2} &= \frac{dm}{dt} + \frac{dn}{dt} \tan \delta \sin \alpha + n \sec^2 \delta \sin \alpha (n \cos \alpha + \mu_{\delta}) \\ &+ n \tan \delta \cos \alpha (m + n \tan \delta \sin \alpha + \mu_{\alpha}) \\ &+ n\mu_{\delta} \sin \alpha \sec^2 \delta + n\mu_{\alpha} \tan \delta \cos \alpha + 2\mu_{\alpha}\mu_{\delta} \tan \delta, \\ &= \frac{dm}{dt} + n^2 \sin \alpha \cos \alpha + \tan \delta \left(\frac{dn}{dt} \sin \alpha + mn \cos \alpha\right) + \tan^2 \delta (n^2 \sin 2\alpha) \\ &+ 2n\mu_{\alpha} \tan \delta \cos \alpha + 2n\mu_{\delta} \sec^2 \delta \sin \alpha + 2\mu_{\alpha}\mu_{\delta} \tan \delta. \end{aligned}$$
$$\begin{aligned} \frac{d^2\delta}{dt^2} &= \frac{dn}{dt} \cos \alpha - n \sin \alpha (m + n \tan \delta \sin \alpha + \mu_{\alpha}) \\ &- n\mu_{\alpha} \sin \alpha - \mu_{\alpha}^2 \sin \delta \cos \delta \\ &= \frac{dn}{dt} \cos \alpha - mn \sin \alpha - n^2 \sin^2 \alpha \tan \delta - 2n\mu_{\alpha} \sin \alpha - \frac{1}{2}\mu_{\alpha}^2 \sin 2\delta. \end{aligned}$$

Replacing m, n,  $\frac{dm}{dt}$ ,  $\frac{dn}{dt}$  by their values for the epoch 1900, and expressing the results in seconds of time and seconds of arc respectively, we finally obtain the following numerical expressions for the centennial variations of the annual variations which figure in the Catalogue:

$$100\frac{d^{2}a}{dt^{2}} = 0^{8} \cdot 00186 + [7 \cdot 81255] \sin 2a$$

$$+ \{ [8 \cdot 47508] \cos a - [6 \cdot 756] \sin a \} \tan \delta$$

$$+ [8 \cdot 11358] \sin 2a \tan^{2}\delta$$

$$+ [8 \cdot 28865] \mu_{a} \tan \delta \cos a + [7 \cdot 11256] \mu_{\delta} \sec^{2}\delta \sin a$$

$$+ [6 \cdot 9866] \mu_{a}\mu_{\delta} \tan \delta,$$

$$100\frac{d^{2}\delta}{dt^{2}} = -[7 \cdot 929] \cos a - [9 \cdot 65117] \sin a - [9 \cdot 28967] \sin^{2}a \tan \delta$$

$$- [9 \cdot 36474] \mu_{a} \sin a + [8 \cdot 7367] \mu_{a}^{2} \sin 2\delta.$$

Columns 8 and 13.—" Proper Motion." These quantities are the proper motions as above described. The numerical values adopted have been taken from Boss's Catalogue for all stars contained therein; from Newcomb when the Newcomb No. 18 given in the last column; and from the Cape Catalogue of Astrographic Standard Stars when marked \*.

Column 14.—"No. of Obs." This indicates the number of observations. When two numbers are quoted, the former applies to the right ascensions, and the latter to the

xxxix

declinations. When a single number only is given, it is to be regarded as applicable to both elements, or in a few cases to the single element observed.

Column 15.—" Epoch 1900 +." The mean epoch of observation, expressed in years in excess of 1900. When the epochs of observation in the two elements are not identical, two epochs are quoted, the former of which refers to the right ascensions and the latter to the declinations.

Column 16.—"Boss No." The number of the star in Boss's Preliminary General Catalogue. For a few stars contained in Newcomb's Catalogue but not in Boss's, the Newcomb number is inserted, preceded by N.

## IV.—FURTHER CORRECTIONS TO THE CATALOGUE RIGHT ASCENSIONS.

The system of right ascensions depends on that of the equatorial clock stars as revised through the medium of the daylight observations. The extension of this system to the higher declinations depends on the assumption that the form of the pivots has remained sensibly invariable throughout the period of observations for the Catalogue. The pivot corrections employed were based on observations made in the years 1902 and 1904, before the commencement of the Catalogue observations.

A new determination has recently been made (1914 July). It will be sufficient here to exhibit the differences between the two determinations as affecting the mean results obtained in the four conditions I. E., I. W., II. E., II. W., as the star observations have been very approximately symmetrically distributed in relation to these conditions.

Denoting by  $\Delta T$  the amount by which a transit is accelerated in consequence of pivot error, Table XVI., p. xli, gives the values of  $\Delta T \cos \delta$ , in the mean of the four conditions, for each 5° of zenith distance in accordance with both the old and new determinations.

This table shows that the effect of wear of the pivots, so far at least as it can affect the mean system of the Catalogue, is insignificant and justifies the use of the earlier determinations throughout.

The equinox of the Catalogue has not been derived from fundamental considerations, but has been based on Newcomb's determination. It remains to examine to what extent the concurrent observations of the Sun indicate a modification of this equinox, *i.e.* by what amount in common all the Right Ascensions should be increased or decreased. The details of the Sun observations will be given in full in a separate publication. To the observed right ascensions and declinations of the Sun, "day corrections" have been applied, derived from observations of bright stars at about the

same time. These "day corrections" were computed from the final star places contained in the Catalogue. Thus the derived right ascensions and declinations of the Sun are in systematic accordance with those of the Catalogue. These

TABLE XVI.—Corrections	on	account	of	the	Form	of	the	Pivots.
------------------------	----	---------	----	-----	------	----	-----	---------

Zenith Distance (South).	Old Determination.	New Determination.	Old-New.	Zenith Distance (South).	Old Determination.	New Determination.	Old-New.
$ \begin{array}{r} - 90 \\ - 85 \\ - 80 \\ - 75 \\ - 70 \\ - 65 \\ - 60 \\ - 55 \\ - 50 \\ - 45 \\ - 40 \\ - 35 \\ - 30 \\ - 25 \\ - 20 \\ - 15 \\ - 10 \\ - 5 \\ \end{array} $	$ \begin{array}{c} 8 \\ + 0.025 \\ + .025 \\ + .024 \\ + .022 \\ + .020 \\ + .019 \\ + .015 \\ + .010 \\ + .005 \\ + .002 \\003 \\007 \\007 \\007 \\009 \\007 \\005 \\005 \\001 \end{array} $		$ \begin{array}{c}         8 \\         - 0.004 \\        002 \\         + .001 \\         .000 \\         + .001 \\         .000 \\         + .001 \\         .000 \\         + .001 \\         .000 \\        000 \\        000 \\        001 \\        001 \\        002 \\        002 \\        002 \\        002 \\        001 \\         + .001 \\         + .001 \\         + .001 \\        002 \\        002 \\        002 \\        001 \\         + .001 \\         + $	$ \overset{\circ}{} + \frac{15}{10} + \frac{15}{20} + \frac{25}{40} + \frac{35}{40} + \frac{45}{45} + \frac{45}{70} + \frac{45}{775} +$	$ \begin{array}{r}                                     $	$\begin{array}{c} & & & \\ & & & \\ & & & \\$	$\begin{array}{r} & & \\$

$\Delta T$	cosd	
------------	------	--

observations have been analysed by a method exactly similar to that given in Cape Annals, vol. ii., part 5. The resulting correction to the equinox derived in the different years over which the Sun observations extend in the sense in which it is to be applied as a uniform correction to the right ascensions of the catalogue are as follows:— .

			S
1907	•		 -0.069
1908			-0.104
1909			-0.104
1910			-0.082
1911		•	-0.056

The discordance between the results obtained in different years, and more especially the pronounced fall in value for the year 1911, where a different method of observing was used, indicate that but little weight can be attached to the results. Separating out C. F. C., 1900. 6

the residuals as obtained from the observations by different observers, we obtain the following more extended table :---

Observer.	1907.	1907. 1908. 1909. 191					
	s	s	S	S	S		
C	-0.067	-0.014	-0.026	-0.094	-0.064		
AP	-0'105	-0.141	-0.146	-0.124	-0'018		
$\mathbf{RC}$	-0.000	-0.000	-0.003	-0'109	-0.067		
AW	-0.077	-0.014	-0.102	-0.000	-0.034		
Μ	-0.080	-0.080	-0.000	-0.080	-0.001		
JW	-0.038	-0.028	-0.001	-0.082			
JJ	-0.100	-0'120	-0.106	-0.104	-0.021		
S					-0.060		

Separate Determination of Equinox Correction by Different Observers.

If we disregard the final column, the quantities in the same horizontal line for the most part give a satisfactory agreement, showing that the discordances between quantities in the same vertical column depend to a greater extent on systematic personality in observing than on accidental errors of observation. Combining the observations of 1907-10, where the same method of observing was used throughout, we obtain the following determinations, each based on homogeneous series of observations :—

Observer.	1907-10.	1911.
~	s	s
С	-0.028	-0.064
AP	- 136	018
RC	092	067
AW	085 -	034
М	079	- '001
JW	066	
JJ	- '108	021
S		060

Assuming that the accidental errors of these determinations are insignificant in comparison with the systematic errors, and that each determination is equally liable to such systematic error, we may advantageously combine these with equal weight and derive

### $\Delta a = -o^{s} \cdot o68 \pm o^{s} \cdot oo63.$

The probable error here derived from the residuals represents the combined effect of accidental and systematic error.

This correction has not been applied, as it appeared preferable to await the result of a more definitive correction to Newcomb's equinox, which it would seem can scarcely be reliably determined without the combination of observations from several observatories and extending over longer intervals.

## V.—FURTHER CORRECTIONS TO THE CATALOGUE DECLINATIONS.

The declination system of the Catalogue has been based purely on fundamental considerations, except in one respect, viz. that the Pulkowa refraction tables have been adopted. The latitude of the Observatory is not sufficiently high to permit of a fundamental determination of the refraction constant being made by means of declinations observed at both eulminations. Recourse must therefore be had to comparison of the declination system with results derived from northern observatories. The most recent and comprehensive data available for the purpose are those of Boss's Preliminary General Catalogue.

Arranging the results in order of declination, we obtain the following comparison :---

Limits of Declination.	Δδ (Cape Funda- mental—Boss).	No. of Stars.	Limits of Declination.	Δδ (Cape Funda- mental-Boss).	No. of Stars.
above $+30^{\circ}$ + $30^{\circ}$ to $+20^{\circ}$ + $20^{\circ}$ ,, $+10^{\circ}$ + $10^{\circ}$ ,, $0^{\circ}$ $0^{\circ}$ ,, $-10^{\circ}$ - $10^{\circ}$ ,, $-20^{\circ}$ - $20^{\circ}$ ,, $-30^{\circ}$ - $30^{\circ}$ ,, $-40^{\circ}$	$ \begin{array}{r}  + 0^{\circ}49 \\  + 0^{\circ}36 \\  + 0^{\circ}28 \\  + 0^{\circ}10 \\  + 0^{\circ}02 \\  \cdot 00 \\  + 0^{\circ}10 \\  + 0^{\circ}18 \\ \end{array} $	46 110 112 121 101 93 115 93	$ \begin{array}{c} -40^{\circ} \text{ to } -50^{\circ} \\ -50^{\circ} , , -60^{\circ} \\ -60^{\circ} , , -70^{\circ} \\ -70^{\circ} , , -80^{\circ} \\ -80^{\circ} , , -90^{\circ} \\ \text{below pole} \\ -90^{\circ} \text{ to } -80^{\circ} \\ -80^{\circ} , , -70^{\circ} \end{array} $	$ \begin{array}{r} + 0^{''}29 \\ + 0^{'}27 \\ - 0^{'}02 \\ - 0^{'}12 \\ + 0^{'}03 \\ - \Delta\delta \\                                   $	176 104 72 55 35  33 24

Comparison between the Declinations of the Catalogue and Boss's Preliminary General Catalogue.

Equating these differences to the expression

$$-\Delta\phi - \Delta k \tan \zeta$$

where  $\Delta \phi$  denotes a correction to the latitude consequently on an alteration  $\Delta k$  in the refraction constant, and weighting the resulting equations proportionally to the numbers in the final column, we derive the normal equations

$$129^{\circ}0\Delta\phi + 31^{\circ}6\Delta k = -20''^{\circ}33$$
$$31^{\circ}6\Delta\phi + 133^{\circ}6\Delta k = -16''^{\circ}97$$

with the solution

$$\Delta \phi = - \circ'' \cdot 134$$
  
$$\Delta k = - \circ'' \cdot 095.$$

The refractions used in the formation of the Catalogue are taken from the Pulkowa Tabulæ Refractionum. For atmospheric conditions which correspond closely with the mean conditions under which the observations were made the refractions computed from these tables are given in the second column of the following table :---

Comparison of Mean Refractions from Pulkowa and Paris Tables.

\$,	Pulkowa.	Paris.	Diff.	$\Delta k$ tan $\zeta$ .
•	"	"	"	"
o	0.00	0.00	0.00	0.00
IO	10.04 '	10.05	+ .02	·02
20	20.73	20.69	.04	.03
30	32.87	32.81	•06	.05
40	47.76	47.68	.08	.08
45	56.89	56.79	.10	.10
45 50 55 60	67.77	67.64	.13	.11
55	81.14	80.99	.12	.13
60	98.30	98.13	.17	.19
65	121'49	121.28	.51	.20
70	155.11	154.85	•26	•26
	209'2	208.8		*35
75 80	311.8	311'2	·4, •6	.54

Barometer 30 inches. Thermometer 60° F.

The third column gives the refractions for the same atmospheric conditions derived from the tables of the Connaissance des Temps, 1916. It will be seen that these are slightly smaller than those from the Pulkowa tables, but that the differences shown in the fourth column correspond almost exactly with the value  $\Delta k$  tan  $\zeta$  in the fifth column as derived from a comparison of the present Catalogue with Boss. Thus it appears that the refractions used have been too large and that a very close agreement between the results of the Cape observations and those of northern observatories would have been secured had the Paris tables been used instead of the Pulkowa tables. A similar but slightly larger reduction from the Pulkowa values is indicated by a recent discussion of Pulkowa observations (v. Backlund, *Die Deklinationssysteme der Pulkowoer Kataloge 1885, 1892, 1900*, Mitteilungen der Nicolai-Hauptsternwarte zu Pulkowo, Band VI. 1).

It remains to examine the effect of the modified constant on the derived value of the latitude of the transit-circle. From a comparison between observations of upper culminations and lower culminations of circumpolar stars, using the Pulkowa refractions, the value obtained above, (§ II.) p. xxxiv, was

-33° 56' 2".36.

The discussions of this section indicate a correction to this quantity amounting to

-0".134,

yielding as the definitive value of the latitude of the transit-circle from the observations for the present Catalogue

-33° 56' 2".49.

We may compare with this the values derived from previous series of observations. These have all been obtained with different instruments, but the difference of geodetic latitude has been accurately derived from measurements at the surface, showing that the position of the new transit-circle is in latitude 1".05 to the North of the old.

The latitude of the old transit-circle derived from observations between 1879 and 1885 is discussed in the Introduction to the Cape Catalogue, 1885 (p. xlvii.), and the definitive value arising from this discussion is

Again, from zenith telescope observations by the Talcott method between the years 1886 and 1891, the latitude of instrument, mounted in the same geodetic latitude as the old transit-circle, was found to be

(Introduction to Cape Catalogue, 1885, p. xlvii.)

The result derived for the old transit-circle for the period 1885-95 (Introduction to Cape Catalogue, 1890, p. xxiv.) is

The mean of these three determinations, regarded as of equal weight, amounts to

or, on applying the correction for the difference of latitude of the two instruments, we obtain for the latitude of the new transit-circle

in almost exact accord with the value derived from the discussion of the observations for the present Catalogue.

7

C. F. C., 1900.

# NOTE.

The Right Ascensions of the Catalogue depend on Newcomb's equinox, but have in other respects been fundamentally derived.

To refer the observations to an absolute system based on concurrent Cape observations of the Sun, a correction of

should be applied throughout.

The Declinations are based on the Pulkowa refractions (*Tabula Refractionum*), and the value

for the mean latitude of the transit-circle, derived from the observations themselves.

A re-determination of the refraction constant and latitude from comparison of the results with Boss's Preliminary General Catalogue indicates the following correction to the declinations

Dec.	Δδ.	Dec.	Δδ.
+40 + 30 + 20 + 10 0 - 10 - 20	$ \begin{array}{r} -0.47 \\ -33 \\ -27 \\ -23 \\ -20 \\ -18 \\ -16 \end{array} $	$ \begin{array}{r} -30^{\circ} \\ -40^{\circ} \\ -50^{\circ} \\ -60^{\circ} \\ -70^{\circ} \\ -80^{\circ} \\ -90^{\circ} \end{array} $	$ \begin{array}{r} & -0.14 \\ - 0.13 \\ - 0.11 \\ - 0.09 \\ - 0.07 \\ - 0.04 \\ 00 \end{array} $

corresponding with the resulting value

for the latitude of the transit-circle.

Uriv. of California

A

# CATALOGUE OF 1293 STARS

# REDUCED WITHOUT PROPER MOTION

TO THE

# EQUINOX 1900<sup>.</sup>0.

# 

# CAPE FUNDAMENTAL CATALOGUE OF STARS FOR 1900'0,

No.	Mag.	Name.	Mean R.A. 1900°0.	$\mu_a \Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900 O.	Sec. Var. 1900'0,	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
		Dissium	h m s 0 0 12.978	S	8	8	ś	ê	- ".87	+20"1 37		+ ".090		0.61	
1 2	4·8 6·6	33 Piscium 5 Ceti		+ .013	+ 3.0709	- ·0014 + ·0005	- · co13 + · 0003	-61559.94 - 3015.60	+ .02	20.040	- '009 '015	- '005	19 16:17	9.64 9.32	9
3	2.0	21 Andromedæα	3 13.099	087	3.0931			+28 32 16.89	+ 1.32	19.884	.012	161	22	8.18	10
4	7.4	Lacaille 9745			2.751	- '397		-86 35 44.54		20.045	.014		45:68	11.22 : 10.11	
5	3.9	Phœnicis	4 20.374	092	3.0221	0288	+.0115	-46 17 58.07	+ 1.52	19.857	.017	186	22:23	8.22:8.15	16
	_			1010	1	10105	1.00010				Loor	1 1017			
6	5.7	Sculptoris	0 6 29·826 6 39·175	- '104	+ 3.0532 3.0574	- ·0137 - ·0190	+.0013	-28 21 23.93 -35 41 33.28	- ·13 - ·96	+20.056	- °021 °022	+ '017 + '120	23 22	7'93 8'03	23 24
7 8	5°3 2°9	88 Pegasiγ	8 5.132	.000	3.0846	+ '0102	1	+14 37 39.38	+ '11	20.021	'024	- '013	22:21	8.52 : 8.27	27
9	5.9	Lacaille 23		+ .08	2'351	- '200	007	-85 33 2.25	68	20.089	.023	+ .059	31:30	11.24	32
10	6.1	35 Piscium		061	+ 3.0873	+ .0068	+.0066	+ 8 15 56.22	+ '22	20.004	.028	- '024	17	9.31	35
		Octantiso		- • 21	- 0.768	+ 2.376	+.019	-88 55 8.34		1	- '002	+ .002	46 : 130	11.20 : 8.01	
11 12	7.5	8 Ceti		+ '011	+ 3.0573	- '0022	0015	- 9 22 42.05	- ·04 + ·29	+20.022	.036	032	22	9.16:8.92	47
13	3.7	Toucani	14 54 339	-2.546	3.1249	0661	+ 2723	-65 27 33.75	-10'90	21.121	Ú Ú	+1.166	17	9.35	53 55
14	5.7	41 Pisciumd	15 27.085	+ .004	3.0840	1		+ 7 38 5.59	- '13	20.012	.039	+ .014	16	9.21	56
15	7.1	Lacaille 75		438	2.9841		+.0572	-51 35 29.58	-	19.712	·046	260	24	7.65	72
16	6.0	44 Piscium	0 20 16.563	+ .015	+ 3.0739	+ .0037	0013	+ 1 23 9.34	+ .12	+19.952	048	016	19:18	9.28 : 9.22	72
17	2.8	Hydri	20 38.34	-8.12	3.2208			-77 48 59.46		20.200	.048		20:58	11.64 : 9.86	73 74
18	2.3	Phœnicisa		-	2.9746			-42 50 59.56		19.229	·049	- '401	22	8.42	78
19	6.7	10 Ceti		046	3.0758			- 0 36 12.31	- '02	19.961		+ .002	17	9.48	79
20	6.3	12 Ceti	24 56.146	004	3.0614	+ .0003	+.0002	- 4 39 35.53	+ .02	19'921	.057	007	32:30	8.23:7.61	90
21	5.4	Piazzi 0'91	0 25 22.691	+ .022	+ 3'0029	0095	0026	-24 20 26.94	13	+19.940	057	+ .016	22:23	8.33 : 8.23	91
22	6.6	Lacaille 109		+ .000	2.0400	- '0206		-41 29 33.94	13	19.936	.056	+ .014	18:19	9'29	92
23	5.0	Phoenicis $\lambda^1$	26 35.785	112	2.9051		+.0130	-49 21 23 12	- '12	19.926	.058	+ .014	20:21	8.82 : 8.76	99
24	4.6	Toncani	26 57.894	123	2.7707	0443	+.0131	-63 30 32.99	+ .21	19.854	•056	054	16	9.41	100
25	5.7	Lacaille 125	28 44 278	+ .018	2:9731	- '0126	0023	-30 6 33.20	+ .55	19.861	.063	028	24:25	7.78 : 7.73	109
26	7.2	Lacaille 133	0 29 28.538	067	+ 2.9206	- '0213	+.0080	-42 58 59.50	19	+19.904	063	+ .023	21:22	8.35 : 8.29	113
27	5.7	Lacaille 137	29 42.638	224	2.8675	-	+.0238	-52 55 31.77	- '23	19.904	.063	+ .025	17:18	9'42:9'25	114
287	5.4	13 Cetim	30 6.326	258	3.0869	+ .0014	+.0272	- 4 8 36.42	+ '17	19.856	•068	- ·018	20	9.48	116
29	4.4	29 Andromedæ	31 32.275	014	3.1939	+ .0244	+.0012	+33 10 7.89	1	19.848	.072	009	21	8.03	123
30†	5.9	Lacaille 147m	32 13.393	854	3.0860	- '0104	+.1055	-25 19 2.88	+ '07	19.840	•073	009	25:28	8.36 : 8.07	127
31	4.5	30 Andromedæe	0 33 15.982	+ .154	+ 3.1612	+ .0208	0173	+28 46 - 5.60	+ 2.31	+19.288	075	248	18	8.93	130
32	3.4	31 Andromedæð	33 58.755	082	3.1985	+ .0224	+.0102	+30 18 49.12	+ .66	19.741	.077	086	21:22	7.67 : 7.62	132
33	8.0	Lacaille 228			0.184			-85 48 4.67		19.793	.013		35:53	11.22 : 10.12	
34	4.7	Phœnicisµ	36 36.087		2.8446			-46 38 1.81	+ .18	19.766	.074	026		7.45:7.36	142
35	2'0	16 Ceti	38 34.367	- 128	3.0133	- '0054	+.0100	-18 32 7.38	- '32	19.803	.082	+ .039	23:22	8.02:8.10	147
36	4.6	Phœnicisη			+ 2.7104	0318	0009	-58 0 40.69	06	+19.766	075	+ .007	20:21	8.65 : 8.50	148
37	6.1	Sculptoris $\lambda^2$	39 22.278	124	+ 2.9070		+.0192	-38 58 20.29	- 1.03	19.868		+ .119	· 18	8.91	153
38	6.8	Lacaille 248			- 0.446	+ .284		-86 14 57.36		19.746	+ .003		49:77	11.00 : 10.51	
39	5.4	Lacaille 193		+ .031	+ 2.9730		0034	-22 33 20.41	- '79	19.832	083	+ .087	18	9'07	155
40	6.0	Lacaille 207	41 4'343	145	2.8246	- '0232	+.0128	-48 6 .3.35	66	19.807	082	+ .081	20:21	8.12:8.09	158
41	4.2	34 Andromedæ		-	+ 3.1721	+ .0180		+23 43 22.88	+ .72	+19.631	093	080	19	8.96	164
42	6.0	Mayer 24		480	3.1432			+ 4 45 48.52	+10.92	18.549	• 096	-1.144	16	9.59	171
43	4.6	63 Pisciumδ			3.1000	1 -		+ 7 2 26.54		19.643	·094	044	18:17	9.67 : 9.64	173
44 45	5.1	Hydriλ 20 Ceti	45 7.493 47 53.827	315	2.0993 3.0638	$- \cdot 0307$ + $\cdot 0037$	+.0355	-75 28 4·36 - I 4I 14·25		19.644	.069	- ·016	19 21 : 22	8.88	182
1.2							1.00	1.0		19.292				9'55 : 9'52	191
46	5.2	Toucani $\lambda^2$		+ .012	+ 2.2524	0325	0023	-70 4 4.64		+19.210	081	037	26	7.45	204
47 48	5.7	68 Pisciumh Sculptorisα	52 25.292	008	3.2374			+28 27 6.34		19.513	.112	- '012	17	9.38	209
40	4'4 4'5	71 Piscium	53 47°317 57 45°109	- ·004 + ·042	2·8942 3·1099			+7295352.18 +7216.40	- '0I - '22	19'498 19'442		+ '001	27:28	7'35:7'32 7'83:7'71	212 226
50	6.3	Lacaille 288			2.5487			-57 32 26.84		19 442		+ .012		8.48 : 8.41	220
	1		5, 457		- 54-7		1			1		1			
10000															

28. 5'9, 6'6; very close binary. 30. 6'6, 6'7; very close binary. .

2

,

FROM OBSERVATIONS IN THE YEARS 1905-11.

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_a \Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	<sup>*</sup> Epoch 1900+.	Boss No.
			hm s	8	s	8	8	0 1 11		+19.360					
51	6.2	26 Ceti 72 Piscium	0 58 40°268 0 59 48°552	- ·077 + ·002	+ 3.0851			+ 0 49 50.65 + 14 24 30.63	+ ":33		- •122	034	22	9.73	230
52 53†	5.9	Phœnicism. B		+ .002 + .032	2.6840	+ .0129	)	-47 15 14.75	+ '10	19.412 19.313	·127 ·112	+ .044	16 : 17 27 : 30	9°37 7°58 : 7°41	234
54	5.7	80 Pisciume	0. 0	+ .171	3.0868	+ .0078		+ 5 7 12.84	+ 1.71	19 313	130	181	16	9.42	245 252
55	5.5	Toucani	3 21.143	101	2.3884	0248	1	-62 18 34.05	- '13	19.301		+ .012	18:20	9.10:8.08	254
1											52.				1
56	3.2	31 Cetiη 43 Andromedæβ	1 3 33 705		+ 3'0172	·0000 + ·0289		-10 42 45.66	+ 1.20	+19.148	- '129	133	30:35	8.96:9.00	255
57 58	5.3	82 Pisciumg	4 7.975 5 35.730	- '144 + '016	001	+ .0289	1	+35 5 25.01 +30 53 34.13	+ 1.11 + .13	19.122	• 143	- ·115 - ·014	16 16:17	9.68	259
59	4.8	84 Piscium	5 35 730 6 4.640	013	3 2904			+20 30 10.86	- · · · 06	19 217	·144 ·142	+ .006	10:17	9°45 9°39	269
60	4.6	83 Pisciumτ	6 9.080	043	3.2936	+ .0238			+ .29	19.180	142	038	25:28	7.78:7.57	270 271
	-				1.1		1.000			1000					
61 62	6.2	Lacaille 328 86 Piscium pr. 5		055	+ 2.7692	-		-38 23 11.28	+ .26	+19.134	- '127	033	25:26	7.81:7.73	280
63	5.5	37 Ceti	8 30°450 9 21°880	- ·089 - ·079		+ .0001		+7247.11 - 82734.35	+ 51 - 2.21	19.106	•143	052 + .268	21:23	9'96 : 9'82	282
64	5.4	89 Piscium		+ .032					+ '22	19°404 19°025	°140 °148	023	17 17:18	9°37 9°40	285
65	4.8	90 Piscium		- '013					+ '09	19.000	.160	012	25:27	7.83:7.80	295 300
	12.30			Ū		14 EXC11	10 C C C C C C C C C C C C C C C C C C C								300
66	5°5 6°0	91 Pisciuml	1 15 35.482	012	+ 3.3064			+28 12 56.33	+ .71	+18.890	164	076	16	9.36	303
67 68	100 C 100 C 100	Lacaille 384θ	18 51.784	- '002 + '041	2°7966 2°9978	0080	+.0002		+ .45	18.817	•146	022	21	8.13	311
69	3°7 5°8	Lacaille 409	19 1.457 21 37.831		2 9978	+ .0018		84159'33 -645322'10	+ 1.20	18.654 18.789	·155 ·114	- '213	23:24	7.67:7.47 7.66:7.60	313
70	3.3	Phoenicis		+ .025	2.6096	0125	- '0028	-43 49 51.53	+ 1.98	18.496	114	 — `219	18:19	9'04	329
															3~9
71	5.2	48 Ceti			+ 2.8796	1	+.0040	-22 8 47.64	- '02	+18.692		+ .005	17	9.31	331
72	5°2 3°8	98 Piseiumμ 99 Pisciumpr. η	24 56·869 26 7·882	188	0 00.			+ 5 37 41.54	+ '43	18.642	.174	- '044	18 16:18	9.67 : 9.69	332
73†	30	Phœnicisδ	20 7 002	019	3.2042	+ .0142	+.0020	+14 49 49.38	+ '09 - 1·28	18.638 18.766	·179 - ·144	- °010 + °149	20	9'31 : 9'32 8'60	335 336
75	8.1*	Gilliss P.Z. 939	30 25.79		-10.684	+ 5.883	T 0129	-87 51 40.28	- 1 20	18.206		T 149	16	11.20	
1		at a set of the set of the set			1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1										
76	5.8	102 Piscium		+ .048	+ 3.1745	+ '0125		+11 37 48.96	35	+18.497		+ .032	18	9.38	356
77 78	6·3	Lacaille 505a Eridania	0 0 00	+ .000	0.3428	+ .1158		-79 0 45'12	+ .89	18.301	.027	118	47 19	7°52 9°08	358
79	4'7	106 Piscium		+ .011	2.2391	- ·0130 + ·0091		-57 44 41.24 + 4 58 53.99	+ '26	18.355 18.307	·139 - ·193	- ·029	33:34	7'74:7'41	363 378
So	8.7*	· Gilliss P.Z. 1219	37 25.69		-27.909	+29.324		-88 58 50.65		18.263			55 · 54 II	11.20	5/0
		The second se			in the second						-				
81	5.9	Lacaille 501			+ 2.6473		0032		+ .12	+18.236	- 167	- 019	23:21	7.91 : 8.07	386
82	3'5	52 Cetiτ 110 Pisciumο	39 24.419					-16 27 44.07		19.046		+ .856	23	7'73	391
84+	4'4 5'5	Sculptorispr. e						+ 8 39 16.83 -25 33 8.74		18·215 18·076	•203 •183	+ ·051 - ·057	23:24 16	9.39	393 396
85	6.5	Hydri $\tau^1$		100				-79 39 7.37	+ '54	18.142		+ .021	35	9'26:9'30	399
						1000			314						
86	5.6	Lacaille 520		-	+ 2.3542			-51 18 58.49	+ .18	+18.068	126		18:17	9.44	400
87 88	5°3 5°8	Eridaniq <sup>2</sup> Lacaille 634				- ·0109 + 1·172		-54 1 25.80	- '57	18·144 18·080	- ·153 + ·241		17 : 18 56 : 150	9°32 11'64 : 8'87	401 406
89	4.8	53 Ceti		- '22				-85 16 29.04 -11 10 52.38	- ·26 + ·74		197	and the second	17:16	9'41 : 9'40	411
90	6.1	54 Ceti		+ .042	3.1283		and the second	+10 32 53.33		17 913		027	16	9'52	414
22.5	-		101000						10.5316			100			122.00
91	3.8	55 Ceti			+ 2.9602			-10 49 44 91	+ '29	+17.888		032	29:25	9'37 : 9'15 8'44 : 8'25	416
92	3°4 4°8	2 Triangulia		011	3.4094			+29 5 28.22		17.654		- ·232 + ·025	20:22 22:19	8'44 : 8'25 9'25 : 9'14	421 426
93 94	2.7	6 Arietisβ		- ·014 - ·067				+ 2 41 38.72 + 20 19 8.35	$- \cdot 23 + 1 \cdot 11$	17·872 17·706		111	17:19	9 25 . 9 14 9 84 : 9 96	420
95	4.3	Phœnicis	49 0 921		2.4084	0088			+ .79	17.704		092	19:20	8.70:8.58	429
1.5	2					NO.	Sec. 10		1						
96	5.3	Phœnicisφ Eridaniχ		+ .033	+ 2.4923	0080			+ .58	+17.740		- °032	19:20	8.81 : 8.73 8.07 : 7.87	433
97† 98	3°7 4°9	9 Arietis $\chi$		586 + .064	2.3383			-52 6 21.15 +23 6 30.27	- 2.24	17·983 17·668		+ ·285 - ·018	23:25 16	9.39	438 441
90	49	Hydri $\eta^2$		- '101	3°3340 1°5142			+23 0 30.27 -68 8 20.27	$+ \cdot 17$ - $\cdot 84$	17 008		+ .093	18:21	9.18:9.03	442
100	4.1	59 Ceti		083	2.8270			-21 33 44.46		17.543		020	21:23	8.92:8.62	453
							,5			, , , , , , ,			-		

53. 4'1, 4'1; close binary. 73. 3'8, 11 1"'0 15° 84. 5'5, 9'5 4"'7 54° 97. 3'7, 12 6":2 198° 1901 °0. 1902 °9. 1900 °0.

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_a \Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
		Lossille ree	hms	9	S	s	S	· · · · · · · · · · · · · · · · · · ·	+ "85	+17":449	-"181			Queo . Queo	
101 102	5°6 2°9	Lacaille 599	I 55 31.514 I 55 37.480	+ .046	+ 2.4751	- '0072 - '0034	0055 +.0355	-42 30 47 <sup>29</sup> -62 3 22 <sup>3</sup>	- '40	17.594		- °104 + °044	20:21 19:21	8'30 : 8'15 9'11 : 9'09	456 458
103	4.8	Fornacis	2 0 0.211	007	+ 2.6909	- '0035	+.0009	-29 46 35.84	- '02	17.363	- '204	+ '002	25:26	7.89:7.79	474
104	8.1	Lacaille 760			- 5.524	+ 1.524		-85 31 16.35		17.317	+ .399		30:46	11.62 : 10.11	
105	2.0	13 Arietisa	1 32.167	113	+ 3.3728	+ .0204	+.0132	+22 59 21.50	+ 1.10	17.148	257	146	25:24	8.22:7.94	477
106	7.2	Lacaille 764	2 3 13.30		- 5.138	+ 1.347		-85 14 3.97		+17.218	+ .376		37:59	11'61 : 10'30	
107	3.0	4 Trianguliβ	3 35.525	101		+ .0302	+.0123	+34 30 51.98	+ .38	17.156	- '274	046	21:22	8.21:8.28	 482
108	7.0	Lacaille 641		+ .002	2.4445	0060	0002	-42 21 17.58	+ .16	17.163	.101	019	20:23	8.85:8.48	484
109	6.1	15 Arietis		059	3.3178	+ .0177	+.0062	+19 1 42.95	+ .27	17.107	•259	028	16	9.47	491
110	7.2	Lacaille 657		025	2.3526	0001	+.003*	-45 56 19.01	+ '41	17.109	• 186	05*	20	8.24	
III	4.6	65 Cetiξ <sup>1</sup>		+ .016		+ :0116	0012	+ 8 22 39.95	+ .07	+17.008		- :007	02.005	0.65.0.65	FOF
111	5.4	Fornacis	2 7 41 908 8 30·289	+ .016	+ 3.1747 2.6438	0032		-31 11 33.99	- '01	16.979	- ·252 ·212	- ·007 + ·002	23:25 25:26	9.65 : 9.67 7.54 : 7.49	505 506
113	6.2	Lacaille 682		+ .026	2.4289	0050	1 ·	-41 37 57.01	+ .27	16.851	.198	033	20	8.25	512
114	4'1	9 Trianguli		- '029	3.5234	+ .0292	-		+ .44	16.792	•288	021	20:21	8.64 : 8.55	517
115	5.9	67 Ceti		058	2.9906	+ .0049	1	- 6 52 59.39	+ '99	16.705	•245	108	19	9'43:9'15	518
													0.000		
116	5.7	22 Arietisθ		+ .000	+ 3.3292	+ .0180		+19 26 19.33	+ .05	+16.780	272	006	18	9.01	521
117 118	3.8	Eridaui	12 56.306	073	2.1438	- '0045		-51 58 30.01	+ '26	16·739 16·466	179	- '029	19	8.99:8.85	524
110	var. 5.5	Fornacis		+ .001	3.0279	+ '0062	- · 0001 + · 0147	-32555.99 -241614.82	+ 2.27	16.462	·251 ·235	- ·237 - ·061	20:18	9*68 : 9*57 7*66	530
120	5.6	24 Arietisξ	19 27 344	008	3.2100	+ '0127			+ .14	16.434	235	- '015	24 17:18	9.40 : 9.29	543 546
120	30					T 012/	7 0000		1 14		2/5		17.10		10.5
121	4.3	Hydriδ			+ 1.0236	+ .0290	- '0102	-69 6 52.08	11	+16.435	094	+ .015	20	8.75	548
122	5.0	72 Ceti		+ .010	2.8961	.0035	0012	-12 44 29.09	+ .00	16.356	•251	009	16	9.63	551
123	5.6	Horologii		+ .092	1.6731	*0043		-60 45 35.38	+ 1.00	16.180	.148	132	21:24	8.06 : 7.87	557
124	6.2	Hydri	22 16.041		0.3202	•0766	0230	-74 5 55.35	10. +	16.306	•032	- •001	19	9'32 : 9'17	558
125	4.4	73 Cetiξ <sup>2</sup>	22 50.483	- '024	3.1847	.0110	+.9050	+ 8 0 42.93	+ .04	16.274	.278	004	24:22	9.11 : 0.19	560
126	4.5	Eridani	2 23 19.222	011	+ 2.2000	0033	+.0012	-48 9 9.32	+ .12	+16.240	195	013	18	9.23	563
127	6.6	27 Arietis	25 21.488	024	3.3209	+ .0165		+17 15 41.36	+ .01	16.021	•294	- '097	16	9.41	568
128	4'9	76 Cetiσ		+ .025	2.8419	+ .0023	0022		+ 1.10	15.927	•254	112	16:18	9'42	575
129	6.1	Fornacis $\dots \lambda^1$	28 56.836	+ .020	2.2016	- '0022		-35 5 23.18	+ .12	15.940	•227	020	22:24	7'76:7'63	579
130	6.2	Lacaille 799	30 30.210	+ .014	2.0438	0013	0019	-51 31 53.21	+ .18	15.856	•188	- '021	19:21	8.65 : 8.43	587
131	6.2	Piazzi II. 123	2 30 36.938	-1.172	+ 3.2848	+ .0123	+.1208	+ 6 24 49.01	-14.25	+17.335	310	+1.463	16	9.74	588
132	5.1	78 Oeti'		+ .021	3.1439	.0103		+ 5 9 24.77	+ .29	15.842	•286	029	20:16	10'07 : 10'02	589
133	5.6	32 Arietis		+ .000	+ 3.3984	.0192	0006	+21 31 44.66	+ .24	15.213	313	023	17:16	10'35 : 10'28	597
134	7.9	Lacaille 1884	33 13.90		-37.440	29.094		-88 49 42.60		15.731	+3.377		37	11.62	
135	5.2	Hydriµ	33 47 280	- '402	- 1.3882	+ . 2507	+ '0430	-79 32 44.63	+ .28	15.671	+ .112	030	33:35	9.36 : 9.44	601
136	5.4	Horologii	2 34 6.533	077	+ 1.9765	0002	+.0079	-52 58 33.20	+ .22	+15.661	187	022	17:16	9'71 : 9'82	603
137	4.1	82 Cetiδ	34 21.358			+ .0082		- 0 6 10.08	- ·01	15.670		+ .001	16	10.20	604
138	7.8	Lacaille 1029		+ .24	- 9.450	+ 2.492	021	-86 9 42.10	04	15.612	+ .860		41 : 116	11.65 : 8.71	N166
139	4.1	Eridaniı	36 43.405	103	+ 2.3673	- '0021	+.0100	-40 17 0.07	+ .27	15.211	226	029	16:19	9.67 : 9.22	614
140	5.9	34 Arietis μ	36 43 578	023	3.3750	+ .0179	+.0022	+19 35 7.11	+ '49	15.492	.318	047	15	10.42	615
141	4.7	35 Arietis	2 37 34.889	003	+ 3.2101	+ .0233	1.0002	+27 16 54.08	+ .11	+15.479	331	013	16:17	8.76 : 8.65	620
142	4'3	Hydri	38 3.082	- 172	0.0086	·0334		-68 41 43.67	17	15.482	·092	+ .019	16	10'30 : 10'42	621
143†	3.2	86 Cetiseq. 7		+ .093	3.1044	.0092		+ 2 48 50.49	+ 1.43	15.312	•294	120	17	9.2	622
144	4.4	89 Cetiπ		+ .005	2.8539	.0033			+ .14	15.379	•273	014	16	10.10	627
145	4.3	87 Ceti		176	3.2377				+ .25	15.356	.311	027	19:20	9.25 : 9.15	629
146	4.8	39 Arietis	2 41 57 219	- •108	1. 1. 1. 1. 1.				+ 1.18	+15.121	244	- '125	16		624
140	3.5	41 Arietis	44 5.751	- '046	+ 3.2015	+ .0244		+28 49 53.65 +26 50 53.29	+ '98	15.011	- '344	- 125		9°43 9'11 : 8'67	634 643
148	4.5	Fornacis	44 5 7 51	040	2'5121	0004		-32 49 31.59	- 1.23	15.237	•248	+ .160	17:18	9'76:9'58	645
149	5.7	43 Arietisσ	45 58.211	- '021	-	+ .0149		+14 40 12.23	+ .33	14.983	.326	033	18:16	9'94:9'93	648
150	4.9	2 Eridaniτ <sup>2</sup>	46 30.107			+ .0017	0037	-21 24 58.27		14.973	269	012		8.33:8.07	650

118. Mira. L, 1'7-9'6; P, 331<sup>d</sup>'6. 143. 3'5, 7'4 3''1 291° 1903'1.

.

× .

## FROM OBSERVATIONS IN THE YEARS 1905-11.

-						7	1				1			1		
P	To.	Mag.	Name	Mean R.A. 1900'0.	$\mu_{a}\Delta E$	Annual Variation 1900°0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900 '0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
-									1			1				
				h m s	S	8	S	5								
1	51	7.3	Lacaille 1076	2 47 54.87		- 8.138	+ 1.766		-85 26 27.21		+14.903	+ ".787		26:47	11.61 : 10.08	
	2	4.0	3 Eridaui	0 0 000	020	+ 2.9289	+ .0020		- 9 17 47.50		14.474	- •297	212	30:24	9.24 : 8.49	665
	3	4.8	91 Cetiλ Eridanipr. θ	54 21°247 54 28°178	006	3.2108	+ '0118		+ 8 30 31.96		14.214	*329	- ·006 + ·034	17:16	9.27	679
I	54	3°5 5°2	Horologii	56 54 354		2 2/41 I·I2II			-64 28 7.95		14°547 14°393	·234 ·120	+ .034	16 20:19	9°55 8°73 : 8°79	680 690
10					1				1.000 2.00							
I		2.7	92 Cetia 11 Eridani			+ 3.1317	+ '0097	1 -	+ 3 41 50.32	1		- '325	022	20:19	9.08 : 9.04	691
19		4°2 8°1	Lacaille 1203		+ .100	+ 2.6446	·0016 2·743	0104	-24 0 59.69 -86 16 6.71		I4.253 I4.242	275	042	16 19:32	9.63 11.60 : 9.88	696
1		5.9	Lacaille 974		-	+ 2.0508	- 743 ·0011		-47 22 0.26	+ .13	14 242	217	012	19:32	8.94:8.83	 701
16		5.3	Horologii		+ .001	1.4083	.0110	0099	-60 7 32.33		14'031	151	066	17	9.10	706
16		5.7	Hydriθ	1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2 1 2	080	1.	+ .0715	+.0089			+14.084		1 1006			
	12	8.4	Lacaille 1848		+ .72	+ 0.0950	+ .0715	0039	-72 17 34.63 -88 34 20.98	$- \cdot 32$ + $\cdot 25$	13.907	- '017 +3'726	+ .036	17:18	9'03:8'88 11'65:9'43	711 N198
16		4.6	57 Arietisδ	5 54.635		+ 3.4233	·0171		+19 20 54.82		13'799	369	006	23:25	8'51:8'06	718
16	641	5.3	94 Ceti seq.	7 40.295	- 127	3.0201	.0077	1.	- 1 34 13.20		13.638	• 333	055	17:18	9'39:9'38	722
16	5†	6.3	Lacaille 1016. AB	8 55.079	074	2.1068	.0013	+.0085	-44 47 40.22	+ .04	13.608	.232	002	19:21	8.71:8.47	728
16	6	4.9	58 Arietis	3 9 9.077	+ '017	+ 3.4412	+ .0176	0012	+20 40 25.46	+ .75	+13.523	374	075	16	9.99	730
16	7	6.0	Lacaille 1040	10 1.101	013	1.2123			-57 41 44.85		13.246	•169	+ .004	21:23	8.69:8.50	733
16	18	7.2	Lacaille 1020		030	2.3592	•0009	+.0030	-35 55 46.03	21	13.218	• 260	+ .022	17:18	9.84 : 9.53	737
16	-	5.0	13 Eridani	10 58.215	+ .003	2'9120	.0022		- 9 11 27.58	- '42	13.225	• 320	+ .042	16	9*27	739
17	0	2.1	96 Ceti	14 7.081	- •156	3.1420	.0092	+.0122	+ 3 0 13.26	83	13.320	·351	+ .094	16 : 17	8.80 : 8.87	752
17	1	6·1	Lacaille 1058	3 14 11.009	+ .013	+ 1.9538	+ .0028	0015	-48 7 4.96	13	+13.288	219	+ .017	19:23	8.35 : 7.89	754
17	2	5.4	61 Arietisτ	15 27.134	020	3.4565	+ .0174	+.0051	+20 47 11.65	+ .27	13.128	*385	030	21:16	9.20:9.08	761
17		4'4	Eridanie		-2.338	+ 2.3994			-43 27 1.67	- 6.02	13.904		+ .748	19:22	8.58 : 8.15	764
17		5.7	Hydri			- 1.5826	+ .1927	+.0347	-77 45 11.77	22	13.056		+ .000	36:42	8.45 : 8.39	776
17	5	3.6	1 Tauri	19 25 794	+ '042	+ 3.2238	+ .0114	0042	+ 8 40 36.61	+ .20	12.846	- • 364	028	20:19	9.33 : 8.98	778
17	6	3.7	2 Tauri ξ	3 21 44 940	038	+ 3.2466	+ .0110	+.0040	+ 9 23 2.89	+ .38	+12.727	371	- •041	17:18	9'40:9'25	784
17		6.7	Lacaille 1107	22 37 288	- •019	2.1442	.0020		-41 59 14.19	26	12.738	•247	+ .029	17:19	8.99 : 8.84	793
17		5.3	4 Tauris		+ .002	3.2742	°0I22			+ .16	12.233	*377	019	16:17	8.56 : 8.66	801
17		4'4 6'3	5 Tanri	25 21.053 27 24.527	- ·009 - ·068	3°3067 1°9248			+12 35 38.93	+ .03	12.520	·382	- '004 + '016	21:25	7'18:7'08	804
								T 0002	-47 42 59.68	13	12.399	•227	+ .019	19:21	8.33 : 8.23	811
18		4.9	Reticuli	3 27 38.263		+ 1.0362		+.0542	-63 17 20.03	- 3.30	+12.737	- 130	+ .369	19	9'12 : 8'94	812
18		3.7	18 Eridaniε 19 Eridaniτ <sup>5</sup>	28 12.542		2.8247	•0056		- 9 47 47.89	11	12.340		+ .013		8'75 : 8'25	814
18		4 3 5 <sup>.</sup> 8	Lacaille 1144	29 22·248 29 35·857	020	2.6489 1.7845			-21 58 5.68 -50 43 3.50		12.227	·311 ·212	- °020 + °086		9'17 : 9'33 8'89 : 8'80	816 818
18		4'4	10 Tauri		+ .141	3.0283	•0076				12°318 11°599	359	- ·482	19:21 16	9.06	825
18		4.6	and the second sec	Constant of the Sec						1.	and sold the					
18		5.8	Eridaniy Brisbane 593y	3 33 30·340 33 37·084		+ 2.1222	+ •0023				+11.019	- :257	- '043	21:23	8.36:8.32	827
18		6.2	11 Tauri	33 37 084 34 47 827		-2.2950 + 3.5758	•2315 •0188		-78 41 11.92 +25 0 22.23		11.923 11.853	+ ·264 - ·425	- ·028 - ·015	48:49 18	8.04 : 7.95 9.25	828 836
18		3.8	38 Perseim. o	38 2.747	007	3'7524			+25 0 22 23 +31 58 17.63		11 053	- 425 ·450	- '024	21:22	8.50:8.40	844
19		5.1	Fornacisδ		+ .003	2.3847	•0023		-32 15 27.63	06	11.629		+ .007	20:21	8.03:7.94	846
19	I	3.7	23 Eridani	3 38 27.370	+ .052	+ 2.8719								18:19	9.11	848
19		3.8	17 Tauri	38 56.142	- ·014	3.222			-10 5 59.94 +23 47 55.81	- 6·77 + ·50	+12.322	- • 346 • 428	÷ ·743	16 : 19	9.97	852
19		2.8	25 Tauriη	41 32.306	010	3.5288			+23 47 45.36	+ .34	11 323	.432	048	23.	7'27:7'11	869
19		4'3	27 Eridaniτ <sup>6</sup>	42 32.622	+ .102	+ 2.5798	.0025		-23 32 47.03		10.792	- '314	523	19:18	8.87:8.86	873
19	5	8.1	Lacaille 1414	42 37.13		- 9.659	1.430		-85 2 47.76		11.310	+1.128		36:53	11.60 : 10.31	
19	6	3.8	Reticuli	3 42 57.046	- '437	+ 0.7377	+ .0282	+ .0472	-65 7 16.51	68	+11.359	099	+ .073	17	9.25	875
19		3.7	27 Tauri	43 12.866	013	3.2001				+ .43	II'217	•434	020	17:19	9.04:8.69	877
19		5.2	28 Eridani $\tau^7$	43 21.615	-	2.5789			-24 11 3.42	54	11.310		+ .054	16:17	9:94 : 9'91	880
19		4'2	Eridanig	45 42.692		2.2441			-36 30 10.96		11.036		020	26 : 28	6.98 : 6.94	888
20		2.8	44 Perseiζ	47 50.619	009	3.7622	•0220	+.0010	+31 35 12.34	+ .14	10.013	•464	- '017	20:22	8.50 : 8.45	894
-																
						164. 5	3, 11.5	4".6	250° 1896*	2.						1
		~				165. Al	B 6.7, 7.5 C 10.5	0".8	182° 1901'	Ι.						
1.	2.					189. 3.	8, 8.5	3″.5 o″.9	211° 1900'0 51° 1901''							
1000			-													

								_							
No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_{a}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annnal Variation 1900'0,	Sec. Var. 1900 o.	Proper Motion.	No. of Obs.	Epoch 1900 <b>+</b> .	Boss No.
								1							
201	3.1	Hydri	h m s 3 48 47 080	e - • 088	s - 0.9798	s + '1071	s +.0110	-74 32 42.84	- "92	+10.977	+ ".115	+ "116	21:22	8.01 : 7.92	899
202	5.3	Eridanii	49 50.317	- :024	+ 2.2854	.0026		-35 1 40.31	+ .15	10.765	286	018	16:17	8.43:8.53	903
203	6.4	Lacaille 1 304	51 56.348		1.2692	.0078		-52 58 54.90		10.658	•198		20 ; 22	7.48:7.32	N249
204	4° I	46 Perseiξ	52 28.422	010	3.8830			+35 30 13.22	+ .11	10.226		- '012	16:17	8.97:9.03	913
205	3.1	34 Eridani	53 21.847	040	2.7977	.0042	+ 0046	-13 47 35.21	+ .97	10.410	.352	115	22:24	8.62	915
206	var.	35 Tauriλ	3 55 8.330	+ .003	+ 3.3192	+ .0114	0004	+12 12 28.15	+ '12	+10.322	- '418	- ·014	21:23	8.03 : 8.28	920
207	7.4	Lacaille 1318	55 23.356	014	1.2160	.0001		-49 53 45.66	- ' .26	10.403		+ .035	20	7.99	922
208	4.7	36 Eridaniτ <sup>9</sup>	55 39.657	002	2.5562	.0035		-24 17 59.15	03	10.324	*324	+ .004	18:20	8.76 : 8.47	923
209	4.2	Reticuliδ	57 9.695	007	0.9406	.0194		-61 40 57.14		10.180	·122 ·404	- '019	16	9.55	930
210	4'0	38 Tauri	57 50.160	001	3.1826	*0092	1	+ 5 42 43.25	+ .02	10 180	404	- 007	23:26	7'95:7'61	932
211	4.2	37 TauriA	3 58 46 953	063	+ 3.2407	+ .0121		+21 48 31.25		+10.025	- '451	064	17	9.35	936
212	5'3	42 Tauri	4 0 49 342	+ .055	3'7019	.0187		+28 43 51.24	03	9.964	.472	+ .003	17	8.82	944
213	5'7	Lacaille 1344 43 Tauri		100	2°4713 3°4895	·0031 ·0136	+.0147	-27 55 30.79 +19 20 41.51	- '70 + '39	10°013 9°726	·319 ·449	+ .103	26:29 19:22	6.83 : 6.77 8.70 : 8.96	948 952
214 215	5.9 5.6	43 Taurip	3 20°395 4 44°319	+ '018	3 4095	'0168		+26 13 11.97	+ .33	9.625	•479	- '037	19.22	8.95	955
1.00									1						11.5-
216	6.2	Lacaille 1376 38 Eridaui		049	+ 1.8593	+ .0047	+.0001	-46 7 44°29 -7 5 53°01		+ 9.010	- '243	+ .000	25:26 20:22	6'94 : 6'92 7'58 : 7'47	959 963
217	4'2	38 Eridani	6 59°001 9 38°220	005	2.9263	•0050		-10 30 17.87	1	9°571 9°128	330	- 157	16	8.10	903
210	4.4	49 Tauri	10 6.193	- '017	3.2544	*0094		+ 8 38 30.59		9.229	•426	- '020	16	8.81	981
220	4'5	40 Eridani	10 38.714	1 '	2'7611	.0016	1484	- 7 49 3.01		5.770	•342	-3.435	21:18	9'70 : 9'47	984
	3.8	Horologiia		026	+ 1.9865	+ .0035	+.0037	-42 32 28.31		+ 8.988	262	- '215	26:28	7.07:7.00	985
22I 222	3.3	Reticuli	4 10 41 · 294 13 8 · 128	- '044	0.7624	0035		-62 43 26.36		9.071	104	+ '059	19	8.10	905
223	4.4	Doradûs		069	1.2672	.0079		-51 44 18.07		9.171	.209	+ .180	21:22	7.00:6.98	995
224	5.2	54 Persei		+ .023	3.8863	.0206		+34 19 31.57		8.937	.210	012	16	9.27	999
225	3.8	54 Tauri		072	3.4097	·0113	+.0081	+15 23 10.20	+ .23	8.910	.450	027	21	8.93 : 8.66	1000
226+	3.6	41 Eridanim. v <sup>4</sup>	4 14 6.650	- "042	+ 2.2688	+ .0031	+.0047	-34 2 32.54	+ .03	+ 8.933	300	003	16	8.96	1001
227	5.5	Lalande 8205		- '021	2.6170	.0037	+.0031	-20 52 41'18		8.760	• 347	005	26:28	6.84 : 6.72	1012
228	4'0	61 Tauriδ	17 10.021	061	3.4553	.0118	+.0011	+17 18 28.80	+ '26	8.663	.458	033	25:28	7.88 : 7.91	1017
229	4.4	68 Tauri		- '061	3°4661	.0112		+17 41 57.04		8.471	•462	025	17	8.09	1029
230	4.0	43 Eridanid	20 16.831	033	2'2517	.0033	+.0042	-34 14 55.80	40	8.202	• 302	+ .022	21:23	7'31:7'26	1032
231	5.3	Reticuli	4 20 48.442	102	+ 0.6383	+ .0237	+.0125	-63 37 23.54	- 1.22	+ 8.585	090	+ .122	19:20	8.59 : 8.58	1035
232	3.6	74 Taurie			3.4988			+18 57 31.22		8.213	•469	038	23:22	8.63 : 8.30	1044
233	5.4	Cœli			1.8321	.0048		-45 10 6.00		7.839		015		7.77:7.69	
234	4.8	86 Tauri	28 10.404	026	+ 3.4000	00100		+14 38 3.04		7.793	461	- '026	17	8.10	1067
235	7'9	Lacaille 1839			-17.030	2*287		-86 29 26.36		7.741			23:30	11'46 : 10'28	
236	0.9	87 Tauria		032	+ 3.4385	+ .0105		+16 18 28.58		+ 7.466	467	101	22	7.31	1077
237	4'I	48 Eridani	31 19.298	.000	2.9954	.0058		- 3 33 24.92		7.563	•408	002	19:17 18	9'48:9'34	1079
238	3.8	52 Eridaniυ <sup>2</sup> Doradûsα	31 39.698		2.3305	·0032		-30 46 1·33 -55 15 5·70		7.533	·318	004	16	9°29 10°04	1080
239 240	3.4	53 Eridani		- '061 + '050	1°2926 2°7456	·0097 ·0040	+ · 0001	-14 29 59.36		7.219		- '161	18	9.35	1001
														8.96	
241	6.9	Lacaille 1543	4 34 3'907	018	+ 1.9517	+ .0042				+ 7.388	$- \cdot 268$ + $\cdot 982$	+ .046	17	8'90	1094 1096
242 243	7°1 5·8	Lacaille 1544		+ .13	- 7.246	·527 ·0034	0010	-83 6 55.59		7.207	- · 342	+ .019	17	8.12	1104
243	4.3	94 Tauriτ			3.2962	0034		+22 45 54.55	1	7.142	*493	- '022	27:28	7.69:7.67	1107
245†	4.6	Cœlipr. a	37 20.277			.0040		-42 3 17.07		6.985	•265	089	20	8.18	1110
246	5.4	Pictoris		- T		+ .0068		-50 40 9.29		+ 6.870	- '212	+ .031	23	7.07 : 7.24	1119
240	4.2	57 Eridani			2.9981	·0054		- 3 26 16.56		6.805				8.10:8.01	1123
248	5.5	Doradûs			+ 0.8955	·0141		-59 54 57 73		6.659	126	+ .037	22	7.11:2.08	1130
249	5.9	Mensæ	44 3.621	+ .003	- 0.6222	.0477	0004	-71 6 51.21	- '24		+ .083			7.83:7.80	1138
250	3.5	I Orionisπ <sup>3</sup>	44 24.913	265	+ 3.2544	.0071	+.0316	+ 6 47 12.04	19	6.213	- '456	+ .050	19:18	8.40 : 8.31	1140
-				,						-					
-					206.	L, 3.8-4.2;	P, 3 <sup>d</sup> 95								

.

FROM OBSERVATIONS IN THE YEARS 1905-11.

.

															1
No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_{\alpha}\Delta E.$	Annual Variatiou 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900*0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
1	-										1				
							1			1.5.2					
251	5.2	97 Taurii	hm s	- ·050	+ 3. 5062	+ .0002	8	+18 40 10.95	+ ":30	+ 6:364	- "488	- "036	18:19	8.40 : 8.30	1143
	3.8	3 Orionis $\pi^4$	4 45 31.421	-		.0067		+ 5 26 3.34		6.365		006	26	6.77	
252	1			+ .005	3.1926				+ .04					8.22	1147
253	5.1	4 Orionis	46 52.460	.000	3.3907	.0083			+ '48	6.229		029	16		1149
254	3.8	8 Orionisπ <sup>5</sup>		+ .001	3.1558	•0060			+ .02	6.102	10.	003	25:26	7.02:6.88	1159
255	4.8	7 Orionisπ <sup>1</sup>	49 23.423	030	3.3004	.0072	+.0032	+ 9 59 29.73	+ 1.08	5.945	•462	- 134	22:23	8.08:8.06	1163
256	2.8	3 Aurigæ	4 50 28.785	005	+ 3.0010	+ .0141	+.0007	+33 0 28.90	+ .19	+ 5.961	545	027	23:24	7.03:7.00	1167
257	5.9	98 Taurik		018	3.6677	.0108	5	+24 53 45.91	+ .48	5.798		060	21:22	8.00 : 7.97	1177
258	4.8	102 Tauri	57 7.063		+ 3.5825	.0002		+21 26 49 95	+ '32	5.385		047	26:27	6.88 : 6.86	1194
259	5.4	Mensæ		010	- 1.7626	.0726		-75 5 25.91	- '44	5.406	+ '245		21:22	8.06	1197
260	4.8	11 Orionis.		-	+ 3.4255				+ '31			036	19:20	8.57:8.55	1203
200	40		4 50 51 215			0070	T 0013	T-15 15 55 32	T 31	5°249	404	- 030	19:20	0 57 . 0 55	1203
261	5.6	Pictoris	5 0 11.687	+ .038	+ 1.5659	+ .0056	0056	-49 17 33.86	09	+ 5.185	- '222	+ .013	22	6.72	1207
262	3.2	2 Leporis	I 13.642 ·	012	2.5386	.0031	+.0010	-22 30 19.60	+ .54	5.018	• 360	067	23:25	7 .93 : 8 .08	1211
263	5.0	Pictoris	2 22.572	024	1.5488	.0056	+.0035	-49 42 47.96	05	4 '995	.221	+ .007	20	6.85	1218
264	2.8	67 Eridani		+ .052	2.9482	.0043		- 5 12 57.36		4.861	.418	079	19	8.84	1220
265	4.8	Doradûs	00 00		+ 1.0233	1010.	0054		88	4*972		+ 105	20:22	8.40 : 8.37	1225
-					1		54				1				
266	5'4	Mensæβ	5 4 0.390 -		- 0'7945	+ .0398	- '0022	-71 27 2.92	38	+ 4.896		+ .047	19	8.13	1228
267	4'3	69 Eridaniλ		005	+ 2.8698	.0040	+.0002	- 8 52 56.18	+ .00	4.811	408	- '008	23	7.76	1231
268	3.5	5 Leporis	8 26.349 -	- '023	2.6936	.0033	+.0028	-16 19 26.02	+ '22	4.444	*385	028	27:26	8.11: 2.03	1241
269	0.0	19 Orionisβ	9 43 915 -	001	2.8817	.0039	+.0001	- 8 19 1.71	+ .01	4.361	·412	- '001	31:32	7.11: 7.07	1250
270	3.7	20 Orionis	12 45.028 -	+ .008	+ 2.9117	.0039	0011	- 6 57 8.88	+ .05	4.097	- '417	002	24:26	7'20:7'19	1262
	0														
271	4.8	Doradûsθ			- 0.0286	+ .0210	0010	-67 17 51.91	36			+ .048	21	7.60	1269
272	5.0	Columbæo		023	+ 2.1655	.0024	+.0062	-34 59 37.63	+ 2.81	3.661		- '346	20	8.15	1270
273	7.8*	Cape 1880. 2449	13 56.57		-34.228	3.972		-87 59 21.25			+4.888		31:41	11'43 : 10'68	
274	4'3	6 Leporisλ	14 58.045		+ 2.7630	.0034	.0000	-13 16 48.07	03	3.012	— ·397	+ .003	16	8.43	1277
275	5.9	Lacaille 1796	15 24.205 -	+ .000	2.3898	.0029	0008	-27 28 17.74	+ .10	3.862	*343	014	22:23	6.97 : 6.93	1279
276	4.7	22 Orionis	5 16 39.407 -	+ .003	+ 3.0010	1 100.00	10001	- 0 28 52.03	1 100	+ 3.766	440	003	27:28	8.62 : 8.59	1284
	5.7	Pictoris				+ '0042	0003		+ '03			+ .216			1287
277	7'2	Lacaille 1836	51.5.5	004	1.4681	.0029	+.0006		- 1.62	3.963			25:26	7'44 : 7'48 6'44 : 6'41	1298
278				- '024	1.4124	*0055	+ 0037	-51 40 20.50	12	3.288		+ .023	23:24		
279	4'9	25 Orionis		+ .002	3.1110	*0042			+ '14	3.201		018	18:19	7.76	1302
280	1.0	24 Orionis	19 46.018 -	+ .004	3.5125	.0046	0002	+ 6 15 32.91	+ '14	3.482	•463	019	20:22	7'81 : 7'61	1303
281	1.6	112 Tauri	5 19 58.197 -	020	+ 3.7900	+ .0076	+.0024	+28 31 21.69	+ 1.48	+ 3.307	546	177	19	8.35	1 304
282	6·1	Lacaille 1850			1.7840			-44 18 51.83		3.308		005	29:30	7'10:7'14	1317
283	6.1	Lacaille 1862,	0 . 0		1.9229			-41 1 46.78		3.234		+ .088	26': 27	7.04:7.02	1322
284†	2.7	9 Leporisseq. B		003		'0027						- '094	25	6.81	1323
285	4'9	25 Aurigæχ		-	2.2703					3.045		016	-	8'14 : 8'18	
	49		20 13 0/0	co4	3.9027	•0076	T 0005		+ .13	2.928	503	010	20:19		1333
286	2'2	34 Orionisδ	5 26 53.838 -	100	+ 3.0638	+ .0036	+.0001	- 0 22 23.26	+ .02	+ 2.882		003	27:28	7.37:7.38	1339
287	5.7	Lacaille 1888	27 24 497 -	+ .003	1.6458	*0035	0004	-47 8 59.60	+ 1.13	2.688		123	20	7.40	1341
288	3.9	Columbæe		019	2'1292	.0028		-35 32 37.69		2.770		049	18:19	8.60	1344
289	2.6	11 Leporisa		001	2.6451	.0029		-17 53 37.65		2.765	*383	+ .003	20	7.42	1347
290	4.5	37 Orionis	29 19.805 -		3'2920	.0042		+ 9 25 18.96		2.667		008	23	7.25	1353
1										1000					
291	5.2	43 Orionisθ <sup>2</sup>	5 30 28.219 -	002	+ 2.9460	+ .0035		- 5 28 54 54	11	+ 2.201		+ .012	18:19	7.35	1365
292	2.9	44 Orionis	0 0 .0	- '002	2.9338	.0035			+ .03	2.266	•425	004	20	7'93:7'84	1366
293	1.0	46 Orionise	31 8.332	.000	3.0430	•0034		— I I5 56.67		2.216		- '002	19	8.10	1370
294	3.0	123 Tauri	31 40.049 -	- '002	3.2838	.0021	+.0002	+21 4 53.84	+ .23	2.444	.519	028	20 : 22	8.30:8.09	1375
295	3.8	Doradûsβ	32 45 336 -	+ .000	0.2101	.0001	0011	-62 33 18.47	- '11	2.391	•076	+ '014	20	7.90	1384
2967	3.8	48 Orionism. σ					1. 2.0							6.80 . 6.00	1389
			5 33 43 531	.000	+ 3.0102		.0000	- 2 39 27.73	01	+ 2.294		100.	26 : 27	6.80:6.77	
297	5.2	Mensoe	000	- '207	- 2'4001	•0459		-76 24 42.08	- 2.34	2.410		+ .301	40	7.88:7.78	1400
298	2.6	Columbæa	36 1.641 -		+ 2.1718	.0022		-34 7 38.20	+ '24	2.058	-	032	28:31	6.93 : 6.94	1401
299	3.7	13 Leporisγ	40 17.483 -		2'5012	'0020			+ 2.58	1.346		376	32:34	6.81 : 6.86	1420
300	6.6	Lacaille 1981	40 50.875 -	008	1.2005	.0032	+.0011	-45 52 42.31	62	1.201	•248	+ .088	26	7*08	1421
-	-														
1															1. 1. 1.

284. 2'7, 9'6 2"'7 296° 1898'I. 296. 3'8, 5'7 ; very close binary.

.

.

.

-															
No.	Mag.	Name.	Mean R.A. 1900°0.	$\mu_{\alpha}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
			h m a	8	s	s	s							1	
301	5.7	130 Tauri	5 41 36.305	·001	+ 3.4975	+ .0032	+.0005	+17 41 29.87	+ ".07	+ 1.597	509	- "010	23:24	7.43:7.44	1424
302	3.6	14 Laporis		+ .010	2.7176	.0025	0015	-14 51 33.14	02	1.238	*395	+ .005	33:32	7.96 : 7.88	1432
303	2°I	53 Orionis	43 0.807	- '002	2.8446	*0026		- 9 42 18.49	+ .04	1.429	.414	002	20:22	7.52	1435
304	7.3	Lacaille 2005	44 20.666		1.8865	.0028	0010		+ '24	1.332	.275	033	21:22	7'34:7'41	1440
305	4.2	Doradûsδ	44 35 571	+ .023	0.1058	.0082	0062	-65 46 22.67	08	1.322	.012	+ .010	17	8.48	1443
306	3.9	15 Leporisδ	5 47 1 361	146	+ 2.5804	+ .0014	+.0172	-20 53 20.43	+ 5.49	+ 0.482	378	653	19:21	8.47 : 8.41	1456
307	3.0	Columbæß	47 26.093		2.1136	.0033	+.0039	-35 48 17.62	- 3.43	1.493	.309	+ '394	17	8.70	1459
308	4.4	Pictoris		076	1.0875	.0038	+.0082	-56 11 30.48	+ .63	0.980	.160	068	17	9.26	1460
309	7.3	Lacailla 2040		- '024	+ 1.9092	*0027	+.0025	-41 7 44.78	12	1.000	279	+ .010	16	9.20	1463
310	6.2	Lacaille 2296	49 33.79	+ .12	-11.211	.152	013	-84 50 6.23	72	0.999	+1.402	+ .086	41 : 144	11.45 : 8.38	N381
311	var.	58 Orionia	5 49 45 489	012	+ 3.2474	+ .0026	+.0010	+ 7 23 18.53	06	+ 0'904	474	+ .008	19:20	7'93:7'86	1468
312	4.9	139 Tauri		+ .001	3'7220	.0028	0001		+03	0.214	•543	- '004	22:23	7.91:7.88	1475
313	3.7	16 Leporisη	51 50.977	+ .010	2.7319	.0024	0028	-14 11 8.38	00	0.845	•398	+ '132	32	6.81	1476
314	4.4	Columbæ	53 59.465	+ '002	2.1263	'0024		-35 17 38.01	02	0.528	.310	+ .002	24:25	7.98:8.00	1490
315	3.9	Columbæ	56 5.135	- '014	1.8359	*0025	1.	-42 49 14.57	+ '19	0.312	• 268	025	22	7.53	1497
	X 21				-								10 . S		-
316	4'2	61 Orionis μ	5 56 52.861	010	+ 3.3002	+ .0020		+ 9 38 50.26	+ .24	+ 0.244	- '481	- '029	17:18	8.39	1501
317	5.7	Mensæ	57 1.438	+ .026	- 4.0604	.0148	0067	-79 22 42.59	- '48	0.312	+ . 593	+ .057	37:38	8.41 : 8.42	1502
318	4°3 7°2*	1 Geminorum		+ .002	+ 3.6464	.0012			+ .96	0.063	532	108	20	8.89	1508
319		Cape 1880. 2901			-43.595	.122		-88 21 34.71			+6.357		34:41	11'47 : 10'90	
320	5.9	66 Orionis	5 59 41.298	+ '002	+ 3.1694	.0018	-*0002	+ 4 9 51.80	+ .10	0.012	- '462	015	17:18	8.43	1514
321	6.6	Lacaille 2137	6 I 35.637	+ .070	+ 1.7262	+ .0030	0080	-45 2 7.80	- 2'00	+ 0.089	250	+ .229	17	8.73	1521
322	4.4	67 Orionis	1 51.723	005	3.4256	.0012	+.0006	+14 46 49.57	+ •27	- 0.199	•499	036	23:25	7.68:7.53	1525
323	5.9	Lacaille 2130	2 14.483	017	2.3108	'0021	+.0022		+ '33	0.238	*337	042	19:21	7.87 : 7.91	1528
324	5.7	Columbæ $\pi^2$	4 46.531	+ .002	1.8631	.0022	0003	-42 8 17.70	+ .10	0.433	.272	012	26:27	6.99 : 6.96	1539
325*	5.1	Lacaille 1766	6 8.590	021	+ 0.2480	+ '0012	+.0028	-62 8 12.38	+ .23	0.608	•080	071	20	7'43	1546
326	7.1	Lacaille 2512	6 6 9.38	+ .30	-15.735	- 117	026	-85 55 52.80	01	- 0.238	+2.297	+ '001	35:95	11.42 : 8.96	1547
327	4.4	70 Orionisξ	6 15.214	005	+ 3.4118	+ .0011	+.0000	+14 13 52.62	+ .28	0.281	- '497	034	17:18	8.34 : 8.36	1548
328	5.6	Lacaille 2182		+ .030	1.9339	.0023	0042	-40 20 5.66	38	0.554	•281	+ .053	24	7.22	1553
329	6.4	Lacaille 2191	7 47 393	+ .009	1.7234	·0021	1	-45 15 35.06	01	0.680	.250	+ .001	18:20	8.02:7.91	1555
330	4.9	Pictorisδ		+ .018	1.1662	·0018	- '0025		+ .06	0.738	.169	008	20	7.38	1558
331+	var.	7 Geminorumseq. η	6 9 101156	1	1	1	100.17			0.000	526	012	11.17	7.11 . 7.10	1561
332	5.3	74 Orionisk		+ .032	+ 3.6220	+ .0002		+22 32 9.35	+ .12	- 0°790 0°754	- 520 ·491		44:47 22:23	7'11 : 7'12 7'85 : 7'84	1577
333	4.5	Columbæĸ		- '047	3.3694				- 1.21		- '310			6.92:6.88	1587
334	5.2	Mensæa	50 0		- 1.7838			-35 6 24·97 -74 43 8·82			+ .256	- '215	31	7'02	1589
335	5.3	7 Monocerotis		- ·192			0002		+ 1.21	1 3/0	- '420	002	23:24	7.87:7.86	1598
	100.0														
336	3.0 ,	I Canis Majoris	6 16 28.431		+ 2.3027			-30 I 7°93	- '01	- 1.439	- '334	+ .001	31:32	6.99 : 6.96	1601
337	3.0	13 Geminorumµ		031				+22 33 53.20	+ .78	1.201	.528	- 113	21:22	6.97 : 6.91	1604
338	1.8	2 Canis Majoris	18 17.711		2.6414			-17 54 22.68	.00	1.200	•383	.000	31:33	8.66 : 8.60	1609
339	4.5	8 Monocerotis	18 28.126	1.	3.1292				+ .02	1.617	•461	003	· 28	7.06	1611
340*	-0.9	Argûsa	21 43 923	013	1.3313	.0009	+.0012	-52 38 27.05	- '08	1.888	.195	+ .010	22	7.76	1622
341	5.1	10 Monocerotia	6 23 1.265	+ .001		+ .0000	0002	- 4 42 1'22	10	- 1.996		+ .014	36:37	7'18:7'22	1634
342	4*1	18 Geminorum	23 1.207	+ .002	3.2630	0011	0006	+20 16 31.88	+ .12	2.032	.216	021	16:17	8.05 : 8.04	1635
343	4.2	Canis Majorisλ	24 27 690	+ .012	+ 2.2230			-32 31 1.18	11	2.150		+ .019	27	6.92	1641
344	5.2	Doradûs $\pi^2$	26 19.818	+ .034	- 0.2089			-69 37 57.54	- 1.32		+ .076		29	6.40	1648
345	4.8	13 Monocerotis	27 29.776	- '002	+ 3°2450	0003	+.0002	+ 7 24 22.46	+ .08	2.409	- •469	010	28	7'90	1657
346	5.8	Lacaille 2349	6 28 58 221	058	+ 1.3986	+ .0002	+.0084	-51 45 20.24	64	- 2.435	202	+ .092	34	6.96	1671
347	4.6	5 Canis Majoris	30 51.882	003	2.5137			-22 53 7.69	09	2.678	•362		40:41	7.18:7.20	1682
348	1.8	24 Geminorum		022	3.4671				+ .33	2.831	.500	047	27	6.99	1690
349	4.4	CarinæN		+ .012	1'3220			-52 53 37.89		2.866		009	23	7.30	1696
350	3.1	Argûs		- '001	1.8358			-43 6 29.47		3.044			35:36	7.02:7.06	1702
	1				0.00	1									
			2 2 2 2 2												

311. L, 0.6-1.1; P, irregular. 325. Lacaille's R.A. is 1<sup>h</sup> too small. The fictitious  $\mu$  Doradûs. 331. L, 3'2-4'2; P, 233. 331. Var., 8'8; close binary. 340. Magnitude from Harvard Annals, vol. l.

. .

9

.

8

.

FROM	OBSERV	VATIONS	IN	THE	YEARS	1905-11.
------	--------	---------	----	-----	-------	----------

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_{\alpha}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900 '0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
351+	5.2	15 Monocerotisseq.	h m s 6 35 28.249	8	s + 3.3021	s - '0012	s + .0002	+ 9 59 17.60	+ "05	- 3'097	- ":475	- "007	32:35	7'16 : 7'24	1706
352	3'1	27 Geminorume	37 46.767	•000	3.6936			+25 13 48.93	+ .14	3.310	.530	020	30	6.99	1717
353	6.9	Brisbane 1331	38 4.203	+ .002	1.6312			-47 31 35.17	- '08	3.304	• 233	+ .011	32:34	7.28 : 7.20	1719
354	3.3	31 Geminorumξ		+ .055	3.3686		1	+13 0 11.51	+ 1.42	3.654	•481	- '201	23:26	7'10:7'05	1725
355*	-2.0	9 Canis Majorisa		+ .257	2.6441	1.1.1.1	0366	-16 34 51 93	+ 8.43	4.751	•372	-1.500	38:39	7'03: 6'99	1732
356	4.8	18 Monocerotis		+ .003	+ 3.1298	- •0008		+ 2 31 17.99	+ .17	- 3.734	- '447	- '025	42:44	6.94 : 6.93	1740
357 358	3.8 3.6	13 Canis Majorisк 34 Geminorumθ	46 6°286 46 11°S61	+ .000	2·2407 3·9588	+ ·0014 - ·0075	-·0008 +·0005	-32 23 34.58 +34 4 54.65	- ·01 + ·37	4.005	·318 ·564	+ '001	34:35	7 °05 6 °94 : 6 °85	1761 1763
359	3'2	Pietorisa	47 9.787	+ .082	0.6176	0020	-	-61 49 59.84	- 1.86	3.837		+ .200	24.25	7.23:7.15	1769
360	2.7	Argûsτ	47 27.278	- '021	1.4886			-50 29 43.58	+ .68	4.207	.211	086	22:23	7 .95 : 7 .88	1772
361	4.3	14 Canis Majoris	6 49 32.557	+ .068	+ 2.7876	+ .0003	0003	-11 54 48.12	+ .11	- 4'315	394	012	44:49	7.28:7.19	1783
362	6.2	Lacaille 2530	51 17.762		+ 1.8903			-42 14 19.58	- '12	4.433	267	+ .017	29	7.06	1790
363	4.4	20 Canis Majoris	51 40.602		+ 2.6759	+ .0002	0001	-16 55 28.78	09	4.471	- '378	+ .011	22:23	7 '94 : 7 '92	1793
364	5.6	Volantis	0 00 00	+ '012		- '0272	0012	-70 50 19.52	20	4.532	+ .008	+ .028	29:30	7.18:7.12	1795
365	5.8	Piazzi VI. 303	54 29.989	+ .000	+ 2.4577	+ .0015	0013	-25 16 42.07	10	4.708	- '346	+ .014	36:37	7'05 : 7'08	1802
366	1.2	21 Canis Majorise	6 54 41 . 720				+.0001	-28 50 9.22	+ .01	- 4.740	332	001	29	6.88	1804
367	6.1	Piazzi VI. 305		- '095	3.8173			+29 30 11.36	+ 6.49	5.771	*539	823	22	7.88	1809 1810
368 369	3.8 var.	22 Canis Majoris 43 Geminorum ζ		+ .006	2°3894 3°5613	+ .0012		-27 47 29'74 +20 43 1'57	+ .02	4.996 5.042	°335 °500	+ ·001	25 31:33	7 °97 6 °53 : 6 °44	1815
370	3.0	24 Canis Majoris		+ .004		+ .0010	0002	-23 41 13.65	+ .05	5.097	•351	006	18	8.12	1817
371	4.1	23 Canis Majoris	6 59 14.041	+ .001	+ 2.7143	f .0004	0001	-15 29 7.51	+ .10	- 5.138	- '380	014	34:36	7 '59 : 7 '45	1819
372	5.4	Lacaille 2642		+ .011	1.1180			-56 35 51.99	01	5'392	.155	+ .002	32:35	7 '01 : 6'90	1833
3731	5.7	45 Geminorumpr.		+ .002	3.4428	0047	-	+16 5 24.72	+ .86	5.522	•481	111	18	7.74	1835
374	7°1	Lacaille 2631	2 47 204	+ '023		+ .0002	0035	-42 10 25.62	06	5.415		+ .000	22:23	7.27 : 7.21	1836
375	5.6	Mensæ $\dots$ $\theta$	2 53 902	+ .048	- 3.7266	- '1452	- * 0058	-79 16 35.96	+ .02	5.436	+ .225	003	44	8*23	1837
376	1.8	25 Canis Majoris	7 4 19 505	+ .003	+ 2.4391	+ .0011	0004	-26 14 3.85	01	- 5.551	- *339	+ .005	27:28	7'13:7'17	1839
377	6.1	Lacaille 2651		+ .024	1.4375		0033	-51 48 40.17	38	5.542	• 198	+ .053	32:33	7*23:7*17	1842
378	4.2	22 Monocerotis 51 Geminorum		- ·002	3.0652		+.0002	-01937.67 +161943.04	09	5°746 5°880	•426 - •478	+ .011	26 : 27 24 : 26	7 °98 8 63 : 8 79	1853 1856
379 380	5°3 3°8	Volantis	9 35.790	- '007 - '040	+ 3°4478 - 0°4916	-	+.0008	-70 20 10'36	+ •44	5.901	+ .070	+ .094	24 . 20	7.90:7.88	1867
1.000				1.1.1.1										6.88 : 6.80	
381 382	4°5 3°5	PuppisΙ 54 Geminorumλ		+ .098	+ 1.7100 3.4207	·0000 - ·0057	0143	-46 35 30'97 +16 43 14'98	$- \cdot 61 + \cdot 33$	- 5.915 6.272	- ·233 ·475	+ .089	34:36 36:35	7.03:6.95	1869 1886
383	2.5	<b>Arg</b> ûsπ						-36 55 3.87		6.331	•290	- '002	30:31	7.14:7.07	1896
384†	3.4	55 Geminorumseq. 8	14 9.042		3.2873	0074	0013	+22 9 59.51	+ .12	6.390	•493	012	24:25	7.27 : 7.22	1598
385	5.0	29 Canis Majoris	14 30°513	+ .002	+ 2.4975	+ .0008	0010	-24 22 34.37	+ .06	6.411	- '342	008	21:22	7.08:2.02	1899
386	3.9	Volantisδ		+ .025	- 0.0195	0253	0035	-67 46 26.70	+ .04	- 6.605	+ .000	006	30	7.09	1917
387	3.9	60 Geminorum	19 30.915			- *0104			+ .63	6.907	- • 508	090	25:26	7.06:6.96	1931
388	2.3	31 Canis Majorisη 3 Canis Minorisβ				1100. +	0007	-29 6 29.03 + 8 29 26.96	- "03			+ .004	22	7.61	1934
389 390	6.7	Lacaille 3274	21 43 <sup>.</sup> 647 22 1 <sup>.</sup> 70	+ .026 03	+ 3.2559 -19.815	- °0043 - 2°648	-•0034 +•003	-86 52 11.36	+ 33 - 06		- °441 +2°709	043 + .007	26 : 30 47 : 150	7'79:7'73 11'31:8'70	1944 1947
	1.2.1	-													
391 392	4°3 5°2	62 Geminornmρ Lacaille 2829	7 22 40·866 23 47·909	- <sup>•</sup> 093 + <sup>•</sup> 004	+ 3.8649		+ °01 17 - °0006	+31 59 1.36	- 1°46 + °02	- 6·893 7·171	- • 526 • 207	+ .183 003	20 32:35	7°97 7°18 : 7°10	19 <b>5</b> 2 1960
392	4.9	6 Canis Minoris		002	3.3425			+12 12 48.20	+ .12	7.222	.452	019	32·35 19:21	7.85:7.84	1962
394	2.9	Argûsσ		+ .046	1			-43 5 54.50	- 1.43	7.172		+ .180	27	7.97	1972
395	4'2	69 Geminorum	29 45.611	+ .012	3.7029	- '0114	-'0020	+27 7 4.63	+ .89	7.768	•495	119	21	7.21	1987
396	4.6	Lalande 14810		+ .034	+ 2.5665	+ .0006	0048	-22 4 47.96	- '32	- 7.608	342	+ .042	41:42	7.07 : 7.04	1988
397	5.2	25 Monocerotis		+ .037	2.9841			- 3 53 15.67	- '14	7.840		+ .018	55:67	7.78	1999
398	5.1	71 GeminorumQ		+ .012	3.9246			+34 48 49.11		8.006	.522	- '122	17	7.90	2001
399 400	4°9 4°7	Puppis		- ·010 + ·022	1°4850 2°2189		and the second se	-52 18 37°99 -34 44 36°67	+ '14 - '10	7°949 7°953	-	- '021 + '014	23:24 20:21	6'95 : 6'87 7'46 : 7'36	2003 2004
		11	55 4 5		- 2109	1 0012	0029	54 44 30 07		1 933	-93			14-1755	
					351. 5	.2, 8.0	2"*9 2	17° 1902'7.							

В

•

\$

No.	Mag.	Name.	Mean R.A. 1900'0.	μ <sub>α</sub> ΔΕ.	Annual Variation 1900°0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900°0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
401*	0.5	10 Canis Minorisα	hm s 7343.724	+ · 325	s + 3.1433	s '0055	s -••0466	+ 5 28 44.79	+ 7.20	- 9'029	- "409	-1.030	24:26	6.98 : 6.99	2008
402	4.1	26 Monocerotis		+ .043	2.8671		0021	- 9 19 4.39	+ '20	8.215	*377	- '024	33:35	8.46 : 8.53	2008
403†	3.5	77 Geminorumseq. к	38 24.656		3.6280			+24 38 16.14	+ .46	8.408	•477	062	18:20	7.61:7.41	2029
404	1.1	78 Geminorumβ	39 11.442		3.6776	0128		+28 16 3.89	+ .47	8.466	•477	058	17	8.09	2031
405	4.1	3 Puppisl	39 47 560	+ .005	2.4083	+ .0011	- '0002	-28 42 56.48	+ .08	8.466	•314	010	16:17	8.62 : 8.48	2035
406	5.1	Lacaille 2945	7 40 17.706	085	+ 2.0427	+ .0005	+.0115	-40 41 22.81	+ 1.43	- 8.685	268	189	19	7.58	2039
407	5.4	80 Geminorumπ	41 3.243	005	3.8775	0165	+.0003	+33 39 40.35	+ .28	8.596	• 507	040	22:23	7'21:7'12	2049
408	5.2	4 Puppis		+ .000	+ 2.7628	.0004	0015	-14 19 14.57	- '02	8.576		+ .003	22	7.87	2051
409	8.4*	Gilliss P.Z. 5752	41 31.05		-86.829	-54·772 + '0011		-89 13 49.78			+11.443		13:16	11.39:11.41	
410	3.5	Puppisc	41 41 499	+ .012	+ 2.1365		0051	-37 43 32.86	+ .06	8.614	- *277	008	18:20	7.02:6.90	2052
411	3.8	Volantis	7 43 3.121	014	- 0.7121		+.0050	-72 21 56.86	01		+ .092	+ .005	26 : 29	7 '04 : 7 '02	2056
412†	3.4	7 Argûsseq. ξ	45 5-299	1	+ 2.5229	+ .0008	0002	-24 36 31.57	- ·01	8.872	-	+ .001	38:43	7.02:6.96	2065
413 414 <sup>†</sup>	4°2 5°5	PuppisP 9 Puppism		$+ \cdot 008$ $+ \cdot 034$	1.8280 2.7789	.0000	-·0011	-46 7 16·59 -13 38 0·70	+ .06 + 2.88	8.969	•234	009	27:28 20:23	7.08:7.00	2070
415	5°I	83 Geminorumφ	47 22.635		3.6784	0133		+27 1 28.36	+ '30	9°373 9°090	· 357 · 474	- '339 - '037	17	8.16	2075 2078
	3.6							10 m m m m m							
416 '	5.9	Puppisa Lacaille 3083a		T 011	+ 2.0620	+ •0009 - •0247	-'0016	-40 19 4·19 -65 56 24·64	+ •05	- 9°170 9°180	- ·263 ·049	- •008	22:23 21	6.70 : 6.64 7.24	2087
418*	4'3	Lacaille 3068		+ .001	1.7639	.0007	0002	-47 50 31.27	+ .13	9.303	•224	018	31:33	7'42:7'31	2095
419	6.1	1 Cancri		+ .012	+ 3.4106	·0086	0019		+ .39	9.407	436	049	27:29	8.00	2098
420	420       8·1       OctautisA       53       2·01       + '49 $-44^{\cdot}246$ 16·886 $-^{\cdot}043$ $-88$ 34 $24^{\cdot}36$ $-^{\cdot}09$ 9·481 $+5^{\cdot}690$ $+^{\cdot}009$ 46:98       11·40:9·49       210         421       3·5       Argus														
421	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$														
422	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$														
423	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$														
424	5.5	Geminorumx			3.6922	0120			+ .37	9.876	•465		45:49	7'25:7'18	2131
425	2.0	<b>Argûs</b> ζ	8 0 4.164	+ '022	2.1079	+ .0013	0030	-39 43 16.61	02	10.025	• 261	+ .002	47 : 51	7'33:7'21	2141
426	5.2	10 Cancri	8 1 52.814	014	+ 3.5365	- '0120	+.0018	+21 52 19.25	+ .64		440	081	24:26	7.85:7.86	2146
427	8.2	Brisbane 2007	3 3.05		-12.040	- I·773		-85 39 16.05			+1.213		41:48	11.32 : 10.82	
428*	2.8	15 Argûs	3 17.060	+ .020	+ 2.5545	+ .0010		-24 0 57.05	34	10.226	- '314	+ .042	43:53	7.63:7.59	2153
429	1.6	14 Cancri	4 25°782 6 27°024	+ .041	3.6207 1.8496	- °0147 °0000	-·0051 -·0004	+25 48 36.62	+ 2.84	10.211	·447 ·225	- ·354 - ·003	26 36:39	8°03 7°14 : 7°07	2157 2167
								-47 2 30.34							
431	5.5	20 Puppis			+ 2.7580	0004	0000	-15 29 12.63	+ .02	-10.682	- :336	002	62:68	7.84 : 7.73	2183
432	3·7 5·3	17 Cancriβ 18 Cancriχ	11 5°532 13 59°424		3°2568 3°6525	·0072 ·0167		+ 9 29 37 41 + 27 32 26 .74	+ '41	10.906 11.452	*394 *439	- ·054 - ·388	54 : 60 28	7:49:7:53 7:89	2195 2202
433 434	33 4°5	Puppisg	13 59 424			+ '0020		$-36\ 20\ 56.85$	59	11 452	• 439	+ .085		6.88 : 6.99	2202
435	6.1	20 Cancrid <sup>1</sup>	17 38.294		3.4406	- '0114	-	+18 39 11.82		11.301	•408	032	24:25	7.76	2218
436	1.4	Argûse			+ 1.2356	0001	0038			-11.218	142	+ .014	22.24	7 .92 : 7 .97	2222
430	3.9	Bradley 1197	20 39.812		2.9996	.0033	0039	-59 11 15.39 -3 34 48.92	+ .10	11.210	- ·142 ·352	$+ \cdot 014$ $- \cdot 025$	23:24 16:20	7.65:7.43	2233
438	6.2	29 Caneri	23 2.558		+ 3.3527	·0098		+14 32 31.07	+ .14	11.733	- '392	018	25:26	7.87:7.90	2253
439	4.5	Chamæleontis $\theta$	23 38.45	+ .21	- 1.7221	• 1648	0447	-77 9 43.74	- '20	11.736	+ .214	+ .025	32:78	11.33 : 9.03	2255
440	3.2	Volantisβ	24 38.953	+ .048	+ 0.6644	°0266	0062	-65 48 11.85	+ 1.18	11.992	072	- •166	24:26	7.20 : 7.08	2258
441	7.0	Lacaille 3353	8 25 22.620	+ .017	+ 2.0921	+ .0019	0023	-42 15 15.43	04	-11.874	241	+ .000	24:25	7.26:7.35	2262
442	5.8	31 Cancri	25 53.641		3.4268	0119	0032	+18 25 56.57	+ .54	11.986	• 396	069	18	7.89	2265
443	6.3	Lacaille 3368				+ .0011		-45 59 48.25		11.974	•223	012	23:26	7.06 : 2.01	2270
. 444	5.7	33 Cancri	26 55 576		3.4762	- *0132		+20 46 51.23	+ .40	12.044	•401	054	32:30	7.52:7.42	2271
445	4*2	4 Hydræδ	32 21 7 30		3.1290	0066			+ -10	12.378	•359	011	33:39	8.45 : 8.73	2295
446	6.1	Lacaille 3443			+ 1.7923				+ .08	-12.414	- '200	011	26:28	6.93 : 7.04	2297
447	4.6	5 Hydræσ	33 31.857					+ 3 41 32.89		12'467	* 353	020	16	7.94	2302
448 449	4°1 5°3	Velorume 6 Hydræ	34 7°659 35 17°150		2°1084 2°8428			-42 38 20·84 -12 7 18·61		12°494 12°571	·235 ·317	- •006 - •004	22 28 : 29	7.41 8.05 : 8.06	2307 2315
449	3.9	Pyxidis	36 11.285		2.3472			-34 57 12.06		12.648	• 260	019	29:29	6.95 : 6.94	2318
	1		<u> </u>	- 5	017					-					
					401. 3	Reduction to		0 <sup>8</sup> 017, + 1"*22.							SPITE
						3'5, 8'0 3'4, 13'7		236° 1903'3. 224° 1898'3.							1000
					414. 6	o•o, 6•6; ▼	ery close l	oinary.							
1						S Navis : in		ria Argentina. Bradley.							6
	-					-									

FROM OBSERVATIONS IN THE YEARS 1905-11.

.

No.	Mag.	Name.	Mean R.A.	μ <sub>α</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var.	Proper Motion.	Mean Dec.	$\mu_{\delta}\Delta E.$	Annual Variation	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch	Boss No.
			1900'0.		1900 0.	1900 0.		1900'0.		1900'0.	1900-0.			1900+.	
			h m s	s	S	S	s					\$			1
451	3.6	Velorumb	8 37 18.459		+ 1.9901		0002	-46 17 34.26	+ ".11	-12-718	- "219	- "013	20	8.09	2324
452	4'7	43 Cancriγ	37 29.928	+ .057	3.4788	0143		+21 49 41.26	+ '39	12.768	• 386	020	17	7.81	2327
453	4'4	Carinæd 47 Cancri	38 <b>24</b> ·438 39 0·184	+ .022	1.3277	- ·0082 - ·0128	0032	-59 24 14.61 +18 31 17.71	+ .04	12.785	•143	- '006	27	7.02	2331
454 455	4'I 7'0	47 Canell		T 008	-12.466	- 2.640	1	-86 13 22.21	T 1 03	13.058 12.849	- '377 +1·401	- •239	23:24 52:63	6·99 : 6·83	2336
			ESCINSION IN		1.000		1000								
456	3.6	Pyxidisa VelorumD	8 39 34 409	+ .007	+ 2.4100	+ .0010	- '0009	-32 49 33°07 -49 27 39°70	05	-12.851	- • 264	+ .000	23	7.75	2342
457 458	5°2 4°2	48 Cancri	40 32.495 40 38.814	+ '011	3.6399	- '0195		+29 7 32.67	- ·04 + ·36	12.917 12.979	·204 ·400	+ .002	26:27 16	7.33:7.27	2347 2348
459†	3.4	11 Hydræ. AB	41 28.759	+ .094	3.1808	- '0071		+ 6 47 8.21	+ .40	13.038	• 346	054	22:24	7'39:7'44	2354
460	3.9	Veloruma		+ .011	2.0328	+ .0023		-45 40 32.44	+ .09	13.075		013	34:35	7.26 : 7.25	2358
461	5.3	14 Hydræ	8 44 20.212	+ .012	+ 3.0169	0036	0014	- 3 4 19.23	+ .22	-13.198	326	024	33:42	8.23:9.05	2365
462	5.7	Chamœleontis	44 43.606	+ '120	- 1.9313	- '2217	0151	-78 36 1.30	26	13.167	1	+ .033	46	7.92	2366
463	8.8*	Gilliss P.Z. 6020	45 26.22		-27.819	-11.600		-88 8 24.43		13.246	+3.050		II	11.38	
464	4.2	Pyxidisγ	46 17.188	+ .073	+ 2.5451	+ .0022	0	-27 20 19.70	- *57	13.351	— ·27 I	+ .081	31	7.07	2375
465	6.2	55 Caneri	46 38 224	+ '290	3. 5828	0196	0365	+28 42 44.45	+ 1.92	13.221	•381	- '245	21	7.95	2380
466	6.8	Lacaille 3577	8 48 13.378	+ .008	+ 2.2203	+ .0034	0012	-40 36 37.37	+ .05	-13.435	234	007	30:31	6.96 : 6.94	2386
467	3.5	16 Hydræ	50 6.440	+ .020	3.1746	0070		+ 6 19 34.13	07	13.241	• 335	+ .000	28:33	7.18:7.66	2393
468	5.8	60 Cancri	5	+ .002	3. 2813	- '0096		+12 0 29.33	+ .12	13.292	• 346	021	18	8.11	2394
469	5.4	Lacaille 3596 Carinæc	50 29.273	+ .012 + .023	2.0110	+ .0025	0019	-47 8 24.76	+ .30	13.017	•210	042	30:31	7'24:7'19	2395
470	3.9	a design of the second s		+ .053	1.3634	0011	0034	-60 15 44.88	- '39	13.662	.139	+ .022	24:26	6.83 : 6.81	2406
471	4.4	65 Cancriα	8 53 1.120	- '017	+ 3.2861	0098		+12 14 41.44	+ .27	-13.226	343	039	27:28	6.93 : 6.81	2407
472	5.3	Carinæb <sup>1</sup> 69 Cancri		+ .013	1.4708	0024	0019	-58 50 35.41	- *02	13.831	•149	+ .005	18	8.35	2414
473	5.6	Velorumc	8 56 53°557 9 0 42°268	·000 + ·047	3.5158	0172 + .0035		+24 50 47 39 -46 41 58 34	+ .04 + .16	13.987	· 361 · 206	- ·005	31 : 33 30	7'97:7'96 6'87	2426 2438
474	5.5	18 Hydræ		+ .011	3.1612			+ 5 29 31.09	+ '02	14 243	.319	003	19:20	7'79:7'77	2439
476	4·1	Volantisa		+ .016	+ 0.9559	0223	0023						05 . 05	6.84 : 7.00	1 3
470	5.3	76 Cancri		+ .011	3.2241	- '0094		-65 59 49.21	+ '71 + '10	-14°331 14°330	- ·092 ·326	- ·102	25:27 24:25	8.76 : 8.86	2440 2445
478	5.3	77 Cancriξ	3 36.635	002	3.4569	- '0159	-	+22 27 0.16	+ .06	14.404	345	007	20	8.03	2449
479	1.8	Argûsλ		+ .012	2.2046	+ .0045		-43 1 43.46	04	14.435	.217	+ .005	31:32	7'00:7'06	2452
480	4.4	CarinæG	4 53.043	+ .038	0.1868	- '0625	0022	-72 12 1'01	+ .05	14.482	°012	008	25:26	6.87 : 6.82	2458
481	3.4	Carinæa	9 8 20.001	+ .043	+ 1.5788	0030	0052	-58 33 25.94	+ .04	-14.686	- 150	005	18	8.36	2473
482	4.0	22 Hydræθ	9 9.808	073	+ 3.1244	•0060	+.0087	+ 2 44 7.67	+ 2.61	15.042	304	312	19:18	8.41 : 8.35	2479
483	5.2	Octantis		+1.12	- 7.860			-85 15 46.67			+ .786	+ .033	66 : 189	11.39 : 8.82	2486
484	1.2	ArgûsB	12 5'974		+ 0.6720	.0328		-69 18 18.10	80	14.806		+ .098	21	8.21	2493
485	6.9	83 Caneri			3.3221	.0132	- 18001	+18 7 44.84	+ .92	15.112		138	24:28	6.88 : 6.87	2501
486	2'0	Argûs		+ .024	+ 1.6062	0023	0034	-58 51 20.19	+ .01	-15.040	- '148	002	21:22	7-15:7-13	2503
487	5.4	VelorumK 40 Lyncis	14 46.173		1·9934 3·6676	+ .0040		-50 37 48.86	- '04	15.024	• 185		21:22	7.21: 2.22	2504
488 489	3·3 5·0	20 Lyneisθ		+ .011	3 0070	0265 + .0035		+34 48 56.10	- ·08 + ·06	15.060	·344 ·246	000	21 33:36	7°54 7°13 : 7°18	2507 2516
409	2.4	Argûs	19 0.991		1.8558	+ .0026		-25 $32$ $23$ $45-54$ $35$ 0.53	+ .03	15.306	• 168	005	29:30	6.93 : 6.88	2526
-							10.00								
491 492	6.0 2.0	28 Hydræα	9 20 24 043 22 40 418	1 -	+ 3.0007 2.9488	- '0027	0011	- 4 41 10.43 - 8 13 30.32	+ .08	-15·389 15·475	- ·274 ·266	010 + .031	26:28 66:78	7'99 : 8'02 7'62 : 7'79	2529
492	4.6	Antliæ	25 7.030		2 9435	+ .0029		-35 30 50.06	+ 14	15.4/5	200	- ·019	33:35	7.18:7.15	2533 2544
494	5.2	5 Leonisξ	26 33.355		3.2383	0100		+11 44 33.02	+ .77	15.806	•286	087	27:33	8.60 : 8.81	2555
495†	3.2	Argûs $m$ . $\psi$	26 45.568		2.3599	+ .0062		-40 1 43.36	46	15.664	. 205	+ .000	24	6.95	2558
496	2.8	VelorumN	9 28 10 929	+ .039	+ 1.8213	+ .0027	0048	-56 35 35.36	03	-15-803	156	+ .004	21	8.14	2567
497	5.2	Lacaille 3900	28 21.054		2.3783	•0068	+.0001	-40 12 24.70	+ '14	15.835	•206	019	23:25	7.24 : 7.17	2568
498	5.3	Lalande 18817		+ .014		+ .0028	0019		+ .01	15.832	•240	002	20	7.42	2569
499	5.9	33 HydræA	29 33.271	•000	2.9943	0023	.0000	- 5 28 7.60	+ .47	15.937	•260	057	20	8.19	2572
500	5.6	CarinæH	30 51.356	+ .020	0.4265	0228	0068	-72 38 14.37	+ .02	15.960	•034	010	24	7*30 : 7*26	2579
				1.00				•							

459. 3'9, 4'4. Very close binary; C not seen. 495. 3'7, 5'7 0''.5 339° 1902'2.

.

No.	Mag.	Name.	Mean R. A. 1900°0.	$\mu_{a}\Delta E.$	Annual Variation 1900 °O.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900°0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
			hm s	s	8	s	S	<u> </u>	- ".08				1		
501	4'1	Carinæh 1 Sextantis	9 31 32.565	+ .013	+ 1.7398		0018	-58 47 1.41		-15.975		+ "011	22	7°20	2581
502	5°4 4°9	2 Sextantis	31 55°911 33 14°265	+ '036	3.1326	- ·0077 - ·0066	0044	+7172.82 +562.93	+ .06 + .52	16.013	·271 ·265	- ·007 - ·063	21 16:18	8.08 8.37 : 8.23	2582
503 504	49	Velorum	33 14 205					-48 54 24.09	+ 5 <sup>2</sup> - · 17	16.023		+ .023	24		2589
505	5.6	Velorumy	34 7.066		1	+ '0075	-		+ .32	16.162	.196	044	22	7 23	2590 2594
															-394
506	4'1	35 Hydræ	9 34 44 980	1	+ 3.0629		+.0031	- 0 41 20.39	+ .60	-16.55	- *258	072	15	8.32	2595
507	5.1	38 Hydræ	35 30.708	+ .014	2.8759			-13 52 43'26		16.209	*240	- *016	20:22	7.97	2600
508	3.7	14 Leonisο Chamæleontisζ	35 48·807 36 50	+ .062	+ 3.2061	- '0092	0098	+10 20 50'07	-	16·248 16·254	- '267	039	29:32	6.82 : 6.87	2602
509 510	5°3 5°8	16 Leonis		+ .002	+ 3'2722		1	-80 29 30.06 +14 28 45.42	02	16.349	+ 146	+ .007	3.	7.28	2606 2612
510	50	το μεσιμό				- 0115			T 11		- 2/0	- 014	27:29	007.009	2012
511	2.1	Autliæθ	9 39 44 589	+ '031	+ 2.6714	+ .0022		-27 18 41.82	19	-16.382	519	+ .023	27	6.96	2615
512	3.1	17 Leonis	40 10.549	+ .022	3.4140	- '0179	4	+24 14 5.13	1	16.424	•278	024	24:27	7.37:7.44	2618
513	var.	Carinæl	42 29.972	+ .024	1.6472	- *0002	0035	-62 2 47 67	19	16.525	.128	+ .051	21:22	7.52:7.59	2628
514	6.0	Lacaille 4022		+ .002	2.3357	+ .0086		-44 17 33.55	.00	16.221	•185	•000	33:35	7'25:7'29	2629
515	6.9	23 Leonis	45 37 358	- '014	3.2220	0109	+.0012	+13 32 1.79	+ '22	16.725	*255	022	23	8.02	2639
516	6.2	6 Sextantis	9 46 11.709	002	+ 3.0246	0025	+.0000	- 3 46 29.32	+ '22	-16.756	- '236	030	32:31	7'34:7'24	2641
517	4.1	24 Leonisµ	47 4.544	+ .119	3.4208	0196		+26 28 40.55	+ .45	16.832	.265	063	19	7.12	2648
518	4.5	Velorumm		+ .020	2.3129	+ .0003	- '0027	-46 4 42.78	+ .20	16.831	.177	027	25:26	7 * 54 : 7 * 47	2651
519	6.2	Bradley 1393		+ .048	3.1843	- '0085		+ 9 24 25.62	06	16.953	• 240	+ .002	32:33	7.94	2663
520	5.2	27 Leonis	52 50.648	+ .012	3.2318	0102	- '0021	+12 55 18.67	+ .23	17.069	•241	- '029	23	8.00	2672
521	3.2	Argûs	9 53 21.066	+ .012	+ 2.1010	+ .0004	0025	-54 5 29.97	+ .04	-17.060	153	000	23:24	6.86 : 6.83	2674
522	6.7	Lacaille 4092!		+ .018	2.2954	+ .0103	-'0024	-47 56 13.16	+ .23	17.118	•168	031	19:21	7.62:7.52	2676
523	5.3	Antliæη	54 34.746	+ .060	2.5703	+ .0085	0076	-35 24 44.43	+ '21	17.146	.187	- '027	19:20	7.84:7.82	2679
524	5.0	29 Leonisπ		+ .018	3.1740	0080	- '0023	+ 8 31 26.36	+ .21	17.162	•233	027	43:45	8.00 : 7.91	2680
525	6.1	Lacaille 4126	9 59 43 739	+ .072	2.7668	+ .0052	0105	-23 48 5.38	- '16	17:327	•194	+ '022	34:36	7'02:7'05	2688
526	4.8	40 Hydræ	10 0 15.271	+ .020	+ 2.9210	+ .0012	- '0025	-12 34 47.04	09	-17:360	- '205	+ .015	24	7'90	2690
527	3.2	30 Leonisη	I 52.886	+ .001	3.2768	0120	-	+17 15 1.32	+ .09	17.455	*228	- '012	33:35	7'14:7'16	2694
528	1.2	32 Leonisa			+ 3.1996	- '0100		+12 27 21.79	+ '02	17.496	219	003	33:33	7.20	2698
529	5.7	Chamæleontisµ	3 24		- 1.4240	3466	-	-81 43 50.17	22	17.478	-	+ .030	13	7.32	2699
530	5.1	VelorumQ	5 8.726		+ 2.2693	+ .0123	0010	-51 19 14.31	+ .04	17.587	- 151	005	27:29	7.21 : 7.23	2702
1 521	3.8	er Undres	1						1 .60	1.					
531	7.6	41 Hydræλ Lacaille 4342		+ .104	+ 2.9244	+ .0014	- '0137	-11 51 35.94 -86 25 32.30	+ .69	-17.699	- 195	- °093 + °002	44:43	7.59:7.46	2706
532 533	3.9	Velorumq		,		- 2.338		-41 37 34.77	- '02 - '24	1	+ .485		52 : 64 23		
534	3.4	36 Leonis	11 7.781	014	3.3420			+23 54 56.50		17 //3	- 100	- '015	16:17	7.90	2723 2730
535	3.4	Argûs		+ .047	1.4318			-69 32 28.67	$+ \cdot 13$ $+ \cdot 02$	17.838	•087	- '002	16:17	8.87:8.83	2733
10.00						_	1.								-133
536	5'5	22 Sextantis.		+ .089	+ 2.9814	.0000	0108	- 7 34 10.12	- '02	-17.886	188	+ '002	17:19	8.20 : 8.21	2735
537	3.3	Carinæq		+ .052	1.9954	+ .0114	0001	-60 49 57 33	+ .06	17.937	•122	002	21	8.49	2739
538	5.7	Lacaille 4260 42 Leonis		006	2.4418			-47 11 46.67	+ .13	18.045	•148	- '019	27:30	7.09:7.06	2749
539 540	6.4	Velorumr			3.2312	0112			+ '24	18.066	·198 - ·153	- '030	20	8.10	2752
540	4'9			+ '021	T 2.2009	+ .0129	-*0029	-41 8 48.03	37	18.042	- 153	T 055	27:29	7'11:7'04	2758
541	7.5	C. G. A. 14444			-29.872	- 31.819		-89 0 24.06			+1.869		39:49	11.42 : 10.62	
542	5.6	Lacaille 4278		+ .000			0122	-37 30 8.65		18.199	154	063	28	7.07	2763
543	4.0	42 Hydræµ	21 15.160	+ .068	2.8999	+ .0040	0089	-16 19 33.46	+ .66	18.299	•168	084	58:67	7.69:7.85	2771
544	4.0	CarinæI	22 24 637	+ '032	1.2012	0224	0043	-73 31 21.96	+ .19	18.278	•064	- *021	21	7.20	2778
545	4.4	Antliæa	22 34.483	+ .039	2.7416	+ .0097	0023	-30 33 30.63	.00	18.263	•157	.000	20:24	7.27 : 7.22	2779
546	4.0	Carinæ	10 24 12.394	+ '014	+ 2.1944	+ .0163	0020	-58 13 43.43	+ .00	-18:329	- •122	- •008	33	7.12	2784
547	5.3	29 Sextantis	24 23 974	+ .026	3.0481	0019	0035	- 2 13 38.35	+ .12	18.347	• 172	019	20:21	8.03	2788
548	3.8	47 Leonis		+ .004	3.1627	- '0079			+ .04	18.444	.173	000	39:41	7 .03 : 6.96	2804
549	3.4	Carinæp	28 27.995	+ '024	+ 2.1247	+ .0168		-61 10 15.03	05	18.462	112	+ .001	25:26	6.97 : 7.05	2811
550	7.8	C. G. A. 14481	29 6.70		- 4.400	- 1.240		-86 2 52.31		18.491	+ .256		46 : 62	11'42 : 10'45	
	-									-					

513. L, 3.6-5.0; P, 35d.5

.

1

FROM OBSERVATIONS IN THE YEARS 1905-11.

1	1				1		1				1	1			
			Mean R.A.	321	Annual	Sec.		Mean Dec.		Annual	Sec.			Epoch	
No.	Mag.	Name.		$\mu_{\alpha}\Delta E.$	Variation	Var.	Proper Motion.	111111111	$\mu_{\delta}\Delta E.$	Variation	Var.	Proper Motion.	No. of Obs.		Boss No.
1		Second and the	1900°0.		1900'0.	1900'0.		1900°0.		1900.0	1900'0.			1900+.	
							1				1				
			h m s	S	8	s	8	0 / 11						1 3 3	
551	5.2	44 Hydræ				+ .0075	0002	-23 13 47.28	00	-18.488	- "152		23	7.28	2815
552	5.4	48 Leonis		+ .059	3.1325	- •0065		+ 7 28 7.26	42	18.455	.167	+ .025	25	8.16	2816
553	4.9	37 Leonis Minoris Velorumm. p	33 5.628 33 5.788	- °004 + °116	3.3880		1	+32 29 44.87	+ '02	18.626	·175 ·127	- ·003	19	7.81	2829 2830
555	4'0	Chamæleontis		+ .110	0'7440	- '0693	0136	-78 5 20.94	- 14	18.645		+ .010	29 34	7.15	2837
556	4°3 6·8	Velorumx				+ .0196	0023	-55 4 56.54	+ .12	-18.713	117	019	19:20	8.07:7.96	2842
557 558	7.1	33 Sextantis Lacaille 4510		+ .02	+ 3.0528 - 3.023	- ·0019 - 1·078	-·0094	- 1 12 58.47 -85 34 20.89	+ 1.04	18.854	- ·150 + ·165	- ·129	16 54 : 156	8.08 11.44 : 8.76	2846 2849
559	6.9	34 Sextantis	37 27.636		+ 3.1000	- '0045		+ 4 6 20.19	- '19	18.740		+ .021	17:19	9.03:8.99	2851
560	5.1	41 Leonis Minoris	37 58.730	+ .065	3.2697	0164		+23 42 43.03	- '04	18.772	.159	+ .005	17	7.84	2854
561	2.8	A norîte A		+ .023	+ 2.1303	+ .0202			06	-18.811		+ .009	18:19		2862
562	5.4	Argûsθ 42 Leonis Minorisθ		+ '014	3.3463	0202	0033	-63 52 13.57 +31 12 32.65		18.888	- °099 °158	- '041	21	7'10:7'08 6'81	2866
563	6.6	37 Sextantis	40 53.296	+ .004	3.1266	0028		+ 6 54 0.41	+ '30	18.902	• 146	038	17	7.96	2868
564	var.	Argûsη	41 10.786	004	2.3181			-59 9 31.45	- '02	18.871	•106	+ '002	12	8.86	2871
565†	2.6	Argûspr. µ	42 28.123	032	2.5692	+ '0196	+.0022	-48 53 30.53	+ '41	18.970	•116	059	27:28	7'07:7'00	2875
566	5.5	53 Leonis	10 44 0.000	+ .001	+ 3.1574	- '0080	0001	+11 4 27.53	+ .23	-18.988	142	033	25	7.11	2883
567	3.3	Hydræv	44 41.476	048	2.9578			-15 40 11.84		18.781		+ .193	27:28	7.44 : 7.15	2888
568	4.6	Chamæleontis	44 50.91	+ .21	+ 0.6097		0183	-80 0 45.85	+ .04	18.983	020	004	17:19	11.46 : 11.03	2889
569	7.6	Lacaille 4578	45 54.55		- 3.625	- 1.200	<b></b>	-86 22 21.91		19.008	+ .176		34:51	11.41 : 10.08	
570	3.9	46 Leonis Minoris	47 43.279	024	+ 3.3674	0258	+.0074	+34 45 12.74	+ 2.07	19:348	144	- '290	32:35	7.29:7.13	2899
571	3.8	Carinæu	10 49 25.854	060	+ 2.4237	+ .0251	+.0078	-58 19 19.18	17	-19.085	099	+ .022	30	7.75	2908
572	4.7	Antliæ	52 3.449	054				-36 36 1.30		19.311	.111	139	47:49	7'39:7'29	2919
573	4.2	7 Craterisa	54 53.852	+ .265	2.9197	+ .0067	0326	-17 45 57.60	98	19.123	.110	+ .121	24	8.13	2925
574	5.1	58 Leonis d	55 23.805	006	3.0999	0032		+ 4 9 15.69	+ .20	19.277	•117	- '021	45:54	9*25 : 9*48	2927
575	4.6	Velorumi	55 33.905	- '015	2.7442	+ .0186	+.0051	-41 41 21.93	+ .02	19.267	. 103	002	27:28	7'28:7'26	2929
576	4.7	63 Leonis	10 59 51.373	+ .192	+ 3.0920	0055	0233	+ 7 52 35.49	+ .38	-19.407	108	047	19	8.25 : 8.11	2942
577	6.4	Octantis	11 0 ò·86	+ .48	- 0'291	330	042	-84 3 21.48	+ .12	19:378	+ .021	014	52:155	11.49 : 8.77	2944
578	5.2	Hydræx		-		+ .0112		-26 45 13.69		19.388	099	013	15	8.20	2947
579 <sup>†</sup> 580	5.8	65 Leonispr. p <sup>4</sup>	1 47 970	+ .208	3.0019	- '0026	_	+ 2 29 53.26	+ .72	19.491	.103	087	22:23	8.27 : 8.25	2950
1.28	5.9	Lacaille 4625		+ .042	2.1488	+ .0314	0064	-70 20 13.01	+ .10	19.449	*068	014	28:31	7'31:7'20	2955
581	3.9	Carinæx			+ 2.5466	+ .0301		-58 25 59.91	+ '02	-19.460	081	002	28:29	7.61 : 7.60	2960
582	4.6	11 Crateris			+ 2.9463					19.609	001	101	36:34	7'10:6'72	2964
583 584	7°2 5°8	Lacaille 4708			- 0.241		1	-85 12 25.13	1		+ .026		44 : 69	11.48 : 9.96	
585	5.6	Lacaille 4649 Lacaille 4657		+ .032	+ 2°7198 2°4690			-48 33 28.71 -63 37 33.37	- ·20 + ·05	19.209	- ·081 ·072	+ '024	17 17:18	8·23 8·56	2965 2969
		and the second second	and the second se		3102-51	11111111	-					1910	= 01		2909
586 587	2.5	68 Leonisδ			+ 3.1970			+21 4 16.67		-19.693	095	- '145	16	8.70	2972
588	3.3	70 Leonisθ Lacaille 4731θ	8 59·569 10 29·42		+ 3.1525 - 0.727	- °0098 - °569				19.638	093	086	16:17	8.57 : 8.46	2974
589	5.7	73 Leonisn		+ .002	-0.727 + 3.1422			-85 41 14.92 +13 51 11.04	+ '22	19.281 19.610	+ ·031	- '026	37:57	11.51 : 10°01 8°46 : 8°44	 2978
590	4.6	74 Leonis		+ .064	3.0496		1	- 3 6 18.03	1	19.645	- 090 ·085	044	16	8.57	2970
591†	1.20						1								-
591	3.5	54 Ursæ Majolispr. ν 12 Craterisδ	11 13 4°695 14 20°361	+ '014	+ 3°2513 2°9967	- ·0225 + ·0065		+33 38 24.52	12	-19.613	- ·088 ·078	+ .012	18	7'99:7'81 8'79:8'66	2985
593	4.2	77 Leonisσ		+ .021	3.0925			+ 6 34 38.28	- 1.69 + .12	19.455 19.693	•078	012	16 : 17 22	8.17:8.02	2989 2990
594	4.2	Centauri		+ .031	2.7216			-53 56 35.04		19 093	·066	018	20:21	7.76:7.70	2990
595†	4.0	78 Leonis	18 42.829	077	3.1297			+11 4 47 99	+ .62	19.807	.074	085	29	7.30	2999
596	6.7	Lacaille 47 36	11 19 34 930	+ .012	+ 2.8582			12	+ .14	-19.754	065	018	21	7.60	3003
597†	4.2	15 Craterispr. γ	19 53.050					-42 7 11 51 -17 8 5 41		19.744	- 005	003	17:18	8.17	3003
598	5.5	Lacaille 4739	20 38.410	-	1			-35 30 50.03		19 744		+ .007	16	7.89	3010
599	6.7	83 Leonis	21 41 236		3.0380	0020	0482	+ 3 33 30.20	- 1.49	19.594		+ .173	18:19	8.60	3014
600	5.3	84 Leonis $\tau$	22 47 704	011	3.0862	- '0020	+.0013	+ 3 24 24.77	+ .19	19.802	.064	019	23:22	8.62 : 8.46	3021
				1						1	1	1			
		554. 4.5, 5.0	0"'7 261°	1897'1.				FOI 7	.5, 10'0	7".5	146° 1	905'2.			×
	-	564. L, > $I-7.4$ ;	irregular.					595-4	0, 7'2	2".6; bir	ary. No	note of d	uplicity.		
1			2"*2 61° 2"*3 86°	1900°4. 1902°6,			A STATE		°2, 10°5	5"*5	94° 1	898.2.			
and the second			-												

.

															_
No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_{\alpha}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900*0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900°0.	Sec. Var. 1900°0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
									1		1				
60I	8.4*	Gilliss P.Z. 7980	hm s	S	a 	8 -21.838	3	-89° 14 56.25	"	-19.784	+ ": 324	"	25 2	11.52 : 9.87	
602	7.6	C. G. A. 15761			- 6.162	- 6.471		-88 41 35.47		19.796	+ .324 + .150		25:42 32:47	11 52 : 9 87	
603	5.1	87 Leonis	25 12.326	011	+ 3.0620	+ '0012	+.0013	- 2 27 6.61	+ '14	19'833	- '059	012	22:24	8.53:8.47	3029
604	3.6	Hydræξ		+ .112	2.9436	.0166		-31 18 15.66	+ .37	19.904	·051	- '051	27	7.35	3042
605	5.2	CentauriC <sup>2</sup>	31 4.670	- '019	2.8940	.0285	+.0026	-47 5 14.01	+ .40	19.942	°045	055	22	7.20	3053
	Con the			1					1.000						
606 607	3.1	Centauriλ 91 Leonisυ		+ ·050 •000	+ 2.7447	+ .0451	-•0060 •0000	-62 27 59.41 - 0 16 17.74	+ .18	-19.910	- '042	- '022	19:21	8.34 : 8.22	3054
608	4°5 5°8	Chamæleontis $\pi$	31 49°727 33 7°808	+ .183	3.0716	·0004 + ·0680	0250	-75 20 34.47	- ·27 + ·18	19.861	•046	+ .035	27:25	7'95:7'81	3058 3064
609	7.5	Lacaille 4865		- 103	I'443	- '019		-84 55 57.95	T 10	19.933 19.929	·033		39:62	7 '33 11 '48 : 10 '12	
610*	4.9	Hydræo		+ .018	2.9721	+ '0194	0024	-34 11 25.52	01	19 929	.038		39.02	7'48:7'46	3073
	1.1														
611	5.0	27 Crateris		- '020	+ 3.0368	+ .0101	+.0024	-17 47 41.57	+ .32	-20.002	031	039	16:17	8.30 : 8.31	3087
612	4'3	3 Virginis		+ .011	3.0821	- '0030	- '0012	+ 7 5 21.43	+ 1.68	20.163	·029	- • 187	16:17	9.04:9.00	3089
613	5.2	Lacaille 4878		+ .045	2°9535 2°8041	+ .0284	0064	-45 8 5.51	.00	19.976	·028	*000	23	7.03	3091
614 615	3.7	Muscæλ Lacaille 4885		+ .131	2.8041	•0566		66 10 27.45 60 37 20.70	$- \cdot 22 + \cdot 26$	19.949		+ .028	21:23	8'12:7'92	3092
015	4°I			+ .035	2.0/91	.0471	0036	-00 37 20 70	+ '26	20'012	.022	029	16	8.92	3094
616	5.6	Lacaille 4898	11 43 41 967	+ .000	+ 3.0249	+ .0122	0007	-26 11 37.54	+ '21	-20'022	023	026	24:25	7.95:8.00	3100
617	2°2	94 Leonisβ		+ • 289	3.0634	0071		+15 7 50.72	+ 1.00	20.151	•022	- 123	28:31	8.46 : 8.65	3101
618	3.6	5 Virginisß	45 29.592	- '402	3.1223	- '0002		+ 2 19 39.52	+ 2.25	20.286	·02I	279	26:25	8.13 : 8.02	3105
619	4.2	CentauriB		+ .062	2.9835	+ .0288	0095	-44 37 1.49	•00	20'010	.018	.000	27:29	7.26:7.23	3109
620	5.2	95 Leonis	50 31.995	014	3.0001	- '0074	+.0012	+16 12 11.55	+ .00	20.032	.010.	002	30:31	8.31	3123
621	5.7	Lacaille 4959	11 53 11.800	+ .011	+ 3.0126	+ .0436	0012	-55 45 37 94	+ .19	-20.064	005	026	33	7.42	3129
622	6.3	Lacaille 4966		+ .018	3.0271	+ .0375	0026	-51 8 23.07	+ •11	20.022	.003	012	20:21	7.04 : 7.00	3133
623	5.5	7 Virginisb		+ .010	3.0732	0006	0011	+ 4 12 43.28	+ .19	20.060	002	- ·018	18:19	8.85	3135
624	4.7	8 Virginisπ	55 44 935	+ .003	3.0752	0022	0003	+ 7 10 18.62	+ '29	20.076	.000	033	20	8.98:8.83	3139
625	6.0	Lacaille 4991	57 18.62	+ . 59	2.839	+ .296	021	-85 4 29.63	+ .01	20.042	+ .004	001	47:74	11.25 : 10.10	3144
626	4'4	Crucis $\theta^1$	11 57 55.676	+ .122	+ 3.0277	+ .0281	0212	-62 45 22.27	+ .06	-20.024	+ .004	008	23:25	7'32:7'16	3146
627	5.4	Lacaille 4992		235	3.0930		+.0286		+ 1.03	20.171	.006	- '125	21:22	8.20 : 8.25	3148
628	4'3	9 Virginis		+ .100	3.0222	0030		+ 9 17 18.65	- '29	20.000		+ .038	40:46	7'39:7'52	3155
629	5.7	CentauriE		+ .021		+ .0357		-48 8 8.30	+ '24	20.079	.012	034	25	7.10	3163
630	2.7	Centauriδ		+ .032	3.0904	.0382	0041	-50 9 55.70	+ .12	20.062	.012	012	15	8.61	3165
631	6.1	Lacaille 5036	10 2 42:002	+ .021	+ 3.0872	+ .0310	0029	-43 46 5.74	+ .56	-20'111	+ .016	067	17:20	8.64 : 8.41	3167
632	6.3	Io Virginis		- '028	3.0742			+22731.90		20.222		184	17.20	9.39	3169
633	3.1	2 Corvi			3.0794					20.032		+ .001		7.45:7.65	3172
634	6.9	Lacaille 5096		+ .84	4.486	1.234	072	-87 51 33.57	.00	20.030	·035`	.000	21:42	11.55 : 9.22	3185
635	2.9	Crucisδ		+ '042	3.1293	.0532	0055	-58 11 33.25		20.047	·028	010	19:21	7'69:7'72	3187
								State of the second		-	1	1.1011		the second se	1.0
636	2.6	4 Corviγ Muscæε		+ .099	+ 3.0801	+ '0116		-16 59 12.06 -67 24 15.60	- 'IO	-20°014 20°063		+ .011	20 : 19 18 : 20	8·84 : 8·81 8·18 : 7·91	3191
637 638	4°I	Chamæleontisβ	12 9°522 12 28°392	+ '331	3-2022	1865		-78 45 25.18	+ .35	20.003	•033 •036	- ·044 + ·012	18:20 35:37	8.95:8.84	3197 3199
639	4'3 4'0	15 Virginisη	12 28 392 14 47 339		3.0684	-	1 - 1		+ .19	20.003	·030	025	33:37	7.88:7.64	3210
640	5°I	16 Virginis	14 47 339		3.0465	*00028		+3529.23		20.030		- ·078	33 · 3 <sup>2</sup> 21 : 23	8.64 : 8.61	3213
		a substantia da la secola de la s								1 1 1 1 1 1					
641	3.4	Crucia'e		+ .189	+ 3°2081	+ .0282	0243	-59 50 54.39	60	- 19.920		+ .078	21:23	7.77:7.68	3218
642	4'9	12 Comæ	17 28.725		3.0212				+ •12	20.003	·042	014	20:22	8.52 : 8.50	3224
643	6.7	Lacaille 5107		+ .19	4.388	+ '720	019	0 00 10 0	+ .06	19.994	•057 •048	- ·006 - ·04*	45:69	11°59 : 10°10 6°57	3225
644	6·3 6·0	Lacaille 5141 Centauri $m. x^2$	19 50.416 20 5.478		3°1762 3°1499		-•013* -•0033	-41 57 34.48	+ °26 + °10	19°972 19°983	·048 ·049	- '04* - '013	19 17:19	7.72:7.69	 3232
645†		001100111	20 3 4/0		and the state	1- 0240		-34 37 55.77	10					1	
646	5.2	14 Comæ			+ 3.0049	- '0119			+ .16	- 19.978	+ .049	018	16	8.96	3240
647	5.8	Lacaille 5154		+ .008		+ '0228			+ .35	19.992	°052	039	16:17	8.94 : 8.88	3241
648	4.6	15 Comæ		+ .062	2.9954				+ .82	20.042	•050	- *087	16	9'44	3242
649	4'I	Centauri		+ .026	3.2238			-49 40 36.54	1 1	19.977	·055	- *028		6.81 : 6.76 9.08 : 9.04	3245
650	6.3	Mayer 525	22 43.628	+ .049	3.0263	0053	- 0054	- 4 3 43.34	+ .08	19.957	·053	- •009	16 : 18	9 00 : 9 04	3247
			,	-											

610. Greek letter not in Auwers' Bradley. 645. 6'7, 6'9 0'''2 41° 1897'5.

FROM OBSERVATIONS IN THE YEARS 1905-11.

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_a \Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0,	Proper Motion.	Меан Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
651 652	3.0	7 Corvi	h m s 12 24 41 221 24 41 857	8 + •110 - •022	+ 3.0990 3.0186	s + ·0119 - ·0080	+.0027	-15 57 32.67 +21 26 59.36	+ 1 <sup>"</sup> 10 + 36	- 20 <sup>°</sup> 074 19 <sup>•</sup> 976	+ "057 .056	- "143 - ·046	27 : 29 16 : 20	7*66 : 7*70 8*21 : 7*93	3256 3257
6 <sub>53</sub> 654	1.3 4.0	Crucis γ Muscæγ	26 29.339	- •018 + •074	3·3003 3·5260	+ ·0549 ·1188	0092	-71 34 49.85	+ 2.12	20°194 19°926	•062 •068	- ·272 - ·013	17 23:25	7.92 8.00 : 7.96	3263 3269
655 656	2·8 4·9	9 Corvi	29 7·987 12 29 52·098	·000	3.1437	+ '0165 - '0085		-22 50 37 ·91 +23 10 47 ·78	+ ·47 - ·06	19°946 — 19°871	·067 + ·065	- °061 + °007	39:36 17	8·20 : 7·76 8·48	3280 3283
657 658	5.3	24 Comæseq. Muscæa	30 6.833 31 12.950		3°0127 3°5285	- ·0062 + ·1013		+18 55 39 <sup>23</sup> -68 35 4 <sup>35</sup>	- ·12 + ·20	19·858 19·882	·066 ·078	+ ·016 - ·021	21 : 22 16 : 17	7 <b>·6</b> 9 : 7 <b>·</b> 62 9·62	3285 3289
659 660	6·1 4·0	25 Virginisf Centauriτ	31 38·263 32 13·777		3°0873 3°2606	·0064 ·0404	-·0020 -·0197	- 5 16 51.09 -47 59 26.73	+ .23 + .12	19·883 19·867	·070 ·075	- ·027 - ·018	16 22	8·59 6·84	3290 3292
661 662†	4·8 2·1	26 Virginisχ Centauri	12 34 5.020	+ .042	+ 3.0931	+ .0077	- · 0051	- 7 26 43.07 -48 24 38.07	+ .31	- 19.863 19.815	+ .075	- ·037 - ·015	21 27:28	8°29 7°04 : 6'96	3298 3302
663 664	5°0 5°8	30 Virginis	36 49.442 38 40.595	053	3.0377		+.0001	+10 47 11.12 -27 46 30.77	+ .86	19.890	•079 •086	- '101 - '054	22 25 : 27	8.67 : 8.53 7.09 : 7.06	3309 3318
665†	3.1	Museæ	40 8.616	+ .039	3'6312	+ .1010	0021	-67 33 38.43	+ .55	19.769	. 101	029	18	7.58	3320
666 667	5°4 7°1	32 V.irginisd <sup>2</sup> Lacaille 5235	12 40 33 849 40 58 76	+ ·066 + ·86	+ 3.0308	+28.711	072	+ 8 13 12.40 -89 15 0.86		19.738	+ .086	011	17 38 : 53	8.64 11.61 : 9.99	3323 3325
668 669	1.1 1.1	Crucis		+ .021	3.4726 3.0542	+ ·0660 ·0022	0003	-59 8 31.60 + 4 7 7.14	+ .10	19'741 19'711	.100 .100	- ·028 - ·012	19 17:18	8.17 8.61	3328 3331
670 671	5.2	Octantist Centaurip	44 27 <sup>.8</sup> 4 12 45 15 <sup>.498</sup>	- ·51	5 <sup>.8</sup> 31 + 3 <sup>.2426</sup>	·866	+.044	-84 34 48·46	- ·09 + ·30	19.661	· 173 + · 101	036	54 : 167 17 : 18	11°58 : 8°80 8°38 : 8°23	3340 3342
672 673	5°1 4°3	31 Comæn		+ .011	2·9264 3·3078	- ·0096 + ·0323	0015	+28 5 5.11 -39 38 6.18	+ ·24 + ·26	19.656	·095	- ·026 - ·038	16 21:23	9·11 6·90	3347 3352
674 675	5.0	40 Virginisψ 43 Virginisδ		+ .013	3.1128	·0093	0016	-85945.43 + 35626.50	+ .17 + .48	19.609 19.625	· 105	- '021 - '064	21 46	8.04 7.71 : 7.51	3362 3367
676	3.2	Muscæδ	12 55 23.669	- • 381	+ 4.0224	+ .1427	+.0536	-71 0 34.67	+ .21	- 19.494	+ .125	030	29	7.10	3377
677 678	2·8 7·1	47 Virginis		+ 144	2.9867 9.435	- '0006 + 2.862		+11 29 47 95 -87 1 22 97	13	19·409 19·423	·115 ·349	+ .012	38:36 57:77	7'79 : 7'66 11'61 : 10'60	
679† 680	6·9 4·4	48 Virginism Centauriξ <sup>2</sup>	12 58 45 195 13 1 4 223	+ ·026 + ·026	3.0878 3.4789	•0066 •0475	-	$\begin{array}{r} -3 & 7 & 31.40 \\ -49 & 22 & 14.19 \end{array}$	+ '34 + '17	19·432 19·364	·122 ·142	- ·040 - ·024	24 28	8.20 7.19	3388 3393
681 682†	6·1 4·4	Lacaille 5398 51 Virginisseq. θ			+ 3.5379	+ .0547	- °0048 - °0026	-52 55 27.56 - 5 0 18.99	+ .24 + .37	-19.357 19.294	+ .145	- ·032 - ·042	26 40 : 37	7°43 8°83 : 8°75	3400 3409
683 684†	5°3 4°7	Lacaille 5422 Lacaille 5418seq.		+ .001	3.4110		0119	-42 50 9.34 -59 23 18.59		19·273 19·254	·148 ·161	- ·043 - ·034	18 17	7·87 8·42	3417 3419
685	4.3	43 Comæ	7 11.977	+ .438	2.8032	0076	- • 0604	+28 23 12.64	- 6.36	18.316		+ .875	22': 23	7'25:7'27	3424
686 687	5.0	Muscæη Centaurir	11 19.783	012	+ 4.0112 3.3201	•0254	- · 0045 + · 0021	-67 21 52.54 -30 58 37.09		-19·178 19·148	+ .180		24 25	7.32	3429 3440
688 689	6.0 5.0	Lacaille 5464σ	12 33.304	+ .008	3°4602 3°0277	·0392 ·0028	0009	+ 5 59 47.96	08	19.112	•163 •146	+ .009	18 16:17	6•98 8·63 : 8·66	3441 3446
690 691	4·8	61 Virginisγ 46 Hydræγ	13 13 29.085		3.1313 + 3.529		-·0754 +·0048	-17 45 28.04 -22 38 39.01	+ 9.72	20·118 	·148 + ·158	-1.084	16 39	8·97 8·60 : 8·59	3448 3449
692 693	2·8 6·2	Centauri Lacaille 5498	16 11.172		3.3584	.0303			+ .71	19.078 18.947	•164 •180	094	23:24 22	7'41:7'53 7'35	3452 3458
694 695	6·7 7·4	Lacaille 5507 Lacaille 5452	17 4.006		3.5622 8.618	.0471 1.580			06	18.916	· 179 · 436	+ .008	23:24 36:59	7'97 : 7'92 11'57 : 10'06	3461
696	0.9	67 Virginisa	13 19 55.418	+ .019	+ 3.1253	+ .0119		-10 38 22.28	+ .25	-18.876	+ .165	036	29	6.91	3476
697 698	8·0 5·7	Lacaille 5444i 68 Virginisi	21 26.045	1. 0	10.091 3.1636	2°356 °0125		-86 12 39.93 -12 11 14.87	+ .19	18.795	·519 ·168		28:40 33:26	11.66 : 10.22 9.00 : 8.52	3481
699 700	5°2 5°7	70 Virginis Octantis		+ .146	2.9341 8.838	1.0001		+14 18 40.99 -85 16 24.77		19.316 18.717	• 160 • 469	1 -	17 : 18 59 : 148	8.76 : 8.75 11.62 : 9.04	3487 3493
					665. 679.	2'9, 2'9; ( 3'7, 4'0 7'6, 7'8 4'4, 8'9	close binan 1''*3 0''*6 6''*8	ry. 341° 1900'4 219° 1899'4 344° 1905'3							
						4.7, 8.5	1".7	349° 1903 3							

.

•

No.	Mag.	Name.	Mean R. A. 1900 °O.	$\mu_a \Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900°0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
7017	3.8	Centaurim. d	hm s 13 25 14.605	s + •010	s + 3.4628	s + '0342	s `0013	-38 53 27.11	+ ":20	-18.701	+ "191	- "025	20	8.01	3496
702	6.1	73 Virginis		+ .055	3.2284		- '0063	-18 12 48.71	+ .20	18.654	.181	023	16:17	8.76	3498
703	5.9	Lacaille 5580		+ .065	3'3294	.0236		-28 10 39.43	+ .19	18.640	.182	- '021	19	7.78	3502
704	3.3	79 Virginis		+ .146	3.0540		-	- 0 5 4.81	- *25	18.200		+ .034	47:45	7.63:7.39	3508
705	6.6	Lacaille 5577	30 38.171	+ .001	5.0096	.2413	0028	-75 10 25.47	+ .58	18.236	• 289	036	21	7.85	3514
706	2.3	Centaurie	13 33 32.963	+ .024	+ 3.7719	+ .0592	0034	-52 57 28.39	+ .19	-18.427	+ .226	027	23:24	7'18:7'14	3521
707	8.7	C. G. A. 18500	35 1.39		13.771	4.327		-87 7 9.08		18.348	·816		28:44	11.67 : 10.25	
708	5.3	82 Virginism	36 21.704	+ .059	3.1440	.0108	0069	- 8 11 54.36	30	18.265	• 194	+ .036	27:22	8.61 : 8.27	3534
709	5.9	83 Virginis	39 6.005	004	3.2300	.0121	+ .0002	-15 40 34.38	+ .04	18.207	. 205	005	17	8.63	3542
710	4.3	I Centaurii	39 59.915	+ .286	3.3959	·0277	0368	-32 32 18.11	+ 1.18	18.321	•215	123	18:19	7.76:7.72	3544
711	4.7	CentauriM	12 40 10.437	002	+ 3.7704	+ .0520	+.0002	-50 55 51.89	+ '31	-18.189	+ '241	032	16	9.22	3547
712	4.6	4 Boötisτ	42 30.310	+ . 295	2.8510	0002	- '0340	+17 57 18.95	22	18.049	.185	+ .026	19:20	8.69:8.54	3558
713	3.5	Centauri	43 30.280	+ .027	3.5798	+ .0380	- '0030			18.062	.235	025	16	9'14	3564
714	3.3	Centauri	43 35.443	+ .018	3.5955	•0392			+ .10	18.052	•236	018	18	8.98	3565
715	5.2	89 Virginis		+ .056	3.2524	.0164			+ .35	18.044	•216	043	35:34	8.12:8.15	3571
	4.8			1 1011	1	1					1			P. of	
716	6.0	4 Centauri	13 47 27°060 48 26°200	+ .011	+ 3.4400	+ .0271	0014		+ .17	1	+ .234	- '021	18	8.07 8.84	3586 3588
717	2.6	Centauri	48 20 200		3.7195	- ·0005 + ·0471		+18 25 32.08	$+ \cdot 11$ + $\cdot 38$	17.857	*256	- ·013 - ·053	17 21		
719	2.7	8 Boötisη		+ .038	2.8567	- '0003		+18 53 52.81	+ 3.11	17 303	• 199	367		7 <sup>.</sup> 19 8.53 : 8.48	3593 3596
720	4.7	Lacaille 5733			4.2927	+ .1003	0023	-63 11 47.51	+ .46	17.817	*297	- '052	33:34 17	8.81	3599
				1 04/					1 40				-/		
721	<b>6.1</b>	92 Virginis			+ 3.0233			+ 1 32 22.42	11	-17.214		+ .015	16	8.77	3600
722	4.0	Centanri $\phi$		+ .012	3.6281					17.712	•256	- '020	19	7.27	3602
723	4.0	Centauri	52 30.028		3.6844	•0430	0031	-44 18 55.78		17.712		035	19	7.83	3603
724	5.3	47 Hydræ		+ .028	3.3266	•0214	0036		+ 32	17.704	•238	041	19	7.88	3604
, 725	6.0	48 Hydræ	54 23.906	+ .153	3*3490	•0214	- '0152	-24 31 21.22	+ .00	17.712	•240	111	19	8.02	3607
726	var.	Apodisθ	13 55 34 288	+ .242	+ 5.6981	+ .2975	0261	-76 18 50.82	+ '37	-17.201	+ .407	- '040	32	9°27	3611
727	4.3	93 Virginisτ	56 33.400	- •009	3.0206	+ .0062	+.0013	+ 2 1 41.98	+ .18	17.535	•224	022	22:23	7'19:7'14	3612
728	6.3	11 Boötis	56 38.365	+ .042	2.7220		-••0060	+27 52 10.48	- '02	17.203	• 200	+ .003	19	7.26	3613
729	0.2	Centauriß		+ .033					+ .30	17*533	55	035	17	9.46	3615
730	4.6	Centauri $\chi$	13 59 56.345	110. +	3.6453	•0378	0019	-40 42 1.79	+ .53	17.398	• 273	034	20	6.62	3621
731	3.4	49 Hydræπ	14 0 40.524	025	+ 3.4059	+ .0230	+.0031	-26 12 3.13	+ 1.27	-17'491	+ .257	160	19	7.96	3622
732	2.0	5 Centauri		+ .404	3.2120			-35 52 45.76		17.854	• 262	528	16	9.27	3623
733	6.9	94 Virginis	0 59 980		3.1714			- 8 24 51.27		17.308	•240	+ .000	16:15	9.29 : 9.24	3624
734	5.0	Apodis	5 39*		7 '2354			-80 32 20.51	+ .59	17.186	• 556	078	5	7.21	3633
735	4.9	12 Boötisd	5 50.221	+ .015	2.7370	- '0017	0017	+25 33 54.74	+ .21	17.12	.215	072	26	7.03	3635
736	4.2	98 Virginis						- 9 48 28.95	- 1.06	-16.891	+ .253	+ .130	37	8.22 : 8.12	3642
730	4 2 4.1	99 Virginis		- °004 + °009	3.1409			- 9 48 28 95 - 5 31 27.57		17.298		- :427	21:23	7.33:7.22	3660
738	4 I 4 I	Octantisδ		+ .60	3 1409 9.087			-3312/5/ -831235.12		16.880		- '013	44:47	11.63 : 11.37	3661
739	0.0	16 Boötisa		+ .578	2.7352				+14.82	18.858		-2.003	24	7'40	3662
740	3.8	Lupi		+ .007	3.8196				+ .06	16.773		008	20	7.58	3668
										16:000	1	1001			1670
741	4'4			-	+ 4.152		- '0029	-55 55 32.64	+ .15	-16°770 16°708		- °021 + °023	21	7.01 8.89 : 8.70	3670
742		100 Virginisλ		+ .013	3.2396			-12 54 39.09	- •20	16.708	3*396		23:20 38:55	11.63 : 10.37	3672
743	8.1	Brisbane 4614	13 48.13		42.175	34.972		-88 55 14.51	+ •17	16.602		 - '022	3° : 55 19 : 20	7.84 : 7.73	3686
744	5.8	Lacaille 5890a		+ .019	4.9054			-67 44 25 49 -39 3 18 44		16.617		040	23	. 6.94	3688
745	4.6			1- 01/	3.6292				1000						
746	6.6	2 Libræ		+ .000				-11 15 27.24		-16.283		064	18:19	9'17 : 9'11	3691
747	7.5	Lacaille 5921		+ .033	3.7128	0.0	-	-40 18 2.94		16.238		042	16	9.11	3693
748	5.2	Lacaille 5929		+ .039	3.4119			-24 21 9.09		16.497	- 1	031	18	7.12	3695
749	4'7	Lupi $\tau^1$		+ .008	3.8305			-44 46 8.56		16.465	0.1	029	17	8.57	3699
750	4'4	Lupi $ au^2$	19 44 843	005	3:8361	•0442	+.0003	-44 55 37.75	+ .19	16.424	327	020	20:21	7.86 : 8.06	3700

701. 4.4, 4.7 0"3 105° 1897.2. 726. L, 5.5-6.5; P, probably irregular.

.

.

## FROM OBSERVATIONS IN THE YEARS 1905-11.

-

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_{a}\Delta E.$	Annual Variation 1900 °O.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
751 752† 753†	5°6 5°1 5°0	22 Boötisf 52 Hydræseq. 105 Virginispr, φ	23 2.864	s + ·045 + ·014 + ·066	+ 2.7900 3.5022 3.0877	+ ·0010 ·0251 ·0088	- · 0019 - · 0089	$+19^{\circ} 40^{\circ} 35^{\circ} 14$ -29 2 32.43 - 1 46 47.49	- "13 + 24 + 07	-16.315 16.337 16.277	• 304 • 269	+ <sup>".</sup> 015 033 010	19 24 23	8·58 7·13 7·45	3705 3707 3710
754 755	4.6 3.8	Lupiσ 25 Boötisρ	25 52.641 27 31.131	+ •039 + •056	4.0141 2.2862	+ ·0539 - ·0015		-50 0 49.68 +30 48 37.86	+ .09 79	16°134 15°925	*354 *232	- ·013 + ·110	28 21	7·20 7·16	3716 3717
756 757 758 759 760	2.4 6.3 4.6 4.1 6.5	Centauriη Lacaille 5994 28 Boötisρ Lupiρ Mayer 592	30 19.688 31 9.425	$+ \cdot 022$ + $\cdot 008$ - $\cdot 131$ + $\cdot 030$ + $\cdot 554$	+ 3.7912 3.7772 2.6132 4.0105 3.1857	$\begin{array}{r} + & \cdot 0390 \\ + & \cdot 0381 \\ - & \cdot 0014 \\ + & \cdot 0515 \\ & \cdot 0131 \end{array}$		-41 43 7.40 -41 4 42.94 +30 10 47.28 -48 59 24.21 -11 52 45.88	$ \begin{array}{r} + & \cdot 25 \\ + & \cdot 23 \\ - & 1 \cdot 06 \\ + & \cdot 31 \\ - & 3 \cdot 36 \end{array} $		+ '341 '340 '240 '365 '287	- '035 - '030 + '120 - '037 + '364	25:26 16 17:18 16:17 17:18	7'17:7'13 7'62 8'80 8'35:8'39 9'32:9'22	3724 3725 3729 3732 3734
761* 762*	0'3 8'7*	Centauri	14 32 44·344 34 18·64	+4.547	+ 4.0412 30.656	+ <sup>.</sup> 0731 14.641	-·4874	-60 25 7.39 -88 16 10.03	- 6.78	-15.025	+ • 338	+ •729	17:18 15	9'33 : 9'30 11'68	3735
763 764 765	3·2 2·3 3·7	Cireinia Lupia Apodisa	35 16.604	+ ·291 + ·018 + ·060	4 7927 3 9674 7 2319	·1123 ·0473 ·4353		64 32 25°21 46 57 32°26 78 37°13°07	+ 2.20 + .24 + .26	15°902 15°647 15°638	• •440 •369 •667	- 236 - 028 - 027	16 : 18 17 : 18 32	9'39 : 9'31 8'69 9'47	3739 3745 3746
766† 767 768 769	3.8 7.8 5.4 .4.1	30 Boötism. ζ Lacaille 5882 Lacaille 6039 Centauri	37 21·748 37 32·324	- · 031  - · 076 + · 056 - · 063	+ 2.8636 15.332 4.7054 3.6543	+ ·0033 3·023 ·0992 ·0303		+14 9 25.60 -86 3 44.25 -62 26 56.43 -34 44 36.82 - 5 13 28.00	+ ·21  + ·71 + 1·76	-15°586 15°545 15°589 15°687	+ ·270 1·418 ·442 ·344	- · 027  - · 085 - · 193 - · 322	15:17 17:37 16:17 17	8.13 : 7.66 11.63 : 9.55 8.32 : 8.39 9.10	3752  3755 3757
770 771 772 773	3.9 6.8 5.0 6.0	Octantis	39 1.652	+2.10	3.1571 +24.564 2.6369 3.9866	·0107 + 8·762 ·0001 ·0471	180 0008	-87 44 31 02 +26 57 10 52	+ 2.82 + .61 + .19 + .26	15°802 -15°480 15°433 15°399	·300 +2·279 ·252 ·379	$ \begin{array}{r} - \cdot 322 \\ - \cdot 067 \\ - \cdot 021 \\ - \cdot 031 \end{array} $	19:18 39:116 16:15 19:20	8.85 : 8.75 11.61 : 9.01 9.17 : 9.18 8.30 : 8.33	3758 3760 3761 3763
774 775	5°3 6°8	Lnpi,b Mayer 596b		+ .036	4°1676 3°3950	·0581 ·0187	0032	-20 45 8.04		15°445 15°446	· 397 · 325	000	16 16	9°43 9°63	3765 3769
776 777 778 779 780	3.8 5.5 2.8 var. 5.3	109 Virginis           8 Libræ           9 Libræ           α           Lacaille 6077           Lacaille 6119	45 9°209 45 20°644 46 28°605	+ .061 + .057	+ 3.0298 3.3112 3.3118 6.6828 4.5824	+ .0074 .0155 .0155 .3127 .0836	- ·0071 - ·0074 - ·0185	+ 2 18 50.69 -15 34 54.37 -15 37 35.30 -76 15 19.09 -59 42 12.77	+ 33 + 67 + 57 + 09 + 81	-15.328 15.141 15.128 14.997 15.017	+ ·292 ·324 ·324 ·652 ·451	- ·038 - ·078 - ·076 - ·011 - ·111	19 17 33:32 35:34 18	8°73 8°58 7'74 : 7'44 8°63 : 8'62 7'27	3772 3784 3787 3794 3800
781 782 783	6.0 5.5 5.8 6.1	13 Libræ Lacaille 6146 15 Libræ Piazzi XIV. 221	49 36°350 51 20°450	- ·012 + ·001	+ 3.2508 3.6693 3.2490	.0130	0001	-11 0 22.33		-14.864 14.816 14.703	+ ·325 ·367 ·328	$- \cdot 022$ $- \cdot 013$ $- \cdot 002$	19 19:22 32	8.56 7.79:7.77 9.54:9.59	3804 3807 3810
784 785 786	6·3 2·7	Piazzi XIV. 221 Piazzi XIV. 212 seq. Lupiβ	51 38.154	- ·657 + ·034	2.8299 3.4952 + 3.9099	·0036 ·0207 + ·0392			$+ \cdot 12$ +15.26 + .40	14.707 16.438 -14.714	·287 ·360 + ·395	$- \cdot 016$ $-1 \cdot 754$ $- \cdot 051$	23 17:18 18	7 · 52 8 · 86 : 8 · 87 7 · 81	3811 3813 3815
787 788 789	3°2 var. 3°4	Centauriκ 19 Libræδ Scorpii 1 Hγ	14 58 12.895	+ .038 + .042	3°8855 3°2000 3°5018	•0377 •0116 •0209	- · 0046 - · 0056	-24 53 20.94	+ ·30 + ·09 + ·41	14.653 14.454 14.340	· 393 · 329 · 364	- ·032 - ·011 - ·055	16 20 27	9°27 8°34 7°42	3818 3825 3837
790 791 792	4°6 5°0 4°0	<ul> <li>43 Boötisψ</li> <li>45 Boötisκ</li> </ul>	15 2 54·621 4 58·746	+ ·100 - ·115 + ·085	2.5702 + 2.6346 4.1479	· 0012 + ·0017 ·0475	+.0138	+27 20 14.75 +25 15 29.49 -48 21 27.61	+ ·15 + 1·53 + ·55	14°185 -14°177 13°930	•270 + •282 •442	- ·020 - ·183 - ·066	20 : 23 23 17 : 18	7.52:7.37 8.35 8.38	3842 3855 3862
793 794 795	3 <sup>.</sup> 4 5 <sup>.</sup> 0 4 <sup>.</sup> 7	Lupiζ Lupi	5 5·816 6 6·413	1	4·2846 4·0093 3·4121	•0548 •0401 •0171	- °0115 - °0036	-51 43 7.73	+ ·49 + ·28 + ·44	13°925 13°825 13°815	·456 ·430 ·367	- ·069 - ·033 - ·049	23:26 18 37:38	7 <sup>.24</sup> : 7 <sup>.17</sup> 8 <sup>.48</sup> 8 <sup>.95</sup> : 8 <sup>.93</sup>	3864 3865 3866
796 797 798	5°1 2°9 4°2	1 Lupi Trianguli Austγ Circiniβ	9 34 056 9 40 823	+ .100	+ 3.6645 5.5340 4.6611	+ ·0251 ·1397 ·0750	0126	-31 8 44.92 -68 18 36.60 -58 25 40.86		-13.658 13.595 13.711	+ ·397 ·598 ·505	- ·018 - ·024 - ·148	19 24 : 26 20	8·18 ,7·30:7·23 7·94	3871 3879 3880
799 800	5.6 3.4	3 Scrpentisδ 49 Boötisδ		1.	2·9793 2·4188	•0066 •0012	0014	+ 5 18 37 56 +33 41 15 13	+ .08	13.538 13.575	•326 •268	- ·009 - ·127	16 : 18 16	8·82 : 8·69 8·93	3882 3887
		752. 5'1, 11	·I 4 <sup>//</sup> ·2 2	79° 1	901.4.				766. 4'4	, 4.8; very	close bina	ry.			

.

752. 5'1, 11'1 4'2 279° 1901'4. 753. 5'0, 9'3 4'''7 110° 1904'4. 761. Reduction to C. G.,  $-0^{3}$ '704, -7'''74. 762. Gilliss magnitude = 7'6. 766. 4'4, 4'8; very close binary. 779. L, 5'5-6'2; P, unknown. 788. L, 5'0-5'9; P, 2<sup>d</sup>'33.

.

C. F. C., 1900.

С

.

17 .

CAPE FUNDAMENTAL CATALOGUE OF STARS FOR 1900.0,

N	o. Ma	g. Name.	Mean R.A. 1900'0.	$\mu_{a}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900°0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900°0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
80	I 2.0	5 27 Libræ	hms 151137.424	s + .060	s + 3.2230	s + .0118	s -•0067	- 9 ó 50 <sup>°</sup> .87	+ "27	-13.468	- "	"	25.120	9'02 : 9'09	3890
80				+ .016	3.9173	.0344	0022	-40 25 18.30	+ '23	13.400	+ .354 .	- ·030	35 : 39 22 : 24	7'29:7'16	3890
80			14 48.396	+ .004	3.9228	•0340	0006	-40 17 7.70	+ .23	13.263		- '032	25:26	7'04:7'06	3896
80				+ .012	4.3323	.0525	0020	-51 22 37.53	+ .17	13.185		- •023	22:23	7.40 : 7.45	3906
80		B Lupi $\phi^2$	16 45 920	+ .014	3.8190	•0294	0018	-36 29 59.91	+ .53	13.135	•427	- °031	18:19	7'55:7'50	3910
80				+ .001	+ 3.3398	+ .0145	- '000I		- ·01	-13.022		+ .001	31:32	9'74:9'68	3913
80 80			20 12.91 20 36.583	-1.04 + .002	13·130 6·4371	1·404 ·2073	+.089	-84 7 53.88 -73 2 33.58	72 + .18	12·792 12·868		+ ·081	41 : 143 19 : 20	11.68 : 8.95 7.93 : 7.86	3924
80	1 .	-	21 9.060	1	2.7804	.0040	0014		+ 10	12.835		- '026	19:20	8.50 : 8.46	3925 3931
81				010	3.3770	•0148	+.0011	-16 22 4.90	+ .38	12.753		- ·043	20:21	8.95 : 8.92	3935
81	1 3.7	3 Coronæ Borealisβ	15 23 42.232	+ .094	+ 2.4732	+ .0010	0133	+29 27 1.94	54	-12.260	+ .284	+ .076	26:27	7.08:7.05	3940
81	10,		27 33.946	025	5°4355	1126	+ 0034	-65 58 50.64	+ .52	12 300		072	24:25	7'25:7'19	3940
81	37 2.8		28 28.506	+ .010	3.9824	.0330	0014	-40 49 50.07	+ .26	12.345	•463	036	r8	7.30	3950
81				+ .019	2.4180	*0020	0020	+31 41 47.40	+ .21	12.306		— ·026	17	8.11	3953
18	5 4'1	38 Libræγ	29 55.896	032	3.3201	.0136	+.0042	-14 27 22.02	+ .01	12.310	·393 ·	001	38:35	8.24 : 8.07	3959
81	6 2.2	5	15 30 27.259	071	+ 2.5391	+ .0024	+.0000	+27 3 2.82	+ .81	-12.274	+ .300	- '102	18	7.94	3961
81			30 57.109	+ .002	3.6325	•0209	- '0007	-27 48 14.04	+ '04	12.143	7-7	005	17	7.77	3962
81			31 18.743	+ .112	4.0249	•0341	0144	-42 14 20'31	- '50	12.021		+ .001	16	8.12	3964
81				+ .032	4.4342	.0213	0036	-52 2 34.13	+ .37	12.149	5	- '042	17	8.90	3965
			35 22.911	+ .019	4.3183	•0445	0022	-49 10 3.21	+ .24	11.860	.213	033	31:32	7'21:7'17	3987
82				+ .027	+ 3.4492	+ .0122	0035	-19 21 18.21	+ .99	-11.889	1 412	- •119	19	8.30	3990
82			37 5.455	+ .042	2.6725	•0036		+19 59 31.82		11.261		055	20	8.26	3994
82			38 32·506 39 20·582	+ .057	2·5188 2·9522	·0026 ·0061	+.000	+26 36 44 82 + 6 44 24 56	- ·23 - ·31	11.508	• 303 • 358	+ .030	22 23:22	7 * 57 8 * 36 : 8 * 28	3998 4001
82			41 34.347	040	2 9522	.0043		+15 44 4.18	+ '46	11 300		057	22	8.11	4009
82			1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1					+18 27 0.30						PIOT - P-66	
82		~	44 23.988	+ .025 + .055	+ 2.6992	+ .0039	- ° 0032 - ° 0059	-3727.96	+ '77	II'209	1 33	- '101 - '028	20 26	7'91 : 7'66 9'25 : 9'28	4015 4016
82			44 36.142	+ .000	3.8014	·0237	- '0007	-33 19 21.66	+ .24	11.194		028	18	8.63	4018
82	9 3.	7 37 Serpentis	45 49.887	068	2.9875	•0065		+ 4 46 43.25	48	11.020		+ .057	20:21	8'20:8'49	4026
83	0 2.8	B Trianguli Austβ	46 19.468	+ .215	5.2436	.0872	0296	-63 7 21.80	+ 2.79	11.429	•639 -	- • 388	22:23	7.26 : 7.19	4030
83	I 5'1	45 Libræλ	15 47 31.656	+ .008	+ 3.4758	+ .0121	0010	-19 52 5.75	+ .28	-10.982	+ .429	034	23	8.30	4033
83			50 42.498	+ .007	3.6954	•0199	0010	-28 55 19.57		10.749	•460	030	32:33	7'43:7'38	4052
83			551		2.7688	.0057	+.0210	+15 59 5.48	+10.29	11.933	·349 ·	-1.297	27	8.46 : 8.32	4055
83			52 48.045		3.6210	.0128		-25 49 34.66		10.000		036	20	8.23	4062
83	57 4.2	-	53 26.745	+ .048	2*4820	.0031	- '0064	+27 10 2.14	+ .20	10.284	.312 -	- •068	24:25	7'43:7'35	4063
83				+ .002	+ 3.2402	+ .0128		-22 20 14.07	+ .30	-10.485	1 115	039	31:26	8.31 : 7.67	4066
83				+ .356	3.3609	.0133			+ 3'23	10.851		400	22	8.08	4067
83 83			56 44.643 59 25.323		2.6936	•0038		+18 5 41.60	- 1'19	10.122		+ •144	24 28:29	8·23 7'35 : 7'45	4075 4084
84			59 25 323 15 59 37 267	- ·001 + ·007	4°2242 3°4817	•0331 •0141	+.0001	-44 54 6·25 -19 31 55·01	$- \cdot 13$ + $\cdot 25$	10.020		+ .017	35:32	8.84:8.52	4086
11												1. 1. 5	1. 2.		
84 84					+ 3.9281	+ .0244	- '0017	-36 31 48.41	+ .27	-10.022	+ .200 -	035	18:19	7.89:7.84 7.59:7.61	4091
84	-		I 32.375 3 33.655		3.5120	*0145 *0041		-20 35 55.31 +17 18 47.40	+ '43	9·963 9·768	•450 - •348 -	- ·056 - ·014	30 : 31 22 : 23	8.13:8.17	4095 4101
84			4 28.516		4.0744	• '0278		-40 51 17.91		9.811		129	29:30	7'44:7'39	4105
84			5 23.566		8.8018		1	-78 26 38.10	+ 34	9.650	00	037	33	9.06	4109
84	5 5.2	Normæκ	16 5 35.351	+ '013	+ 4.7080	+ .0479	0016	-54 22 18.18	+ .32	- 9.640	+ .606 -	043	21:22	8.19 : 8.13	4111
84		3 13 Scorpii	6 8.510	+ .018	.3.6862			-27 40 1.19	+ 33	9'593	.476 -	038	17	8.70	4115
84	3 4.1	Trianguli Austδ	6 19.976	003	5.4231		+.0004	-63 25 48.36	+ .14	9.558		- •018	18	7.56	4118
84		-	9 6.240		3.1401			- 3 26 14.29		9.479	•409 -	123	41:43	7.91 : 7.65	4134
85	ot 5.7	Normæ	12 19.984	+ .002	4.1602	.0278	0009	-42 25 44.65	+ •14	9.094	•544 -	010	24:26	7.26 : 7.15	4144
							-				176-5-				
		813. 3'5, 3'7			0.10			835. 4.2, 12	2"*0	352° 1	905 4.				
		822. 5°4, 5°4 823. 3°9, 6°9	; very close bin		93*5.			840. 2°6, 10 850. 6°1, 6°9	1"'1 0"'4		901'4. 897'2.				
		828. λ in Au	wers' Bradley.							S.U.S.	272				

FROM OBSERVATIONS IN THE YEARS 1905-11.

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_{\alpha}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900 °O.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
851	4'I	Normæ	hm s 16 12 21 211	s + •146	s + 4.4701	+ ·0373	s 8	-49 54 37'25	+ "43	- 9.126	+ ". 583	- "053	21:22	8'10 : 8'02	4145
852	3'1	2 Ophiuchie	13 1.807	045	3.1705	.0081	+.0023	- 4 26 55.71	- '27	8.989	.417	+ .032	23	8'48 : 8'29	4147
853	7.0	Lacaille 6783		+ .002	4.3976	·0341	0009	-47 56 52.22	+ .28	8.972	• 578	034	17:19	8.55 : 8.10	4153
854	5.6	Lacaille 6790		+ .008	4.4664	.0358	0011	-49 20 0.65	+ '23	8.898	•588	031	21:24	7 '53 : 7 41	4156
855	3.0	20 Scorpiiσ	15 6.217		3.6392	.0124	0009	-25 21 10.57	+ .54	8.890	•480	032	25	7.65	4158
856	4'9	50 Serpentisσ			+ 3.0347	+ .0066		+ 1 15 50.34	- '35			+ .043	16	8.02	4163
857 858	5.2	Lacaille 6810 20 Herculis	17 14·874 17 30·466	- ·054 + ·030	4.0534 2.6449	·0237 ·0038	+.0072 0034	-385732.83 +192316.33	+ '14	8·709 8·630	·537 ·351	- ·019 + ·039	22:23 23:24	7'52:7'39 8'86:8'74	4164 4165
859	3°7 5°1	Trianguli Aust	17 30 400		6.3990	.1158	+.0397	-69 51 31.13	83	8.553	.851	+ .100	18	8'31	4165
860	3.8	Apodis	18 5.732		9.0554	• 3205	0400		+ .64	8.699	1.190	077	34:35	8.40 : 8.34	4168
861	4'9	19 Coronæ Borealis ξ	16 18 11.939	+ .069	+ 2.3363	+ .0030	0074	+31 7 27.29	85	- 8.522	+ .310	+ .092	17	9°27	4169
862	6.5	23 Herculis		010	2.3012			+32 33 58.14	+ .17	8.561	.307	018	16	9.56	4176
863†	4.6	24 Herculisseq. w	20 48.033	022	2.7669	.0045	+.0030	+14 15 47.59	+ .47	8.474	• 370	062	23:24	7.30 : 7.26	4182
864	5'4	Lacaille 6824		+ .010	5'3024	•0616		-61 24 43.28	+ .02	8.322	. 707	003	19:20	7.84:7.75	4185
865	5.6	Lacaille 6841	22 27 479	+ '013	4'3323	•0295	- '0015	-46 I 16·54	+ .08	8.287	.578	010	19	8.40	4190
866	6.6	Lacaille 6441	16 22 50.86		-+29.826	+ 4.962		-87 23 34.64		1	+3.967		31:42	11.25 : 10.31	
867†	0.8	21 Scorpiiseq. a	23 16.486	+ '004	3.6720	·0149	0005	-26 12 36.74	+ '28	8.245	.492	033	20	8.34 11.51 : 8.68	4193 4196
868 869	6.3 4.4	Lacaille 6545N	23 34'91 24 50'787	- ·08	21·343 3·9117	2°353 °0192	+.007	-86 10 42.85 -34 29 11.63	+ '01 + '20	8.110	2.845	- '001 - '024	47:157	8.46	4190
870+	4.0	10 Ophiuchim. λ	25 52.151	+ .028	3'0220	*0063		+ 2 12 8'97	+ .74	8.088	.407	084	18:17	8.83	4203
871	2.6	27 Herculis		+ •071		+ .0036		+21 42 26.29	+ .23	- 8.024	+ .347	024	17:16	9'50:9'45	4204
872	5.2	Normæµ	26 58 550	000	+ 2.5770		+.0011		+ '08	7.924	• 573	000	17	8.46	4208
873	4'2	Apodis		+ .710	8.4547	•2440	0880	-77 18 31.67		8.115	1.128	346	40:41	8.02 : 8.10	4215
874	2.8	23 Scorpiiτ		+ .002	3'7279		-•0008	-	+ .30	7.736	. 202	032	19:21	8.30 : 8.22	4218
875	2.2	13 Ophiuchiζ	31 39.098	- •006	3.2996	•0086	+.0008	-10 21 52.70	13	7.221	•449	+ .012	22:21	7.89:7.82	4225
876	5.2	24 Scorpii		+ .014	+ 3.4651	+ .0103	0012	-17 32 55'47	+ .06	- 7.208	+ .474	007	22	8.38	4239
877+	2.8	40 Herculis		+ • 298	2.2608	•0027		+31 47 5.03	- 3.16	6.675	•306	+ .385	22:20	8.16 : 8.22	4246
878 879	1.4 7.0	Trianguli Austa Lacaille 6953	38 4°425 38 46°457	- °025 - °002	6.3107	•0889 •0258	+.0032	-68 50 38.83 -46 20 46.63	+ '20	7·042 6·998	·865 ·603	- ·027 - ·041	18:20 19:21	7'74:7'56 8'25:8'08	4250 4252
880	3.7	Aræη	41 8.922	- '033	4°3863 5°1593		+ 0002	-58 51 46.07	+ 33 + 33	6.807	.712	- '041	28:30	7.59:7.40	4265
1.1		Gilliss P.Z. 11448					1.1.1.1							11.43 : 10.58	
881 882	7·9* 7·1	18 Ophiuchi		 + '007	+66·525 3·6461	+21·427 ·0118		-88 51 49°91 -24 27 54°21	+ .19	- 6.751 6.578	+9.134	- '023	35:45 17:18	8'15:8'13	4271
883	2.1	26 Scorpii		+ .374	3.8783		-		+ 1.97	6.811		258	24:25	7 55 : 7 65	4272
884	4.8	20. Ophiuchi	44 18.102		3.3149			-10 36 23.34	+ .85	6.604	'461	- '102	17	8.29	4273
885	3.1	Scorpii	45 5'721	+ .002	4.0562	•0177	0006	-37 52 32.99	+ .26	6.466	• 563	030	19:20	8.87 : 8.68	4277
886	5.7	47 Herculisk	16 45 28.032	032	+ 2.9109	+ .0048	+.0035	+ 7 25 12.51	+ .07	- 6.413	+ '405	008	16	9.26	4280
887	6.7	49 Herculis	47 31.660	002	2.7292	•0039		+15 8 30.85	+ .06	6.241	.381	007	44:39	8.49 : 8.14	4291
888	3.5	Scorpiiζ <sup>2</sup>	47 32.632		4.2128	•0204			+ 1.93	6.470	. 585	- *237	14:15	8.18 : 8.13	4292
889	6.8	Lacaille 7024	48 25.946		4.6140	.0274		-50 30 44.78 +31 52 1.18	+ .23	6·184 6·120	·642 ·317	- ·025 - ·023	16:18 16	9'49 : 9'29 9'45	4296 4300
890	5.6				2.2734	.0033		1. C	+ '22						
891	4'3	25 Ophiuchi		+ .034	+ 2.8365			+10 19 47.17	+ '41	- 6.134	+ .396	- •046	16:17	8·90 : 8·86 7·93 : 7·84	4302
892 893†	3.0 5.7	Aræζ 24 Ophiuchi	50 20°560 50 46°086	+ .001	4°9483 3°6132	•0342 •0104		-55 49 55.90 -22 59 29.74	$+ \cdot 3^{2}$ + $\cdot 0^{3}$	6.041 5.968	•690 •505	- ·041	22:23 19:21	7 93 : 7 °4 8 '49 : 8 '22	4304 4309
894	4.1	Aræε <sup>1</sup>	51 36.712		4.7675	.0293			- ·01	5.893		+ .001	19	8.81	4313
895	3.2	27 Ophiuchi	52 55.908		2.8376	.0043		+ 9 31 49'12		5.797	• 396	014	26:29	7.85:7.45	4315
896	5.3	Lacaille 7089	16 55 24.570	+ .002	+ 3.8734	+ .0130	0009	-31 59 41.95	+ .47	- 5.637	+ .545	062	23:25	7.83:7.66	4321
897	5.1	30 Ophiuchi	55 47.132	+ .031	3.1602	•0060		- 4 4 22.62		5.632	•444	088	21	8.60	4323
898	3.8	58 Herculise			2 · 2941	.0031	-	+31 4 24.74	1	5.466		+ '021	17:20	8.56 : 8.36	4328
899	5.4	59 Herculisd 60 Herculis		•000	2'2131	•0032		+33 42 46.61		5.376	*313	- '012	19	8·42 7·58 : 7·50	4332 4346
900	4'9	00 H0100113	17 0 44 404	027	2.7805	.0038	+ 0030	+12 52 40.77	+ .13	5'143	• 394	- '017	30:31	7 50 . 7 50	4340
$\begin{array}{cccccccccccccccccccccccccccccccccccc$															

CAPE FUNDAMENTAL CATALOGUE OF STARS FOR 1900.0,

No.	Mag.	Name.	Mean R. A. 1900°0.	$\mu_{a}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation • 1900'0.	° Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900 <b>+</b> .	Boss No.
901†	0.4	35 Ophiuchim. η	h m s	8	8	s + •0071	s + .0022	-15 36 3.60	- "68	- 4.709	+ "489	+ "086	25	8*45 : 7*96	4160
901	2.4	35 Opinteinη Scorpiiη		- ·021	+ 3.4372	•0169		-43 6 28.51	+ 2.30	- 4 /09 5'060	•610	- '294	25 24	7.84	4360
1	3.3	C.G.A. 23027	4 59°394 6 3°75		30.294	2.782		-87 17 47.18		4.674			31:4I	11'43 : 10'59	4361
903 904	7·7 5·7	Lacaille 7202		+ .060	3.8979	•0106	- '0077	-32 32 59.99			4°341 °556	 '055	22	7.78	
904	-	65 Herculisδ			2.4628	*0033		+24 57 23.75	+ '43	4'347	352	163	15:16	9.01 : 8.78	4375 4376
905	3.1	05 Hereuns	10 55 304	+ .019	2 4020	0033	- 0018	T24 3/ 23 /3	+ 1.43	4.423	332	- 103	19:10	901.070	4370
906	5.8	Apodis	17 10 56.518	+ .020	+ 6.6614	+ .0614	0025	-70 1 4.21	+ .14	- 4.276	+ .950	- '017	18	7.98	4377
907	6.1	Lacaille 7088	12 45		11.1508	2378	+.0012	-80 45 58.67	+ '33.	4.147	1.589	043	3	7.28	4387
908†	var.	68 Herculispr. u	13 37.846	+ .014	2.2138	·0031	0016	+33 12 27.87	+ '11	4.041	• 318	013	20	8.82	4388
909	4.5	40 Ophiuchiξ		130	3.5928	.0074	+.0121	-21 0 22.29	+ 1.26	4.112	.218	- 207	25:26	7.62:7.52	4394
910	3.2	42 Ophiuchi $\theta$	15 52.047	100. +	3.0811	·0078	0001	-24 53 59.49	+ .29	3.867	• 528	- '031	27:26	9.60 : 9.37	4399
911	e.e	72 Herculisw			1	+ .0048	+.0099		1 0.12	1.200	1 1224	-1.053	16	8.67	
912	5.5	Lacaille 7247		086	+ 2.2427	+ ·0048 ·0136	0022	+32 35 37.77 -44 3 59.62	+ 9'13 + '27	- 4'799	+ '324	033	10	8.32	4403
913	5.3	Aræβ			4.3410	·0217	0018	-44 3 59.62 -55 26 7.08	+ .27	3.774	.714	033	18:19	8.25:8.05	4404
913	4.2		16 59.146		4.9763		0002		+ 1.02	3.773	.527	- '132	27:28	8.02:7.93	4406
1.		44 Ophiuchid	20 15.762 20 58.087		3.6603	.0073	+.0014			3.200			22:21	8.42 : 8.33	4420
915	4'3			- '012	3.8274	•0083	T 0014		+ 1.30	3.224	.552	120	22:21	0 42 : 0 33	4421
916	4.5	49 Ophiuchiσ	17 21 33.156	002	+ 2.9751	+ .0032	+.0002	+ 4 13 37.38	03	- 3.344	+ .429	+ .003	21:19	9.77 : 9.58	4425
917	3.6	Aræδ	22 4'131	+ .001	5.4042	.0254	0076	-60 36 1.46	+ .68	3.388	.778	085	17	8.02	4426
918	2.7	34 Scorpii			4.0746	•0094		0, 0, ,	+ .35	3.181	. 588	042	18	8.39	4429
919	2.7	Aræa	24 6.625	+ .029	4.6309	·0146	0034	-49 47 48.87	+ .71	3'210	•668	084	21	8.40	4431
920	4.9	51 Ophiuchi	25 18.827	001	3.6575	.0064	+.0001	-23 53 7.68	+ .32	3.060	.528	038	18:19	8.59 : 8.55	4434
921	4.7	76 Herculisλ	17 26 41.700	010	+ 2.4232	+ .0028	+.0011	+26 11 9.38	- '12	- 2.889	+ .351	+ .014	16	8.78	4438
922	1.2	35 Scorpiiλ			4.0697	•0087	0003			2.928	.588	036	20	7.91	4439
923	4.7	Aræσ			4.4602	·0114	0039	-46 26 12.04		2.811	•645	039	21:22	8.38:8.34	4445
924	5.4	Aræπ			4 4002	·0154	0027	-54 26 0.62		2.778	•712	151	17	8.10	4454
925	6.6	Lacaille 7078		+ .29	18.747	.538	026	-85 10 35.89		2.748	2.707	- '131	33:46	11.45 : 10.60	
				T 29	1	530									1.50
926	1.8	Scorpiiθ		003	+ 4.3056	+ .0096		-42 56 2.70		- 2.616	+ .624	011	17:18	8.74 : 8.72	4457
927	2.0	55 Ophiuchia		077	2.7833	.0033	+.0080			2.826	•405	- *235	19:18	9.64 : 9.53	4459
928	3.4	55 Serpentisξ		+ .028	3.4331	.0046				2.526	.497	071	19:17	9'43:9'21	4462
929	5.0	Aræλ		085	4.6291	.0119	+.0111	-49 21 14.69		2.264	•673	129	20	7.69	4466
930	2.4	Scorpii ĸ	35 34.173	+ .007	4.1467	.0072	0009	-38 58 42.48	+ '21	2.129	•602	026	20	8.26	4474
931	4.4	56 Serpentis 0	17 35 47 590	+ .043	+ 3.3700	+ .0040	0048	-12 49 19.65	+ .50	- 2'170	+ .489	056	16	9.00	4475
932	3.5	Pavonis			5.8797			-64 40 33.41		2.158	.853	055	16:17	9'20 : 9'11	4476
933	5.3				4.7589			-51 46 52.20		2.279		- '201	16	9.47	4478
934	2.9	60 Ophiuchi			2.9623	.0027	-	+ 4 36 33.19		1.724	•430	+ .152	33	8.53 : 8.55	4487
935	3.0	Scorpii			4.1935	.0062				1.694			19	7.83	4492
- 10 A				-			and the second				_	1		Store Lan	- 52
936	var.	3 Sagittarii(X.)				+ .0042	0004	-27 47 33.85	+ .18	- 1.659		022	18	8.03 .	4493
937	3.4	86 Herculis			2.3460	.0038		+27 46 37.64		2.276	•338	750	16	9.39	4497
938	3.8	62 Ophiuchiγ	42 52.674		3.0067	.0029		+ 2 44 40.30		1.222	•437	079	17	8.50	4500
939	3.1	ScorpiiG			4*0825	.0021				1.461	• 595	+ .020	17	9.08	4501
940	5.0	Scorpii	43 11.479	004	4.1934	•0056	+.0004	-40 3 29'12	+ .13	1.483	.610	014	17	9.22	4503
941	5.3	87 Herculis	17 44 45.801	+ .007	+ 2.4310	+ .0025	0008	+25 39 21'00	+ '41	- I.377	+ .354	045	17	9.12	4508
942	6.6	Mayer 722	50 2.038		3.5276	·0028		-18 47 4.87		0.892	.514	- '021	16	8.92	4521
943	5.0	Lacaille 7497				.0039		-41 42 6.91		0.842	•621	028	23	8.08	4525
944	5.8	89 Herculis			2.4192			+26 3 57.03		0.421	*353	+ .002	16	9.26	4528
945	3.4	64 Ophiuchi	53 31.271		3.3014	*0025		- 9 45 42.24		0.685	•481	118	26:25	8.27 : 8.03	4536
										01-6-	-	027	17	8.67	4538
946	4.0	92 Herculisξ	17 53 52.768		+ 2.3302	+0023		+29 15 30.60		- 0.202			17		4530
947	4.0	67 Ophiuchi	55 38.190		3.0039	.0021		+ 2 56 10.27		0.396	•438		25:24	9'79 : 9'74 11'42 : 8'45	4540
948	5.4	Octantis		+1.02	35.738	• 369	096	-87 39 52.48		0.474	5.197	- ·131 - ·266		8.92	4550
949	5.8	Lacaille 7473			8.3886			-75 53 39.90		0.204	1.223	028	17 16:17	9'24:9'18	and the second sec
950	3.8	Aræθ	58 50.782	+ .011	4.6694	.0018	0015	-50 5 52.35	+ .26	0.150	000	- 020	10.1/	9 24 . 9 10	4565
1000	-77									1-15-1-1					

901. 2'9, 3'4 0'''5 245° 1905'5. 908. Var. 10 4'''5 60° 1898'4. 908. L, 4'8-5'3; P, 2<sup>d</sup>'05. 936. L, 4'4-5'0; P, 7<sup>d</sup>'01.

.

•

•

.

FROM OBSERVATIONS IN THE YEARS 1905-11.

No.	Mag.	Name.	Mean R.A. 1900'0.	$\mu_a \Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
-															
951 952	4'4 2'8	Pavonisπ 10 Sagittariiγ	h m s 17 58 57 159 17 59 23 023	s - ·024 + ·043	s + 5.7755 3.8526	+ • • 0028 • 0020	s + · 0024 - · 0046	-63 40 22.32 -30 25 32.94	+ 1.83	- 0.283 - 0.248	+ "842	- "191 - "194	16 16 : 17	9 <sup>.</sup> 95 9 <sup>.</sup> 43	4566 4568
953	3.7	72 Ophiuchi	17 59 23 023	+ '037	2.8433	·0018		+ 9 32 58.66	71	+ 0.310	•414	+ .082	17:18	9 43 8.73 : 8.65	4581
954	3.8	103 Herculis	3 38.475	- '002	2.3395	.00%1		+28 44 55.13	02	0.320	.341	+ .002	16:18	9'44 : 9'42	4584
955	4.2	Telescopii	3 48.402	+ .014	4'4531	.0002	0019	-45 58 18.36	+ .32	0.296	•649	032	16	8.60	4588
956	4.4	102 Herculis	18 4 28.872	+ .001		+ .0020		+20 47 54.64	+ .19	+ 0.322	+ .374	012	16	9.21	4590
957 958	4.0 5.5	13 Sagittarii	7 46·985 8 35·616	- ·003 - ·048	001	+ .0007	+.0003 +.0055	-21 5 6·47 -44 14 11·62	+ .05	0.676	·522 ·638	- ·005 + ·019	29 18:19	9'15:9'00 8'69	4604 4610
959	5.6	Lacaille 7608	8 42.088	+ .045	4°3782 5°0517	.0032	- '0047	-56 3 15.95	+ .25	0'735	.735	026	16.19	9.52	4611
960	8.1	Brisbane 6229	10 21.65		23.530	• 303		-86 15 57.90		0.906	3.426		29:37	11.39 : 10.58	·
961†	3.0	Sagittariipr. η	18 10 51.283	+ .092	+ 4.0592	0006	0112	-36 47 31.44	+ 1.37	+ 0.783	+ .589	- 167	20:22	8.30 : 8.21	4617
962†	4.3	Pavonispr: §		+ .008	5.5322	·0058	0010	-61 32 21.21	07	1.233	.804	+ .008	18:19	8.41 : 8.32	4625
963	2.2	19 Sagittariiδ	14 35.569	- '026	3.8412	0009	+'0028	-29 52 14.41	+ .34	1.540	.558	036	17	9.40	4628
964 965	3.3.	58 Serpentisη 20 Sagittarii		+ '343	3°1029 3°9823	+ ·0017 - ·0018	-·0376	- 2 55 36·24 -34 25 55·76	+ 6.32 + 1.11	0.711	°445 °578	- ·699	23:21 17:18	9'12:9'04 8'50:8'39	4638 4645
													1		
966 967	5°9 5°4	Bradley 2308 B.D. + 17° 3555	18 17 58.464 18 23.920	- ·010 - ·043	+ 2.2015	+ .0016		+23 14 4.40	- ·71 - ·07	+ 1.646	+ • 363	+ .075	16 16	9°42 9°60	4649 4651
968	4.0	109 Herculis		- '129	2.5556	+ .0021	1	+21 43 24.42	+ 2.47	1.437	372	261	22:23	9'34:9'45	4656
969	3.6	Telescopiia		+ .009	4.4514	- '0047	0010	-46 1 24.77	+ .46	1.657	•646	052	17:18	8.75	4657
970	6.0	Lacaille 7642	20 4.988	013	7.7250	·0359	+.0014	-74 1 38.98	+ 1.11	1.634	1.131	- '120	16:17	9'37 : 9'26	4658
971	4.2	Telescopii $\zeta$	18 21 7.840	135	+ 4.6247	0056	+.0147	-49 7 31.28	+ 2.36	+ 1.289	+ .673	257	16	9.19	4662
972	2.2	22 Sagittariiλ		+ .033	3.7028	.0015	0032	-25 28 39.03	+ 1.84	1.213	. 536	- • 191	22:23	9'51 : 9'64	4665
973	4·7 8·3	Scuti 2 H Lacaille 7442	23 29 890 23 58 40	- '002	3°4196 20°601	•0006		-14 37 46.87	+ .02	2.044	°495 2°986	- '008	17 30:38	8.65 11.43 : 10.72	4674
974 975	5.6	60 Serpentis	23 58 40	014		- ·517 + ·0004	+.0012	-85 39 49°08 - 2 3 0°76	+ .28	2°093 2°106	2 900 •452	- '031	30:30 16	9.00	 4678
976	4.6	Corouæ Anstθ		029	+ 4.2875	0057	+.0035	-42 23 4.14	+ .23	+.2.274	+ '620	027	21	8.41	4689
977	4'0	Seuti 3 H.		+ .013	3.2645	*0001	-'0015	- 8 18 53.40	+ 2.67	2'279	.471	- '317	21	8.42	4705
978	4'I	Pavonis		+ .021	7.0302	•0429	0022	-71 30 51.06	+ 1.48	2.579	1.014	155	16	9.53	4709
979	6.1	Lacaille 7780	31 40.297	- '012	4.5436	.0092	+.0013	-47 59 45.43	- '22	2.785	•655	+ .024	17	9.31	4710
980	5.9	Bradley 2333		+ .000	3.6495	•0027	-*0006	-23 35 25.11	+ .27	2.798	525	029	16	9.41	4718
981	6.1	Bradley 2335		+ .023	+ 3.5781	- '0022	0022		+ 1.48	+ 2.717	+ .214	- 154	16:17	9.63	4720
982 983	4·8 5·2	Scuti 4 Hλ Coronæ Austλ	36 47 936 36 55 383	003	3°2856 4°1190	'0012	+.0000	- 9 8 53.87 -38 25 10.94	$+ \cdot 04 + \cdot 52$	3.1201	•471 •591	- ·004 - ·059	22 16	9'19 : 9'29 8'73	4731 4732
984	3.2	27 Sagittarii	39 24 599		3.7496	- '0044	1	-27 5 36.98	+ .03	3 157	.538	003	22:21	9.11: 0.08	4732
985	4.3	110 Herculis	41 21.451	+ .014	2.5806				+ 3.28	3.254	• 368	344	17	9.54	4753
986	5.7	Coronæ Aust	18 41 37.532	024	+ 4.3350	- '0104	+.0026	-43 47 19.78	+ .12	+ 3.605	+ .620	016	16	9.18	4755
987	4.5	Scuti 6 H	41 52'122	+ .002	3.1833	- *0010	0007	- 4 51 18.13	+ .22	3.619	•455	023	16	9.55	4756
988	4'3	111 Herculis	42 36.322			+ .0008		+18 4 13.17	- 1.01	3.811	•378	+ .100	16	9.22	4761
989 990	4°3 6°5	Pavonisλ 30 Sagittarii	42 57°141 44 49°818		5.5713 3.6067	- °0294 °0040	-·0023	-62 18 7.38 -22 16 36.38	+ '11 + '21	3°724 3°872	•796 •514	- ·011	16 17:16	9'76 8'66 : 8'63	4762 4767
-	200		and the second second				1.000		1.7765	1					
991 992	6.7 var.	Lacaille 7881 10 Lyræpr. β		+ .014	+ 4.2437 2.2144	- '0103 + '0014	0012 +.0003	-41 49 33.18 +33 14 47.34	+ '22 + '07	+ 3.899	+ '605	- °023 - °007	16 16	· 9·43 9·86	4769 4776
993	2.0	34 Sagittariiσ	49 3.933	003	3.7218	- '0055	+.0002		+ .57	4.193	.528	066	19	8.70	4784
994	5.0	Tolescopiiλ	50 27.855	019	4.8094	°0202	+.0020	-53 4 10.21	- '13	4'393	·683	+ .014	17	9*57	4796
995	4.8	63 Serpentispr. θ	51 14'946	030	2.9826	.0002	+.0031	+ 4 4 24.27	- • 26	4.472	•422	+ .027	17:16	9*52:9'49	4802
996	3.2	37 Sagittariiξ		023	+ 3.2808		+.0023	-21 14 17.60	+ .18	+ 4.471	+ .207	018	22	10.09 : 10.12	4809
997	5'1	Coronæ Auste		+ .092	4.0512	-	0103	-37 14 16.81	+ .81	4.415	• 57 2	093	16	8.73	4810
998 999	4°2 3°2	13 Aquilæε 14 Lyræγ	55 5°001 55 12°128		2·7218 2·2436			+14 55 55 57 +32 33 8.14	+ '74 + '06	4°695 4°775	·383 ·316	- ·077 - ·007	20 16	9'70:9'64 9'09	4823
1000	and the second sec	38 Sagittariim. ζ			3.8199		- '0016	-30 I 23.29	- 00 •00	4 775	'538	.000	16	9'41	4832

961. 3'0, 10'3 3"'9 105° 1897'4. 962. 4'3, 10'0 3"'1 151° 1895'7. 992. L, 3'4-4'1; P, 12<sup>d</sup>'91. 1000. 3'4. 3'6; very close binary.

.

## CAPE FUNDAMENTAL CATALOGUE OF STARS FOR 1900.0,

No.	Mag.	Name.	Mean R. A. 1900'0.	$\mu_{a}\Delta E.$	Annual Variation 1900°0,	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0,	μ <sub>δ</sub> ΔΕ.	Aunual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900 <del>+</del> .	Boss No.
1001	5.3	Telescopii	h m s 18 58 25.003	s - '023	s + 4.7597	s - `0222	s +°0027	-52° 29 15'.82	+ 1.02	+ 4 <sup>"</sup> 937	+ ".670	- "118	17	8.63	4844
1002	5.2	Octantisσ Lacaille 7751σ	18 59 45 56 19 0 28 72	-1.53	102.437	38.819	+.100		+ .08	5.122	14.445	008	61 : 102	11.31:9.66	4854
1003	7.9	40 Sagittariiτ	0 41.819	+ '042	3.7484	·915 - ·0072	- '0045	-27 49 2'21	+ 2.45	5°229 4°987	2°463 525	- '260	35:43 17:18	9'35 : 9'44	4857
1005		17 Aquilæpr. 5	0 48.821	+ .000	2.7570	+ .0004		+13 42 51.74	+ 1.00	5.122	• 386	105	17	9.84 : 9.80	4858
1006	3.4	16 Aquilæλ		+ .016	+ 3.1842	0021		- 5 1 58.61	+ .86	+ 5.178	+ .446	000	18:16	9.61 : 9.57	4859
1000	4.2	Coronæ Austa	2 40.277	067	4.0873	- '0122		-38 3 37.47	+ .95	5'309	- 440	102	16.10	9.02	4868
1008		17 Lyræseq.	3 38.732	094	2.2680	+ .0015		+32 20 38.53	13	5.509		+ '013	16	9.85	4872
1009	5.3	18 Lyræt		+ .000	2.1400	+ .0015		+35 56 35.63	+ .00	5.497	· 298	- '006	16:17	9'79 : 9'81	4873
1010	3.0	41 Sagittariiπ	3 49.055	+ .004	3.2699	0029	0004	-21 10 58.24	+ .39	5'470	•498	040	17:18	9.60 : 9.66	4874
1011	5.6	Lacaille 7997 m	19 7 8.823	002	+ 6.0626	0624	+.0002	-66 50 0.52	- '02	+ 5'792	+ .844	+ .002	16	9.29	4882
1012		Lacaille 8029		054	4.3677	0184	+.000*	-45 21 44.55	+ .63	5.810	•607	07*	17	9.04	
1013	1	19 Lyræ		+ .000	2'2999	+ .0012		+31 6 59.04		5.848	.318	- '007	16	9.26	4885
1014		21 Aquilæ	8 40°192 9 24°616	+ .002	3.0248	- '0015 '0079		+ 2 7 24.50	+ .06 + .35	5'911	·419 ·510	- '006	16 21	9*89 9*92 : 9*90	4887 4891
	1				Ŭ					5.944		00			
1016		22 Aquilæd 43 Sagittariid	19 11 34.085 11 47.088	000	+ 2.9693	0012		+ 4 39 29.42	+ '13	+ 6.145	+ '409	- '014	16	9.36	4902
1017	-	Lacaille 8050		012	3.5125	·0062	-	-19 7 51.77 -51 45 8.20	+ '18	6°158 6°134	·484 ·648	- ·019	16 17	9.68 9.59	4903 4904
1019		Gilliss P.Z. 13504			40.647	7'010		-88 3 49.89		6.233	5.629		12:13	11'38:11'44	
1020	5.3	25 Aquilæ	13 7.357	+ .001	2.8160	.0004	0001	+11 24 53.87	- '10	6.299	.387	+ .011	20:19	9'34 : 9'16	4914
1021	4.0	Sagittarii	19 15 27.013	+ .001	+ 4.3214	0199	0001	-44 38 48.53	+ .17	+ 6.462	+ . 594	010	16	9.14	4929
1022	1.	Sagittariia			4.1644	·0169		-40 48 15.77	+ 1.18	6.480	.571	- '126	16	9.37	4936
1023	+ 5.7	Lacaille 8091br	19 46 294	+ .054	4.8295	.0337	0057	-54 31 28.89	14	6.853	•658	+ .012	16	9.23	4946
1024		31 Aquilæb			2.8611	.0015		+11 43 54.96		7.503	395	1 -	16	9.47	4950
1025		30 Aquilæδ		163	3.0223	.0019	+.0169	+ 2 54 55 43	- '74	6.971	.414	+ .072	17:16	9.62	4953
1026		Lacaille 8107			1 0 1 - 0 -	- '0112			+ .22	+ 6.853		022	16	10.03	4955
1027		Bradley 2459			2.4813	+ .0012		+24 43 49 43		6.331	.334	631	16	10.00	4961
1028 1029		5 Vulpeculæ 6 Vulpeculæ			2.6186 2.4959	·0005 + ·0010		+195356.22 +242743.44	+ '37	6·970 7·115	·354 ·335	039	16 17	9°45 8°95	4965 4976
1029		36 Aquilæe			3.1381	- '0031			1.	7.288	335 '423	013	16	9.49	4970
		Lacaille 8129							_						
1031		6 Cygniрг. β				- '0233 + '0010		-45 29 1.08 +27 44 58.22		+ 7.327	+ .584	- '033 - '009	16 17	9.69 9.14	4984 4986
1033		Telescopii						-48 18 53.64		7 394	•600		16	9.46	4990
1034		8 Cygni		+ .002	2.2288			+34 14 24 62		7.211	• 298		16:17	9'98 : 9'99	4992
1035	4.8	38 Aquilæ	29 12.399	- '137	2.9312	- '0012	+.0143	+ 7 9 57.80	+ 1.45	7.455	*394	- 152	24:23	9'56 : 9'53	4995
1036	5.1	39 Aquilæ	19 31 30'721	002	+ 3.2292	0045	+.0002	- 7 14 59.85	+ .02	+ 7.792	+ .430	002	16	8.76	5003
1037		4 Sagittæ				.0001	+.0010	+16 14 16.86	- 12	+	.361		17	9.39	5010
1038		44 Aquilæσ			2.9614			+ 5 10 11.05					16	9'02	5018
1039		54 Sagittarii						-16 31 21.66		8.019				9'73:9'43	5019
1040	4'5	6 Sagittæ В			2.6939	+ .0001	+.0001	+17 14 38.91	+ .35	8.160			16	9°27	5027
1041		55 Sagittariie						-16 21 30.51					17:16	9'23:9'25	5028
1042		Lacaille 8094 Lacaille 8156	0. 0	- '0I - '0I2	6:9902	1464		-81 36 0.60			1·499 ·924		52:141 16:17	11·39 : 8·51 9·68 : 9·65	5030
1043		10 Vulpeculæ	· 37 53 525 . 39 33 398			1		+25 31 57.10	-		326		16	9.50	5034 5039
1045		Lacaille 8211	39 38.474					-32 8 59'30			1		16	9.64	5040
1046		Telescopii		10.00				-56 36 11.61					17:18	9.99	5041
1040		56 Sagittarii						-20 0 6.78						9'48:9'52	5041
1048		50 Aquilæ		-				+10 22 9.92						9.01	5047
1049		7 Sagittæ			2.6746	+ .0001	+0001	+18 17 14.64	08	8.713	• 348			9.41	5052
1050	$\begin{array}{c} 1050 \\ 5.6 \\ 1050 \\ 5.6 \\ 1050 \\ 1050 \\ 5.6 \\ 1050 $														
	$\begin{array}{cccccccccccccccccccccccccccccccccccc$														

FROM OBSERVATIONS IN THE YEARS 1905-11.

.

No.	Mag.	. Name,	Mean R.A. 1900'0.	μ <sub>α</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0,	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900 0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
1051 1052 1053	var.	<ul> <li>53 Aquilæα</li> <li>Lacaille 8226η</li> <li>55 Aquilæη</li> </ul>	47 22.758	s - `339 - `028 - `005	s + 2 <sup>.</sup> 9274 5 <sup>.</sup> 2717 3 <sup>.</sup> 0573	s - 0018 0638 0032	+.0002		$-3^{3}57$ -15 +08	+ 9 <sup>:</sup> 317 8·956 9·043.	+ <sup>":</sup> 383 .684 .394	+ <sup>"</sup> 380 + ·015 - ·009	16 16:17 16	9°40 9°87 : 9°79 9°17	5062 5064 5071
1054 1055 1056 1057	4°1 4°0 3°8 5°4	Sagittarii	49 1.985	$ - \cdot 007 - \cdot 148 - \cdot 023 - \cdot 012 $	4.1494 7.0172 + 2.9469 2.8406	·0249 ·1646 - ·0015 ·0011		$\begin{array}{rrrrr} -42 & 7 & 50.79 \\ -73 & 10 & 29.10 \\ + 6 & 9 & 19.62 \\ +11 & 9 & 29.12 \end{array}$	$ \begin{array}{r} - & \cdot 48 \\ + & 1 \cdot 21 \\ + & 4 \cdot 81 \\ - & \cdot 08 \end{array} $	9.181 9.049 + 8.805 9.381	·535 ·908 + ·377 ·362	$+ \cdot 052$ $- \cdot 132$ $- \cdot 483$ $+ \cdot 008$	18:19 16:17 23:22 17:18	9'18 : 9'22 9'16 9'86 : 9'95 9'50 : 9'44	5078 5084 5093 5099
1058 1059 1060 1061	5·1 4·4 3·7 5·8	61       Sagittarii $g$ Sagittarii $\theta^1$ 12       Sagitta $\gamma$ 63       Sagittarii $\gamma$	JH J-	- ·004 - ·012 - ·040 - ·020	3.4055 3.9144 2.6674 + 3.3639	$- \cdot 0084$ $- \cdot 0195$ $+ \cdot 0002$ $- \cdot 0081$			$ + \cdot 83 + \cdot 42 - \cdot 15 - \cdot 18 $	9°343 9°462 9°605 + 9°767	·434 ·499 ·338 + ·425	- ·090 - ·044 + ·016 + ·019	16 : 17 17 : 18 20 : 22 16 : 18	9'19 9'45 9'51 : 9'47 9'26 : 9'23	5101 5108 5118 5128
1062 1063 1064 1065	4.6 4.9 6.9 3.6	62 Sagittarii	57 48.775 58 57.017	$ \begin{array}{r} - \cdot 026 \\ - \cdot 039 \\ + \cdot 025 \\ - 1 \cdot 882 \\ \end{array} $	3.6955 2.4698 3.5606 5.9258	- ·0148 + ·0012 - ·0121 ·0933		$\begin{array}{r} -27 59 16.19 \\ +27 28 37.40 \\ -22 52 34.82 \\ -66 26 23.71 \end{array}$	$ \begin{array}{r} - & \cdot 10 \\ - & \cdot 06 \\ - & \cdot 23 \\ + 11 \cdot 21 \end{array} $	9.768 9.800 9.882 8.795	•310 •448	+ .010 + .006 + .025 -1.146	17 16 18 : 20 16	9.67 9.95 9.14 : 9.02 9.78	5129 5132 5135 5138
1066 1067 1068 1069 1070	5.9 5.1 6.3 3.2 6.0	<ul> <li>63 Aquilæτ</li> <li>Telescopiiξ</li> <li>Lacaille 8202θ</li> <li>65 Aquilæθ</li> <li>20 Vulpeculæ</li> </ul>	19       59       43.575         20       3       33.         6       8.784	$ \begin{array}{r} - \cdot 008 \\ + \cdot 026 \\ \dots \\ - \cdot 020 \\ + \cdot 003 \\ \end{array} $	+ 2.9309 4.6160 13.327 3.0967 2.5146	- °0020 •0443 1°060 - °0042 + °0012	-·0027 -·001 +·0021		$ \begin{array}{r} - & \cdot 20 \\ + & \cdot 01 \\ + & \cdot 01 \\ - & \cdot 03 \\ + & \cdot 15 \\ \end{array} $	+ 9.988 10.002 10.290 10.488 10.594	·579 1·663	$+ \cdot 021$ $- \cdot 001$ $- \cdot 002$ $+ \cdot 003$ $- \cdot 016$	16 16:17 4 30:24 17	9·29 9·60 7·28 9'50 : 9'02 9·19	5143 5147 N1306 5171 5178
1071 1072 1073 1074	5°7 5°0 4°5 6°1	66 Aquilæρ           67 Aquilæρ           5 Capricorniα <sup>1</sup> 4 Capricorni	20 8 4.099 9 39.021 12 6.390	- '011 - '033 - '009 - '021	+ 3.0994 2.7759 3.3283 3.5300	- ·0043 ·0005 ·0085 - ·0128	+·0012 +·0036	- 1 18 33°21 +14 53 34°57 -12 49 2°61	$+ \cdot 22$ $- \cdot 47$ $- \cdot 06$ $+ \cdot 31$	+10.604 10.796 10.932 10.895	+ · 378 · 337 · 402 · 427	$ \begin{array}{r} - \cdot 024 \\ + \cdot 051 \\ + \cdot 006 \\ - \cdot 034 \\ \end{array} $	17 16 17 17:16 18	9.09 9.19 9.21 9.24	5179 5182 5197 5198
1075 1076 1077 1078	5.2	<ul> <li>4 Supression</li> <li>6 Capricorniα<sup>2</sup> Lacaille 8400</li> <li>9 Capricorniβ</li> </ul>	12 30'319 20 12 30'494 14 25'102	- :013	2.5668 + 3.3318 4.3721 3.3742	+ ·0011 - ·0085 ·0408	+.0013	+24 21 46·45 -12 51 17·73	+ .19 05 + 2.32 01	10.932	+ ·403 ·521 ·404	$- \cdot 020$ + $\cdot 005$ $- \cdot 258$ + $\cdot 001$	16 22 : 20 17 20 : 18	9.62 10.13 : 10.02 8.99 9.48 : 9.22	5201 5202 5209 5216
1079 1080 1081 1082	5·8 1·8 7·1 6·2	Sagittarii	15 40°290 17 44°350 20 18 48°31	- •055 - •005	3 3742 4 0897 4 7738 +15 054 3 6810	·0297 ·0595 - 1.640 ·0177	+·0059 +·0005 +·031	$\begin{array}{r} 15 & 5 & 50 & 45 \\ -42 & 21 & 53 & 54 \\ -57 & 3 & 20 & 55 \\ -84 & 44 & 49 & 40 \\ -28 & 59 & 15 & 50 \end{array}$	+ .97 + .79 29 05	11.082 11.250	404 •491 •569 +1•803 •436	- ·104 - ·086 + ·033 + ·005	16 17 49 : 165 16	9°29 9°17	5217 5223 N 1326 5232
1082 1083 1084 1085 1086		69 Aquila 41 Cygni	24 25 <sup>.</sup> 493 25 18 <sup>.</sup> 571 27 2 <sup>.</sup> 884	- ·038 - ·006 - ·011	3·1373 2·4505 4·1389	- ·0054 + ·0020 - ·0349	+ · 0042 + · 0007 + · 0013	- 3 13 6.02 +30 2 5.02 -44 51 18.50	+ ·19 + ·04 + ·36	11.792 11.872 11.957	·365 ·283 ·478	$- \cdot 021$ $- \cdot 004$ $- \cdot 041$ $- \cdot 168$	16 16 17:18 16	9 20 9 07 9 06 8 84 : 8 78 9 49	5252 5254 5255 5266 5268
1087 1087 1088 1089 1090	4 · I 5 · d 6 · 3 3 · I	2 Delphini	28 26 154 29 12 484 29 42 495 30 32 204	- · 067 - · 006 - · 063 - · 453 - · 040	+ 5.0029 2.8665 5.0678 7.5576 4.2369	1	+.0067	+10 57 47 37	+ 1.59 + .25 + .59 + .07 61	12.086 12.086 12.175 12.302	+ ·579 ·328 ·583 ·876 ·484	$- \cdot 026$ $- \cdot 063$ $- \cdot 008$ $+ \cdot 061$	28 : 20 16 45 : 46 16 : 17	9 49 9 98 : 978 9 43 8 38 10 13 : 9 92	5272 5274 5277 5281
1091 1092 1093 1094		4 Delphiniζ 6 Delphini	32 51.668 34 3.374	- ·025 - ·067 - ·038 - ·030	+ 2.8047 2.8131 2.6782 3.7751	$ \begin{array}{r} - & \cdot 0005 \\ - & \cdot 0004 \\ + & \cdot 0010 \\ - & \cdot 0229 \end{array} $	+·0026 +·0074 +·0041	+14 19 45·12 +14 14 49·46	- ·04 + ·33 + ·01 - ·07	+12°252 12°364 12°482 12°490	+ ·319 ·318 ·301 ·426	+ ·004 - ·037 - ·001 + ·007	16 23:21 16 16	9°59 9°01 : 8°88 9°29 9°83	5282 5291 5301 5302
1095 1096 1097 1098	5°3 5°4 3°9 3°4	<ul> <li>7 Delphiniκ</li> <li>15 Capricorniν</li> <li>9 Delphiniα</li> <li>Pavonisβ</li> </ul>	34 16.577 20 34 21.491 34 59.653 35 56.977	- ·220 + ·020 - ·043 + ·072	2.9144 + 3.4197 2.7865 5.4607	·0016 - ·0122 ·0001 ·1163	+·0213 -·0020 +·0044	+ 9 44 1.72 -18 29 27.18 +15 33 32.93 -66 33 44.27	$ \begin{array}{r} - & \cdot 12 \\ + & \cdot 21 \\ + & \cdot 08 \\ - & \cdot 12 \\ \end{array} $	12.510 +12.483 12.539 12.625	+ .384	$+ \cdot 012$ - $\cdot 021$ - $\cdot 008$ + $\cdot 013$	16 18:17 16 18	10°35 10°03 9°67 9°14	5304 5306 5310 5315
1099 1100	4.7	Indiη 11 Delphiniδ		141	4°4271 2°8007	·0508	+.0153	-52 16 41.81 + 14 42 56.15	+ .42	12.615 12.754	·497 •308	- ·048 - ·051	15.17 17:18	9 <sup>.</sup> 23 : 8 <sup>.</sup> 85 8 <sup>.</sup> 74 : 8 <sup>.</sup> 78	5318 5323

1053. L, 3'6-4'2; P, 7<sup>d</sup>'18. 1076. BC=10'6 7"'8 152° 1092. 4'0, 5'3; very'close binary.

×.

•

•

## CAPE FUNDAMENTAL CATALOGUE OF STARS FOR 1900.0,

.

No.	Mag.	Name.	Mean R.A. 1900°0.	$\mu_{\alpha}\Delta E.$	Annual Variation 1900 '0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900 <b>+</b> .	Boss No.
1101 1102	4.2	16 Capricorniψ Microscopii	h m s 20 40 10'540 41 42'681	* + ·038 - ·142	8 + 3.5590 4.0826	- <sup>8</sup> - '0167 - '0362		-25 37 50.75 -44 21 11.55	+ 1 <sup>"</sup> .44 + .87	+12.739	+ "391	<sup>"159</sup>	17 : 18 16	9'13 : 9'03 9'13 : 8'97	5328 5332
1103 1104 1105	4·4 2·5 3·8	12 Delphiniγ 53 Cygniε 2 Aquariiε	42 1°112 42 10°166 42 15°856	+ ·022 - ·292 - ·019	2°7832 .2°4265 3°2507	+ .0028	+.0288	+15 45 47.69 +33 35 47.36 - 9 51 43.45	+ 2.00 - 3.27 + .34	12.817 13.352 13.003	• 302 • 267 • 354	$- \cdot 204 + \cdot 322 - \cdot 034$	16 16 18 : 17	9°78 10°15 10°04 : 10°00	5335 5336 5337
1106 1107 1108	4·6 5·3	3 Aquarii Indi	20 42 27.721 44 16.409 45 51.313	+ ·004 - ·004 + ·006	+ 3.1676 4.3616 3.5885	- ·0065 ·0512 ·0184	-·0004 +·0005	- 5 23 3 <sup>8</sup> .77 -51 58 49.70 -27 17 36.03	$+ \cdot 39$ $+ \cdot 32$ $+ \cdot 13$	+13.011 13.133 13.260	+ '345 '474 '386	- ·039 - ·037 - ·014	16 17 : 18 16 : 17	10°05 8'77 9'49 : 9'45	5338 5354 5363
1103 1109 1110	4·1 3·6 5·6	Indi	46 59·868 47 9·945	- '014 - '037	4°7223 3°9179	•0734 •0308	+·0015 +·0040	-58 49 53.15 -40 11 3.83	+ ·24 + ·91	13°323 13°264	• 508 • 420	— •026 — •096	17:18 16:18	9'43 : 9'37 9'37 : 9'49	5367 5369
1111 1112 1113 1114	4·8 6·1 5·3 5·3	6 Aquariiμ 19 Capricorniμ 32 Vulpeculæα	49 8·854 50,17·858	$- \cdot 025$ + $\cdot 035$ + $\cdot 006$ + $\cdot 024$	+ 3°2390 3°3956 2°5554 7°4332	$ \begin{array}{r} - & \cdot 0083 \\ - & \cdot 0128 \\ + & \cdot 0026 \\ - & \cdot 3501 \end{array} $	+ 0025 - 0038 - 0007 - 0028	- 9 21 31.48 -18 18 8.28 +27 40 37.75 -77 24 21.52	+ ·17 + ·02	+13.331 13.469 13.561 13.349	+ ·346 ·360 ·268 ·784	$- \cdot 035$ $- \cdot 019$ $- \cdot 002$ $- \cdot 362$	23:17 16:17 23:20 33:34	10'02 : 9'92 9'17 : 9'15 9'09 : 8'82 8'48 : 8'47	5371 5374 5379 5390
1115 1116 1117	6·0 7·0	Lacaille 8624 C.G.A. 28663γ Microscopiiγ	53 14.650 20 53 43.67	+ .086	4.2920 +15.699 3.6911	·0519 - 2·472 ·0235		-51 39 24 95 -85 36 17 00	- 1°08	13.878 +13.782 13.877	·448 +1.656 ·383	+ · · 127  + · 004	19 56:69 17	8.74 : 8.47 11.40 : 10.53 9.06	5391  5402
1118 1119 1120	5°5 7°4	Microscopiiζ C.G.A. 28714 22 Capricorniη	56 34.681 57 18.07	+ '014	3.8478 16.919 3.4199	•0302 3•036 •0142	0017	-39 1 19·82 -86 3 0·95	+ .99	13.842 14.007 14.052	· 396 1.759 · 348	- ·120  - ·043	19 58 : 68 17	8·23 11·39 : 10 <sup>.</sup> 68 9 <sup>.</sup> 22	5411
1121 1122 1123	7.0	23 Capricorni	. 0 59.161	031	+ 3.3779 4.1714 3.5165	- ·0128 ·0474 ·0178	+.0036	-49 20 25.23	+ .26	+14.129 14.206 14.203	+ •342 •422 •354	- ·066 - ·030 - ·051	22:17 18 18	9'55 : 9'33 8.66 8'77	5427 5429 5430
1124 1125 1126	4.6	Pavonis	4 8.933	052	5.7043 3.2721 + 2.9178	·1704 ·0098	1.		+ .11	14·384 14·417 +14·349	·572 ·326 + ·288	$- \cdot 034$ $- \cdot 013$ $- \cdot 161$	17:18 23 17	9.04 : 8.89 9.03 : 8.62 9.26	5439 5441 5443
1127 1128 1129	5.7 6.0 3.3	3 Piscis Aust Lacaille 8727 64 Cygni	. 7 21.728 8 37.501 8 40.762	- ·065 - ·009	3.5661 4.3094 2.5515	·0200 - ·0588 + ·0040	+.0071 +.0009 0002	-28 I 39.34 -53 40 36.58 +29 48 59.40	+ 1.27 + .41 + .57	14·485 14·657 14·643	· 350 · 421 · 247	- ·138 - ·043 - ·059	16 16 16	9°21 9°55 9°60	5448 5451 5452
1130 1131 1132	4°0 7°0	Lacaille 8551 8 Equulei	21 10 49 580 11 3 981	+ .031	4.1073	- 2°138 - °0027 °0476	0033	$ \begin{array}{r} -85 & 14 & 17 \cdot 64 \\ + 4 & 50 & 2 \cdot 44 \\ -49 & 8 & 2 \cdot 11 \end{array} $	+ .86	14.813 +14.742 14.757	+ •289		58:71 18:16 16	11'40 : 10'72 9'99 : 9'90 9'32	5461 5463
1133 1134 1135	4.4	Microscopii	13 48.325	015		+ .0020	+.0019	$\begin{array}{r} -32 & 35 & 26 \cdot 03 \\ +34 & 28 & 37 \cdot 07 \\ -41 & 13 & 56 \cdot 22 \end{array}$	+ .19	14.856 14.982 15.035	·350 ·232 ·366	- '021	16 18 : 17 19	9°25 9°26 8°28	5464 5471 5473
1136 1137 1138 1139	4°3 † 6°1	32 Capricorni 1 Pegasi Microscopiim. 6 Pavonis	. 17 27·769 18 2·448	- ·065 - ·020	2·7734 3·8405	+ .0019	+.0022	$ \begin{array}{r} -17 & 15 & 38.08 \\ +19 & 22 & 36.20 \\ -41 & 26 & 6.87 \\ -65 & 48 & 59.53 \\ \end{array} $	- ·52 - ·03	+15°174 . 15°271 15°249 16°065			27:20 17:18 17 16:17	9.89:9.53 9.06:8.97 9.16 9.67:9.48	5484 5489 5492
1140 1141	6·4 † 5·8	Indi	19 7.620 21 20 36.812	+ .001	4·3089 + 3·8645	·0642 - ·0373	- · 0001	-55 5 31.93 -42 58 50.98	38	15·349 +15·398	·399 + ·354	+ .041	16 18 : 19	9°24 8°68 : 8°59	5493 5497 5506
1142 1143 1144 1145	4·6 2·9	34 Capricorni	23 1.482	- ·086	3.4325 3.4276 3.1608 3.9063	•0166 •0163 •0071 •0413	+.0002	-22 14 34.09 - 6 0 40.49	+ ·07 + ·07	15.434 15.517 15.698 15.726		- · 008	21 : 23 19 21 : 23 18	9'77 : 9'76 9'09 9'54 : 9'75 8'77	5507 5513 5527 5530
1146 1147 1148	6·5 3·7	Lacaille 8842 Octantis	. 21 30 4·180 30 21·968	- ·021 - ·122	+ 4.8407 6.8523 3.1969	- ·1158 ·3826 ·0082	+.0025	-65 16 18·22 -77 50 3·64	+ .03	+15.905 15.694 16.009	+ •423	- ·003 - ·230	19:20 37:38 16:18	8·44 : 8·37 8·73 : 8·76 9·55 : 9·62	5541 5544
1149	7.7	Lacaille 8751	33 6.97	- ·072	3°1969 11°251 13°149	1.557 2.314	+ 0075	$ \begin{array}{r} - 8 & 18 & 10^{\circ} 43 \\ - 84 & 25 & 10^{\circ} 82 \\ - 85 & 29 & 46^{\circ} 51 \end{array} $		16.009	•974		51 : 63 53 : 65	9 55 : 9 62 11'42 : 10'65 11'42 : 10'68	
					1117. 1126.	1 Piscis À 4'7, 11	nstralis in 2 <sup>''•</sup> 2	Auwers' Bradle 272° 1901.6	y.						

 1126.
 4.7, 11
 2"2
 272°
 1901'6.

 1133.
 4 Piscis Australis in Auwers' Bradley.
 1138.
 6'4, 7'6
 1"0
 292°
 1900'8.

 1141.
 5'8, 8'8
 2"'9
 146°
 1900'6.
 1900'6.

## FROM OBSERVATIONS IN THE YEARS 1905-11.

•

No.	Mag.	Name.	Mean R. A. 1900'0.	$\mu_{a}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900°0.	μ <sub>δ</sub> ΔΕ.	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
1151 1152 1153 1154	5·4 4·8 6·7	40 Capricorni         γ           41 Capricorni         γ           43 Capricorni         κ           Octantis         B	h m a 21 34 33.216 36 19.233 37 4.642 37 41.30	s - ·117 - ·057 - ·093 - ·25	s + 3.3294 3.4235 3.3560 68.391	- <sup>8</sup> •0131 •0174 •0145 88•544		-17 6 50.98 -23 42 55.75 -19 19 19.83 -89 19 3.92	+ <sup>"18</sup> + <sup>.79</sup> + <sup>.07</sup> + <sup>.41</sup>	+16.123 16.142 16.266 16.263	+ <sup></sup> 282 · 286 · 279 5 • 799	- <sup>".</sup> 021 093 007 041	19:20 20 17 67:115		5562 5568 5570 5576
1155 1156 1157 1158 1159 1160	2.4 4.2 5.6 2.8	9 Piscis Aust	38 59 562 21 39 16 514 40 6 996 41 9 212 41 31 552 41 45 794	- ·024 - ·016 - ·023 - ·014 - ·170 - ·133	3.5855 + 2.9464 2.7145 3.2333 3.3163 3.9215	·0259 - ·0005 + ·0047 - ·0100 ·0125 ·0461	+·0024 +·0016		$\begin{array}{r} + & .72 \\ + & .01 \\ - & .02 \\ + & .10 \\ + & 2.80 \\ + & 2.84 \end{array}$	16.285 +16.384 16.429 16.468 16.202 16.202	•295 + •240 •220 •261 •269 •318	$ - \cdot 086  - \cdot 001  + \cdot 002  - \cdot 011  - \cdot 295  - \cdot 307 $	20 17 16 18:19 21:20 17	8·37 9·49 9·42 8·88 : 8·77 9·51 : 9·50 9·24	5582 5584 5592 5596 5600 5601
1161 1162 1163 1164 1165	5.6 5.3 5.3 3.1	Indiο 14 Pegasiμ 51 Capricorniμ Gruisγ 16 Pegasi	THE REAL PROPERTY.	+ ·080 - ·018 - ·195 - ·080 - ·001	+ 5°1500 2°6518 3°2754 3°6474 2°7272	- ·1651 + ·0064 - ·0112	0084 + .0020 + .0211 + .0093	-70 5 40.69 +29 42 30.20 -14 1 21.12	+ 204 + 25 - 08 + 15 - 01	+16.538 16.662 16.814 16.790 16.838	+ ·417 ·207 ·255 ·283	·000 - ·027 + ·009 - ·017 + ·001	16 19 16 19 16	9 57 9 23 9 24 8 63 9 77	5607 5617 5623 5624 5627
1166 1167 1168 1169 1170	8·1 4·6 6·6 4·9	Lacaille 8738δ Indiδ Mayer 939 Indi		 055 009 -4.206 001	+16.730 4.1143 3.3523 4.6243 3.0709	- 4.687 .0660 .0160	 + · 0064 + · 0010 + · 4818 + · 0001	-21 39 36·37 -57 12 10·11	$ \begin{array}{r}\\ + 20\\ + 03\\ +22 \cdot 13\\ + 06 \end{array} $	+16.845 16.936 17.050 14.579 17.176	+1·315 ·313 ·250 ·387 ·223	 - '024 - '004 -2'591 - '006	41 : 49 19 : 21 19 18 : 21 18	11.45 : 10.78 8.59 : <b>8</b> .46 8.53 8.73 : 8.54 9.25	 5635 5645 5654 5655
1171 1172 1173 1174 1175	4.6 5.1	20 Pegasi 31 Aquariiο Gruisλ 22 Pegasiν 34 Aquariiα	21 58 8.554	$ \begin{array}{r} - \cdot 033 \\ - \cdot 008 \\ + \cdot 023 \\ - \cdot 068 \\ - \cdot 009 \end{array} $	+ 2.9218 3.1045 3.6304 3.0265 3.0826	+ ·0014 - ·0050 ·0335 ·0018 ·0041	$+ \cdot 0009$ $- \cdot 0028$ $+ \cdot 0073$	-40 I 34·29 + 4 34 II·6I	$ \begin{array}{r} + & \cdot 51 \\ + & \cdot 10 \\ + & 1 \cdot 00 \\ - & \cdot 86 \\ + & \cdot 06 \end{array} $	+17.137 17.268 17.245 17.481 17.383	+ ·212 ·222 ·257 ·213 ·216	$ - \cdot 056 - \cdot 011 - \cdot 120 + \cdot 092 - \cdot 006 $	19 18 20 17 21 : 24	9'14 9'24 8'34 9'37 9'51	5658 5663 5672 5674 5676
1176 1177 1178 1179 1180	1.7 3.9 4.6	33 Aquarii	1 56.115 2 21.481 2 33.127	- ·025 - ·112 - ·193 - ·051 + ·038	+ 3.2445 3.8011 2.7903 3.5114 2.6547		+·0220 +·0057	-47 26 44.06 +24 51 23.94 -33 28 35.73	$ \begin{array}{r} + & \cdot 60 \\ + & 1 \cdot 56 \\ - & \cdot 16 \\ + & \cdot 36 \\ + & \cdot 60 \\ \end{array} $	+17.346 17.283 17.481 17.432 17.495	+ ·228 ·266 ·194 ·244 ·179	- ·060 - ·162 + ·018 - ·040 - ·072	17:18 17:18 18 19:18 19:18 19	9'97 : 10'00 9'62 8'78 8'96 : 9'01 8'34	5680 5684 5688 5689 5701
1181 1182 1183 1184 1189	6·5 6·7	26 Pegasiθ 29 Pegasiπ 28 Pegasiπ Lacaille 9061 16 Piscis Austλ	5 32.692 5 46.564 8 32.555		+ 3.0266 2.6606 2.8316 3.6809 3.4097	$\begin{array}{r} + & \cdot 0089 \\ + & \cdot 0048 \\ - & \cdot 0359 \end{array}$	0010 0021 +.0489	+ 5 42 21.25 +32 41 14.61 +20 29 11.13 -41 51 26.24 -28 15 45.40		+17.616 17.575 17.595 16.940 17.724		+ •034 - •023 - •013 - •782 - •003	18:20 17 17:18 20:21 18:20	9'50 : 9'20 9'24 9'49 8'44 : 8'51 8'36 : 8'42	5703 5709 5710 5725 5726
1186 1187 1188 1189 1190	4°4 2°9 5°6	Gruisμ1         43 Aquariiθ         Toucaniα         Lacaille 9076         Octantis (C.)ν	11 33 551 11 39 125 11 42 828 12 34 83	074	+ 3.6311 3.1684 4.1482 3.9588 12.836		+ · 0043 + · 0074 - · 0111 + · 0459 - · 040	60 45 29.19 54 6 37.58 86 28 33.33	$ \begin{array}{r} - & \cdot 28 \\ + & \cdot 19 \\ + & \cdot 30 \\ + & 6 \cdot 44 \\ - & \cdot 60 \\ \end{array} $	+17.796 17.825 17.817 17.169 17.952	·203 ·267 ·258 ·835	$\begin{array}{r} + \cdot 031 \\ - \cdot 019 \\ - \cdot 031 \\ - \cdot 681 \\ + \cdot 067 \end{array}$		9'11 : 9'00 9'95 : 9'77 9'75 : 9'74 9'46 11'45 : 8'97	5733 5744 5747 5748 5750
1191 1192 1193 1194 1192	5.6 5.4 3.9 5.1	46 Aquariiρ Indiν 47 Aquariiν 48 Aquariiγ 31 Pegasi	16 5.058 16 5.377 16 29.601 16 35.733	- <sup>•</sup> 073 - <sup>•</sup> 003	+ 3°1595 5°2441 3°3086 3°0998 2°9519	2000	+.0082	- 1 53 28.64 +11 42 4.82	+ ·80 - ·08 - ·05	+17 ·972 17 · 325 17 ·934 18 · 046 18 · 046	.180	$ \begin{array}{r} - \cdot 005 \\ - \cdot 694 \\ - \cdot 087 \\ + \cdot 009 \\ + \cdot 005 \\ \end{array} $	16 15:16 18 17:19 16:17	9°23 10°11 : 10°05 9°17 8°95 : 8°90 9°82 : 9°61	5759 5761 5762
1190 1192 1198 1199 1200	+ 6.0 4.6 5.7	32 Pegasi       Gruis	16 59.765 20 10.254	- ·006 - ·034	+ 2.7654 3.7051 3.0645 3.5320 3.6034	+ ·0083 - ·0433 ·0027 ·0324 ·0386	+·0223 +·0007 +·0040	$\begin{array}{r} +27 \ 49 \ 36^{\circ}47 \\ -46 \ 25 \ 54^{\circ}50 \\ + \ 0 \ 52 \ 11^{\circ}33 \\ -39 \ 38 \ 17^{\circ}34 \\ -44 \ 0 \ 23^{\circ}25 \end{array}$	$\begin{array}{r} + & \cdot 02 \\ + & \cdot 57 \\ - & \cdot 03 \\ + & 1 \cdot 43 \\ - & \cdot 01 \end{array}$	+18.043 17.995 18.178 18.101 18.289	·228 ·181 ·204	$ \begin{array}{r} - \cdot 002 \\ - \cdot 061 \\ + \cdot 003 \\ - \cdot 170 \\ + \cdot 001 \\ \end{array} $	16 16:18 19 20 20	10°38 9°63 : 9°41 9°24 8°44 8°37	5763 5765 5777 5789 5791

1157. 4'7, 5'4; very close binary. 1197. 6'0, 12'5 4"'7 208° 1900'8.

C. F. C., 1900.

.

25

.

D

.

CAPE FUNDAMENTAL CATALOGUE OF STARS FOR 1900.0,

No.	Mag.	Name.	Mcan R.A. 1900'0.	$\mu_{a}\Delta E.$	Annual Variation 1900°0.	Sec. Var. 1900 °O.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0,	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
I 20I I 202	4°9 5°6	57 Aquariiσ 38 Pegasiσ	lı 111 8 22 25 21°377 25 27°327	8 •000 — •024	s + 3 <sup>.</sup> 1784 2 <sup>.</sup> 7392	- <sup>8</sup> .0087 + '0107	s •0000 +•0025	$-11^{\circ} 11^{\circ} 23^{\circ} 42$ $+32^{\circ} 3 38^{\circ} 57$	+ ":29 + :15	+18.332	+ "178	- "030 - °016	21 : 18 16	9*86 : 9*65 9* <b>5</b> 0	5803 5806
1203 1204	4°4 5°2	17 Piscis Austβ Toucaniν	25 49 <sup>•</sup> 343 26 14 <sup>•</sup> 621	- <sup>.</sup> 043 - <sup>.</sup> 023	3°4218 4°0961	- •0247 •0909	+.0046	-32 51 32°06 -62 29 44°94	+ .17 + .30	18·360 18·359	· 192 · 230	- ·018 - ·034	16 . '22	9°34 8°70	5808 5811
1205 1206	5°5 4°1	59 Aquariiν 62 Aquariiη	29 13.639	- ·129	3.2880	·0150 - ·0030	+.0122	-21 13 15.03 -0 37 59.25	+ 1.23	18·347 +18·474	·178 + ·164	- ·148	20 25 : 26	8·31 8·88 : 8·58	5819 5824
1207 1208	6°5 5°4	Lacaille 9181	30 39°044 32 34°646	- ·025 + ·048	3.2182	·0339	+.0028	-41 5 55.42 - 4 44 39.28	+ .77	18.458	·187 ·160	085 115	18 16	9.08 9.30	5828 5835
1209 1210	5.8 4.2	Lacaille 9197 18 Piscis Aust	33 12.569	- ·007 - ·017	3.4014 3.3266	·0251 ·0196	+ · 0008	-33 36 6.14 -27 33 54.58	- ·32 + ·02	18.664 18.686	·175 ·168	+ .038	20 16	8·37 9·47	5836 5849
1211 1212	4'3	Octantis		+ '34	+ 6.4158	6245	0300	-81 54 20.84	06	+18.716	+ · 326	+ .002	29 : 3I 18 : 17	11 <b>'</b> 49 : 11'48 9'86 : 9'82	5850 5853
1213	3.2	Gruis	36 42.046	118	2°9909 3°6017	+ .0024 0435	+.0122	-47 24 27 08	+ .20	. 18.716	•179	021	16	9'40	5854
1214 1215	6·8 3·0	67 Aquarii 44 Pegasiη	38 1.008 38 18.816	- '011 - '007	3°1349 2°8072	- '0062 + '0110		- 7 29 11'41 +29 41 53'28	+ '20 + '31	18.757 18.752	·152 ·135	- ·021 - ·035	17 19:20	9°55 8·84 : 8·80	5863 5865
1216 1217	4'0 3'6	47 Pegasiλ Gruis	22 41 42·842 42 31·093	— <sup>•</sup> 036 — <sup>•</sup> 096	+ 2.8861 3.6476	+ .0084 0517	+.0042	+23 2 21.90 -51 50 34.28	+ '11 + '53	+18.875	+ .133	- ·014 - ·062	21 : 25 19	8.62 : 7.92 8.61	5875 5880
1218 1219	4.2	71 Aquariiτ 48 Pegasiμ	44 17.888 45 10.650	+ .009	3°1803 2°8915	- °0098 + °0092	- '0011 + '0107	-14 7 13.84 -1-24 4 24.30	+ '30	18·927 18·943	·142 ·127	- ·036 - ·045	21 18	8·28 9°07	5884 5885
1220	5.6	Lacaille 9275		022	3.4299	- '0312	+.0022	-39 41 11.06	+ .14	18.978	.152	012	18	9.37	5886
1221	6·4 4·6	Lacaille 9268pr. 22 Piscis Austseq. y		+ '013 + '022	+ 3.9334 3.3461	- •0918 •0242	- '0013 - '0024	-63 43 4.68 -33 24 21.16	+ '42 + '29	+18.959	+ 174	- °043 - °032	16 17:18	9.69 9.27 : 9.13	5858 5893
1223 1224	3.8	73 Aquariiλ Indiρ	47 23°916 47 42°230	003 + .092	3.1320	·0062 ·1447	+.0003	- 8 6 42.13	$- \cdot 32$ - $\cdot 68$	19.085 19.131	·134 ·183	+ .036	20:22 17	8 ·97 : 8 ·88 9 · 30	5895 5898
1225	3'4	76 Aquariiδ	49 20.631	+ .030	3.1880	.0109	0033			19.081	.133		21:23	9.19 : 9.54	5904
1226 1227	7°2 6'1	Gruis $\tau^2$ Piazzi XXII. 250 m.	22 49 26·382 49 59·858	+ ·207 - ·011	+ 3.5157 3.1124	- °0444 °0047	-·0229 +·0012	-49 I 33.35 - 5 31 I4.56	- ·45 + ·03	+19.124	+ '146	+ .020	18 16	9'05 9'29	5906 590.)
1228 1229	1.0 1.0	-24 Piscis Austa 52 Pegasim	52 7°809 54 11°646	- · 207 - ·016	3.3239	- '0211	+.0250	-30 9 9.48	+ 1.34	19.008	·134 •116	- ·166	21:24	8·27 : 5·10 9·40	5916 5922
1229	4.1	Gruisζ	54 58.613		2°9991 3°5667	+ .0038	+.0017		+ '39 + '04	19°185 19°241	•137	005	19:20 24	8.18	5925 5926
1231	5.7	Lacaille 9337		- '022	+ 4.0190	- 1256	+.0024	-69 21 38.85	73	+19.404		+ .080	18:19	9.12 : 9.16	5937
1232 1233	4.5 var.	4 Pisciumβ 53 Pegasiβ		- '007 - '141	3.0528		1	+ 3 16 53.61 + 27 32 26.59		19·328 19·472	·109	- ·008 + ·133	16 16	9°41 9°76	5939 5940
1234	2.6	54 Pegasia	22 59 46.807	041	2.9855	+ .0028	+.0040	+14 40 1.50	+ .46	19.314	.102	045	17:16	10.10	5944
1235	1.2	Gruispr. θ		+ .040	3.3960	0323	-*0043		+ '32	19.328	•117	034	18:19	9'39:9'41	5949
1236 1237	4·7 4·8	86 Aquarii	23 I 18.664 I 58.020	044	+ 3.2301	- °0158 + °0031	+ 0050	-24 17 0.44 + 8 52 9.31	+ .13	+19.393	+ '111	°000 - °014	18:19 16	8·77 : 8·72 9·36	5950 5952
1238	5.7	5 PisciumA	, 3 33.713	- '085	3.0724		+.0001	+ 1 35 1.09	- 1.03	19.22	1	+ .110	16	9*34	5959
1239 1240	3.8 4.0	88 Aquarii	4 7°000 4 42°126	- °028	3°2041 3°4138	·0138 - ·0376	+.0033	-21 42 54 45 -45 47 18 22	- · · 30 + · 36	19°491 19°426	.102 .102	+ <sup>•</sup> 037 - <sup>•</sup> 040	35 : 29 19	8'57 : 8'19 9'02	5960 5965
1241	5'3	59 Pegasi	23 6 41.235	+ .010	+ 3.0270	+ .0030	0011	+ 8 10 37 . 24	+ .00	+19.201	+ .094	- •006	19	9.29	5973
1242 1243	4°4 6°1	90 Aquariiφ Lacaille 9407φ	9 8.673 9 25.993	- ·017 - ·074	3·1081 3·3416	- ·0043	+.0001	-63519.19 -413850.63	+ 1.74 + .98	19·364 19·441	•092 •099	- ·191 - ·120	16:17 21	9.28 : 9.13 8.15	5978 5979
1244	4.4	91 Aquarii	10 39.421	237	3.1456	·0060	+ 0248	- 9 37 57.68	+ '12	19.221	.090	013	16:17	9.57 : 9.56	5981
1245	5.8	Lacaille 9412	10 57.276	- '231	3.6435	.0781	+.0248	-62 32 46.53	+ •33	19.555	.102	035	18	9'31	5983
1246 1247	4°1 3°8	Toucani $\gamma$ 6 Piscium $\gamma$	23 11 35·679 11 59·382	+ ·046 - ·488	+ 3.5299 3.1093	- ·0636 + ·0007	- · 0048 + · 0502	-58 47 1.34 + 2 44 9.15	18	+19.682	+ ·099 ·088	+ .081	17 17	9°50 9°72	5985 5988
1248	5.7	Octantis $\tau$ ,	13 10.15	17	10.929	- 5.236	+.012	-88 1 52.85	12	19.643		+ .013	56:162		5994
-	$\begin{array}{c ccccccccccccccccccccccccccccccccccc$														
	1221. 6.5, 9.5       1":2       20°       1900.8.         1222. 4.6, 8.8       4":1       268°       1900.8.         1227. 6.3, 8.3       0":8       325°       1899.8.         1229. 6.2, 7.7; close binary.       1230. 1., 2.2-2.7; P, irregular.														

.

.


FROM OBSERVATIONS IN THE YEARS 1905-11.

No.	Mag.	Name.	Mean <b>R.A.</b> 1900'0.	$\mu_{\alpha}\Delta E.$	Annual Variation 1900°0.	Sec. Var. 1900'0.	Proper Motion.	Mean Dec. 1900'0.	$\mu_{\delta}\Delta E.$	Annual Variation 1900'0.	Sec. Var. 1900'0.	Proper Motion.	No. of Obs.	Epoch 1900+.	Boss No.
1251	4.6	62 Pegasiτ	hms 231541.217	8 - °020	s + 2.9643	s + •0111	s + '0020	+23 11 34.71	+ "20	+19.652	+ "075	- "ô21	16	9.75	6005
1252	6.0	Lacaille 9448		+ .019	3.2034	0172	0021	-27 32 3.69	+ .41	19.631	.081	046	18	8.93	6007
1253	4.5	98 Aquariib1	17 43.100	+ .074	3.1260	- •0122	0087		+ .77	19.014	•076	093	23:22	8.45 : 8.33	6012
1254	4.6	68 Pegasi	20 23.354	- •115		+ .0114	+ .0137	+22 51 13.26	23	19.776		+ .028	21	8.36	6024
1255	5.2	Gruis	21 0.932	015	3.3220	0428	+.0014	-53 16 29.23	- 1.00	19.875	.075	+ •117	19	8.57	6027
1256	6.6	Lacaille 9476			+ 3.3467	0428	+.0050	-50 42 27.49	+ .13	+19.753	+ .073	013	16:17	9'46 : 9'30	6030
1257	5.0	8 Piseium	21 48.449	059			+.0027	+ 0 42 28.14	+ .90	19.679	•066	090	17:16	10.27 : 9.96	6031
1258	4°5 5°8	10 Pisciumθ Lacaille 9483	22 53.653 23 13.790	+ .086	3.0415 3.2080	+ ·0028 - ·0768	-·0088 +·0044	+ 5 49 46.24 -63 39 40.09	+ '42	19.742	•063 •073	- °043 •000	16 20:22	9.80 8.48 : 8.33	6037 6039
1259 1260	4.7	70 Pegasi	23 13 790	034			+.0038	+12 12 32.34	- '24	19.828	1	+ .027	19:21	8.84 : 8.81	6040
	6.0						+.0032	-77 56 15.16		+19.833					
1261 1262	4.6	Lacaille 9494 Sculptoris	23 26 52·244 27 36·708	- ·025 - ·066	+ 3.9765	- ·2200 ·0258	+.0032	-38 22 16.36	+ .04	+19.033	+ .075	- ·005 + ·011	44:45	7 <sup>.</sup> 89 : 7 <sup>.</sup> 91 8 <sup>.</sup> 58 : 8 <sup>.</sup> 53	6052 6054
1263	4.7	101 Aquarii	28 2.686	+ .005	3'1449	0121	- '0005	-21 28 1.93	10	19.869		+ '017	17	9'42	6057
1264†	5.2	72 Pegasim	28 59.473	038		+ .0166	+.0040	+30 46 24.09	+ .11	19.852	.050	012	17:18	9.53 : 9.38	6059
1265	6.5	14 Piscium	29 0.605	069	3.0821	0008	-·007 I	- 1 47 59.55	+ .09	19.855	.052	009	16	9.67	6060
1266	4.8	Phœnicis	23 29 41.943	010	+ 3.2399	0307	+.0023	-43 10 4.86	+ .07	+19.863	+ .054	000	19:20	8.38 : 8.32	6062
1267	7.9	Lacaille 9464	29 47 .08		6.371	1.750		-86 57 6.05		19.873	114		30:54	11.50 : 9.62	
1268	6.7	Mayer 1003		+ .002	3.0960	•0040	0006	- 8 I 4.40	19	19.900	.020	+ .020	17	9.37	6065
1269	4.9	Lacaille 9535		- '049	3.2447	0339	+.0062	-46 2 44.42	+ .19	19.878	•048	- '024	26	7.86	6068
1270	4.5	17 Piscium	34 48.607	208	3.0841	+ .0035	+.0248	+ 5 5 0.03	+ 3.62	19.487	•041	- •439	22:19	8.38 : 8.24	6077
1271	5.2	Sculptoris		+ .072	+ 3.1228	- '0196	0082	-32 37 34.01	+ .33	+19.893	+ .041	038	19	8.79	6079
1272	4.7	18 Pisciumλ	36 56.221	+ .086	3.0603	+ .0015	0095	+ 1 13 45.50	+ 1.34	19.801	.036	- '144	16	9.33	6084
1273†	4.7	105 Aquariipr. ω <sup>2</sup> Lacaille 9566	37 32.296	053	3'1136	0077	+.0000	-15 5 52.86 -71 2 48.94	+ .22	19.891	.036	060	28:25	8.88:8.68	6087
1274	6·3 5·4	106 Aquarii	38 42·772 39 0·946	- '428	3.4818	·1092 ·0098	+.0484	-18 49 55 27	57	20.025	·040 ·033	+ .065	20:21	8.84 : 8.72 9.42	6093 6095
				- 019			1 0020					000			0095
1276	7.9 5.6	Lacaille 9563			+ 4.204	529		-84 25 5.00		+19.978	+ '042		30:55	11.21:9.82	
1277 1278	5.4	19 Pisciumσ		+ .032	3.0634	$+ \cdot 0023$ $- \cdot 0387$	- · 0034	+ 2 55 55.05	+ .10	19.960	·028	- '020 - '012	16 21:23	9°33 8.65 : 8.42	6102 6103
1279†	4.7	Sculptoris seq. 8	43 43.176	066	3.1322	•0160	-+ · 0080	-28 41 0.28	+ .83	19.895	•024	- '101	22	8.23	6110
1280	7.9	Lacaille 9596	46 10.18		4.372	887		-86 27 8.25		20'010	.030		35:59	11.24 : 10.02	
1281	5.2	81 Pegasi	22 47 22:080	+ .010	+ 3.0466	+ .0110	0011	+18 33 53.85	+ .38	+19.973	+ .016	044	21	. 8.74	6127
1282	6.5	25 Piseium		007		+ .0020		+ 1 32 4.18	+ .06	20.013	.015	006	16	9.33	6133
1283	6.5	Lacaille 9633		- '028			1	-24 47 7.49	+ .01	20.019		001	21:23	8.64 : 8.67	6134
	5.9	Octantis $\gamma^2$	52 4		3.4164	2848	0184	-82 43 33.31	+ .15	20.019	.009	019	8	7.42	6146
1285	4.8	84 Pegasi	52 39.688	+ .029	3.0497	+ .0149	0031	+24 35 8.03	+ .35	20.000	•006	032	16	9.38	6150
1286†	5.2	27 Pisciumseq.	23 53 33.211	+ .036	+ 3.0712	- '0007	0038	- 4 6 39.67	+ .64	+19.972		067	16	9.49	6153
1287	5.5	Phœnicis	53 45.021	044	3.1263	0400	+.0021	-53 18 15.27	- '46	20.093		+ .054	21:22	8.53 : 8.20	6154
1288	4'I	28 Piseiumω	54 10.646	- •091	3.0282			+ 6 18 33.85		19'931	•003	109	21:26	9.05 : 8.76	6156
1289	4.6	Toucaniε Octantisθ	54 43 395	063	3.1486			-66 8 0.40		20.018	+ .002	- '024	17	9.35	6160
1290	4.9		56 27 394	+ .164	3.1449		- '0215	-77 37 5.02			001	126	42:45	7*63 : 7*62	6165
1291	4.6	30 Piscium			+ 3.0772		+.0027	- 6 34 11.71	+ .32	+20.011	002	034	16	9.42	6171
1292	4.6	2 Ceti Lacaille 9710	58 37.089	013	3.0762				+ .08	20°038 20°032	•006 •008	- ·008	15:17	9.66 : 9.55	6179
1293	5.8	Lacame 9710	59 37.149	024	3.0822	-0908	+ .0062	-71 59 36.34	+ .15	20.032	003	- 015	23:24	8.54 : 8.13	6185
			1	Seres H			1				1		TRACE	1.1.1	

1264.6.0,6.0;very close binary.1273.4.7,10.7 $5^{\prime\prime}$ '3 $84^{\circ}$ 1898.7.1279.4.7,11.0 $3^{\prime\prime}$ '3230°1899.7.1286.5.2,11 $1^{\prime\prime}$ .8270°1899.8.

.


đ





## UNIVERSITY OF CALIFORNIA LIBRARY BERKELEY

Return to desk from which borrowed. This book is DUE on the last date stamped below.



Y.H 02473

